

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/435, 14/705, C12N 15/12, 15/63, 15/70, 15/74, 15/79, 15/81, A61K 38/17</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/37675</b>
			<b>(43) International Publication Date:</b> 29 July 1999 (29.07.99)
<b>(21) International Application Number:</b> PCT/US99/01418 <b>(22) International Filing Date:</b> 22 January 1999 (22.01.99) <b>(30) Priority Data:</b> 60/072,151 22 January 1998 (22.01.98) US <b>(71) Applicant (for all designated States except US):</b> THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 12th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> JULIUS, David, J. [US/US]; 50 Clifford Terrace, San Francisco, CA 94117 (US). CATERINA, Michael [US/US]; Apartment 7, 110 Carlotta Circle, Mill Valley, CA 94941 (US). BRAKE, Anthony, J. [US/US]; 2115 Los Angeles Avenue, Berkeley, CA 94707 (US). <b>(74) Agent:</b> FRANCIS, Carol, L.; Bozicevic, Field & Francis LLP, Suite 200, 285 Hamilton Avenue, Palo Alto, CA 94301 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>	
<b>(54) Title:</b> NUCLEIC ACID SEQUENCES ENCODING CAPSAICIN RECEPTORS			
<b>(57) Abstract</b> <p>The present invention features vanilloid receptor polypeptides and vanilloid receptor-related polypeptides, specifically the capsaicin receptor subtypes VR1 and VR2 (VRRP-1), as well as the encoding polynucleotide sequences. In related aspects the invention features expression vectors and host cells comprising such polynucleotides. In other related aspects, the invention features transgenic animals having altered capsaicin receptor expression, due to, for example, the presence of an exogenous wild-type or modified capsaicin receptor-encoding polynucleotide sequence. The present invention also relates to antibodies that bind specifically to a capsaicin receptor polypeptide, and methods for producing these polypeptides. Further, the invention provides methods for using capsaicin receptor, including methods for screening candidate agents for activity as agonists or antagonists of capsaicin receptor activity, as well as assays to determine the amount of a capsaicin receptor-activating agent in a sample. In other related aspects, the invention provides methods for the use of the capsaicin receptor for the diagnosis and treatment of human disease and painful syndromes.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**NUCLEIC ACID SEQUENCES ENCODING CAPSAICIN RECEPTOR****FIELD OF THE INVENTION**

5           The present invention relates to nucleic acid and amino acid sequences encoding a receptor for vanilloid compound and polypeptides related to such vanilloid compound receptors, and to the use of these sequences in the diagnosis, study, and treatment of disease.

**BACKGROUND OF THE INVENTION**

10           Pain is initiated when the peripheral terminals of a particular group of sensory neurons, called nociceptors, are activated by noxious chemical, mechanical, or thermal stimuli. These neurons, whose cell bodies are located in various sensory ganglia, transmit information regarding tissue damage to pain processing centers in the spinal cord and brain (Fields Pain (McGraw-Hill, New York, 1987)). Nociceptors are characterized, in part, by their sensitivity to capsaicin, a natural product of capsicum peppers that is  
15           the active ingredient of many "hot" and spicy foods. In mammals, exposure of nociceptor terminals to capsaicin leads initially to the perception of pain and the local release of neurotransmitters. With prolonged exposure, these terminals become insensitive to capsaicin, as well as to other noxious stimuli (Szolcsanyi in Capsaicin in the Study of Pain (ed. Wood) pgs. 255-272 (Academic Press, London, 1993)). This latter phenomenon of nociceptor desensitization underlies the seemingly paradoxical use of capsaicin as an  
20           analgesic agent in the treatment of painful disorders ranging from viral and diabetic neuropathies to rheumatoid arthritis (Campbell in Capsaicin and the Study of Pain (ed. Wood) pgs. 255-272 (Academic Press, London, 1993); Szallasi et al. 1996 Pain 68:195-208). While some of this decreased sensitivity to noxious stimuli may reflect reversible changes in the nociceptor, such as depletion of inflammatory mediators, the long-term loss of responsiveness can be explained by death of the nociceptor or destruction  
25           of its peripheral terminals following capsaicin exposure (Jancso et al. 1977 Nature 270:741-743; Szolcsanyi, *supra*).

            Responsivity to capsaicin has been used to define sensory afferent fibers that transmit signals in response to noxious stimuli (chemical, thermal, and mechanical stimuli); however, the precise mechanism of action has remained unclear. Electrophysiological (Bevan et al. 1990 Trends Pharmacol. Sci 11:330-  
30           333; Oh et al. 1996 J. Neuroscience 16:1659-1667) and biochemical (Wood et al. 1988 J. Neuroscience 8:3208-3220) studies have clearly shown that capsaicin excites nociceptors by increasing plasma membrane conductance through formation or activation of nonselective cation channels. While the hydrophobic nature of capsaicin has made it difficult to rule out the possibility that its actions are mediated by direct perturbation of membrane lipids (Feigin et al. 1995 Neuroreport 6:2134-2136), it has been  
35           generally accepted that this compounds acts at a specific receptor site on or within sensory neurons due to observations that capsaicin derivatives show structure-function relationships and evoke dose-dependent responses (Szolcsanyi et al. 1975 Drug. Res. 25:1877-1881; Szolcsanyi et al. 1976 Drug Res. 26:33-37)).  
            The development of capsazepine, a competitive capsaicin antagonist (Bevan et al. 1992 Br. J. Pharmacol.

107:544-552) and the discovery of resiniferatoxin, an ultrapotent capsaicin analogue from *Euphorbia* plants that mimics the cellular actions of capsaicin (deVries et al. 1989 Life Sci. 44:711-715; Szallasi et al. 1989 Neuroscience 30:515-520) further suggest that the capsaicin mediates its effects through a receptor. The nanomolar potency of resiniferatoxin has facilitated its use as a high affinity radioligand to visualize  
5 saturable, capsaicin- and capsazepine-sensitive binding sites on nociceptors (Szallasi 1994 Gen. Pharmac. 25:223-243). Because a vanilloid moiety constitutes an essential structural component of capsaicin and resiniferatoxin, the proposed site of action of these compounds has been more generally referred to as the vanilloid receptor (Szallasi 1994 *supra*). The action of capsaicin, resiniferatoxin, and the antagonist capsazepine have been well characterized physiologically using primary neuronal cultures (see, e.g.,  
10 Szolcsanyi, "Actions of Capsaicin on Sensory Receptors," Bevan et al. "Cellular Mechanisms of the Action of Capsaicin," and James et al. "The Capsaicin Receptor," all in Capsaicin in the Study of Pain, 1993 Academic Press Limited, pgs. 1-26, 27-44, and 83-104, respectively; Bevan et al. 1990, *supra*).

The analgesic properties of capsaicin and capsaicinoids are of much interest for their uses in the treatment of pain and inflammation associated with a variety of disorders (see, e.g, Fusco et al. 1997 Drugs  
15 53:909-914; Towheed et al. 1997 Semin. Arthritis Rheum 26:755-770; Appendino et al. 1997 Life Sci 60:681-696 (describing activities and application of resiniferatoxin); Campbell et al. "Clinical Applications of Capsaicin and Its Analogues" in Capsaicin in the Study of Pain 1993, Academic Press pgs. 255-272).

Although capsaicin and capsaicin related compounds can evoke the sensation of pain, cause hyperalgesia, activate autonomic reflexes (e.g., elicit changes in blood pressure), and cause release of  
20 peptides and other putative transmitters from nerve terminals (e.g., to induce bronchoconstriction and inflammation), prolonged exposure of sensory neurons to these compounds leads to desensitization of the neurons to multiple modalities of noxious sensory stimuli without compromising normal mechanical sensitivity or motor function, and without apparent central nervous system depression. It is this final phenomena that makes capsaicins and related compounds of great interest and potential therapeutic  
25 value.

Despite the intense interest in capsaicin and related compounds and their effects upon sensory afferent, the receptor(s) through which these compounds mediate their effects have eluded isolation and molecular characterization. Thus, the development of elegant systems for screening or characterizing new capsaicin receptor-binding compounds, or for identifying endogenous, tissue-derived mediators of pain  
30 and/or inflammation, have been severely hampered. To date the only means of assessing the activity of compounds as capsaicin receptor agonists or antagonists has been to examine their effects on sensory neurons in culture or in intact animals. The present invention solves this problem.

#### SUMMARY

35 The present invention features vanilloid receptor polypeptides and vanilloid receptor-related polypeptides, specifically the capsaicin receptor and capsaicin receptor-related polypeptides, as well as nucleotide sequences encoding capsaicin receptor and capsaicin receptor-related polypeptides. In related aspects the invention features expression vectors and host cells comprising polynucleotides that encode



capsaicin receptor or capsaicin receptor-related polypeptide. In other related aspects, the invention features transgenic animals having altered capsaicin receptor expression, due to, for example, the presence of an exogenous wild-type or modified capsaicin receptor-encoding polynucleotide sequence.

The present invention also relates to antibodies that bind specifically to a capsaicin receptor polypeptide and/or capsaicin receptor-related polypeptide, and methods for producing capsaicin receptor and capsaicin receptor-related polypeptides.

In one aspect the invention features a method for identifying compounds that bind a capsaicin receptor polypeptide, preferably a compound that binds a capsaicin receptor polypeptide and affects a cellular response associated with capsaicin receptor biological activity (e.g., intracellular calcium flux).

In another aspect the invention features a method for detecting a vanilloid compound in a sample, where the vanilloid compound has activity in binding a capsaicin receptor polypeptide, by contacting a sample suspected of containing a vanilloid compound with a cell (e.g., an oocyte (e.g., an amphibian oocyte) or a mammalian cell) expressing a capsaicin receptor polypeptide and detecting an alteration of a cellular response associated with capsaicin receptor activity in the capsaicin receptor-expressing host cell. Preferably, the cellular response associated with capsaicin receptor activity is an alteration of intracellular calcium levels in the capsaicin receptor-expressing host cell. The method can be used to detect vanilloid compounds in samples derived from natural products (e.g., natural product extracts) or can be used to screen candidate compounds for use as analgesics (e.g., vanilloid analogs, therapeutic antibodies, antisense oligonucleotides, capsaicin receptor-encoding nucleotides for replacement gene therapy), flavor-enhancing agents, etc.

Yet another aspect of the invention relates to use of capsaicin receptor polypeptides and specific antibodies thereto for the diagnosis and treatment of human disease and painful syndromes.

In another aspect the invention features transgenic, non-human mammals expressing a capsaicin receptor-encoding transgene, and use of such transgenic mammals for use in screening of candidate capsaicin receptor agonist and antagonist compounds.

A primary object of the invention is to provide isolated polynucleotides for use in expression of capsaicin receptor and capsaicin receptor-related polypeptides (e.g. in a recombinant host cell or in a target cell as part of organochemotherapy) and for use in, for example, identification of capsaicin receptor-binding compounds (especially those compounds that affect capsaicin receptor-mediated activity).

The invention will now be described in further detail.

### BRIEF DESCRIPTION OF THE FIGURES

Fig. 1A shows the putative domains present in the capsaicin receptor amino acid sequence. Open boxes delineate ankyrin repeat domains; right-hatched boxes indicate predicted transmembrane domains; and the left-hatched box indicates a possible pore-loop region. Bullets denote predicted protein kinase A phosphorylation sites.

Fig. 1B shows the predicted membrane topology and domain structure of the capsaicin receptor. Open circles labeled "A" denote ankyrin domains; cylinders denote transmembrane domains; and the

double-lined area indicates a possible pore-loop region. "i" and "o" denote the inner and outer membrane leaflets, respectively.

Fig. 1C shows the alignment of the capsaicin receptor VR1 with related sequences. Identical residues are in cross-hatched boxes and conservative substitutions are in singly-lined boxes.

5 Fig. 2 is a current trace of whole cell voltage clamp analysis of capsaicin receptor-expressing HEK293 cells.

Fig. 3 is a plot of the voltage steps (400ms) from -100 mV to +40mV for the data presented in Figure 3.

Fig. 4 is a graph illustrating the current-voltage relationship of the data from Figure 4.

10 Fig. 5 is a graph of the voltage generated across membranes of recombinant capsaicin receptor-expressing cells when exposed to capsaicin and tested under conditions of varying ionic compositions. a = NaCl; b = KCl; c = CsCl; d = MgCl<sub>2</sub>; e = CaCl<sub>2</sub>.

Fig. 6A through Fig. 6F are graphs showing the effects of extracellular calcium upon capsaicin-induced current in whole-cell voltage clamp experiments.

15 Fig. 7 is graph illustrating the single channel behavior of capsaicin-induced current in capsaicin receptor-expressing HEK293 cells using outside-out (O/O) and inside-out (I/O) patches.

Fig. 8 is a graph showing the current-voltage relationship of the data obtained in Fig. 7.

Fig. 9A is a graph showing the effects of capsaicin and resiniferatoxin upon current in whole-cell voltage clamp experiments in *Xenopus* oocytes expressing the capsaicin receptor. Bars denote duration of agonist application. Membrane currents were recorded in the whole cell voltage clamp configuration ( $V_{\text{hold}} = -40\text{mV}$ ).

20 Fig. 9B is a graph showing the concentration-response curve for capsaicin (squares) and resiniferatoxin (open circles) in VR1-expressing oocytes. In Fig. 9B, the membrane currents were normalized in each oocyte to a response obtained with 1 $\mu\text{M}$  capsaicin and expressed as a percent of maximal response to capsaicin. Each point represents mean values ( $\pm$  s.e.m.) from 5 independent oocytes.

Fig. 10A is a graph showing the effects of capsazepine upon capsaicin-induced current in whole cell voltage clamp experiments. Slash marks represent wash out periods of 2 and 3 min, respectively ( $n = 3$ ). cap = capsaicin; cpz = capsazepine; RR = ruthenium red. Each point represents 4 independent oocytes. Current response were normalized to that elicited by capsaicin in each oocyte (0.6  $\mu\text{M}$ , open diamond). Slash marks denote 2 and 12 minute wash out periods, respectively ( $n=3$ ).

30 Fig. 10B is a graph showing the capsazepine inhibition curve of capsaicin response in whole cell voltage clamp studies.

Fig. 11 is a histogram with corresponding current traces reflecting the relative capsaicin content of several different hot peppers.

35 Fig. 12 is a graph showing induction of cell death in HEK293 cells transiently transfected with capsaicin receptor-encoding cDNA. Open bars = cells exposed to carrier alone (ethanol); filled bars = cells exposed to capsaicin; pcDNA3 = control cells without capsaicin receptor-encoding DNA; VR1 (1:50) = cells transiently transfected with capsaicin receptor-s encoding cDNA diluted 1:50 with control pcDNA3; and

VR1 = cells transiently transfected with capsaicin receptor-encoding cDNA alone. Asterisks indicate a significant difference from ethanol-treated cells.

Fig. 13 is a current trace showing the effect of hydrogen ions upon capsaicin receptor activity in oocytes expressing capsaicin receptor. cap on = time of capsaicin introduction; cap off = time of capsaicin wash out. The pH of the bath solution was changed during the experiment as indicated by the horizontal bars.

Fig. 14 is a graph showing a summary of the current response obtained from nine independent capsaicin receptor-expressing oocytes. The open portion of each bar indicates peak current evoked by capsaicin at pH 7.6, while the cross-hatched portion represents the additional current evoked by changing the bath solution to pH 6.3.

Fig. 15A is a current trace showing the effects of heat and capsaicin upon capsaicin receptor activity in capsaicin receptor-expressing HEK293 cells as determined by whole patch clamp analysis.

Fig. 15B is a graph showing the current-voltage relationship of the data obtained in Fig. 15A.

Fig. 16 is a graph showing activation of capsaicin receptor in capsaicin receptor-expressing *Xenopus* oocytes by noxious, but not innocuous, heat. The asterisk indicates a significant difference from control water-injected oocytes.

Fig. 17 provides representative current traces of the effects of capsaicin, heat, and heat plus ruthenium red (RR) upon capsaicin receptor-expressing *Xenopus* oocytes and control water-injected oocytes.

Fig. 18 is a schematic illustrating the relationship between rat VR1, rat VRRP-1, and the human EST sequence AA321554.

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions

"Polynucleotide" as used herein refers to an oligonucleotide, nucleotide, and fragments or portions thereof, as well as to peptide nucleic acids (PNA), fragments, portions or antisense molecules thereof, and to DNA or RNA of genomic or synthetic origin which can be single- or double-stranded, and represent the sense or antisense strand. Where "polynucleotide" is used to refer to a specific polynucleotide sequence (e.g. a capsaicin receptor-encoding polynucleotide or a capsaicin receptor-related polypeptide-encoding polynucleotide), "polynucleotide" is meant to encompass polynucleotides that encode a polypeptide that is functionally equivalent to the recited polypeptide, e.g., polynucleotides that are degenerate variants, or polynucleotides that encode biologically active variants or fragments of the recited polypeptide. Similarly, "polypeptide" as used herein refers to an oligopeptide, peptide, or protein. Where "polypeptide" is recited herein to refer to an amino acid sequence of a naturally-occurring protein molecule, "polypeptide" and like terms are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule. By "antisense polynucleotide" is mean a polynucleotide having a nucleotide sequence complementary to a given polynucleotide sequence (e.g. a polynucleotide sequence encoding a capsaicin receptor) including polynucleotide sequences associated with the

transcription or translation of the given polynucleotide sequence (e.g., a promoter of a polynucleotide encoding capsaicin receptor), where the antisense polynucleotide is capable of hybridizing to a capsaicin receptor polynucleotide sequence. Of particular interest are antisense polynucleotides capable of inhibiting transcription and/or translation of a capsaicin receptor-encoding or capsaicin receptor-related polypeptide-  
5 encoding polynucleotide either in vitro or in vivo.

"Peptide nucleic acid" as used herein refers to a molecule which comprises an oligomer to which an amino acid residue, such as lysine, and an amino group have been added. These small molecules, also designated anti-gene agents, stop transcript elongation by binding to their complementary (template) strand of nucleic acid (Nielsen et al 1993 Anticancer Drug Des 8:53-63).

10 As used herein, "capsaicin receptor" or "capsaicin receptor polypeptide" means a recombinant or nonrecombinant polypeptide having an amino acid sequence of i) a native capsaicin receptor polypeptide, ii) a biologically active fragment of a capsaicin receptor polypeptide, iii) biologically active polypeptide analogs of a capsaicin receptor polypeptide, or iv) a biologically active variant of a capsaicin receptor polypeptide. Capsaicin receptor polypeptides of the invention can be obtained from any species,  
15 particularly mammalian, including human, rodentia (e.g., murine or rat), bovine, ovine, porcine, murine, or equine, preferably rat or human, from any source whether natural, synthetic, semi-synthetic or recombinant. The term "capsaicin receptor" as used herein encompasses the vanilloid receptor subtype 1 (VR1) described in detail herein, but is not meant to be limited to VR1, and particularly may be generically used to refer to the receptor subtypes VR1 and VR2.

20 As used herein, "capsaicin receptor-related polypeptide" or "vanilloid-like receptor (VLR) polypeptide" means a recombinant or nonrecombinant polypeptide having an amino acid sequence of i) a native capsaicin receptor-related polypeptide, ii) a biologically active fragment of a capsaicin receptor-related polypeptide, iii) biologically active polypeptide analogs of a capsaicin receptor-related polypeptide, or iv) a biologically active variant of a capsaicin receptor-related polypeptide, herein referred to as "VRRP-  
25 1", "VLR1," or "VR2". Capsaicin receptor polypeptides of the invention can be obtained from any species, particularly a mammalian species, including human, rodentia (e.g., murine or rat), bovine, ovine, porcine, murine, or equine, preferably rat or human, from any source whether natural, synthetic, semi-synthetic or recombinant. The term "capsaicin receptor-related polypeptide" as used herein also encompasses any polypeptide having at least about 40% identity, preferably at least about 45% identity, more preferably at  
30 least about 49% identity to an amino acid sequence of a capsaicin receptor polypeptide of the same species (e.g., rat or human capsaicin receptor polypeptide). The term "capsaicin receptor-related polypeptide-encoding sequence" also encompasses a nucleotide sequence having at least about 50% identity, preferably at least about 55% identity, more preferably at least about 59% identity to a nucleotide sequence of a capsaicin receptor polypeptide of the same species. In one embodiment, the capsaicin  
35 receptor-related polypeptide interacts with capsaicin receptor. "Capsaicin receptor-related polypeptide" as used herein encompasses the vanilloid receptor-related polypeptide 1 (VRRP-1) described in detail herein, but is not meant to be limited to VRRP-1.

As used herein, "antigenic amino acid sequence" means an amino acid sequence that, either alone or in association with a carrier molecule, can elicit an antibody response in a mammal.

5 A "variant" of a capsaicin receptor or capsaicin receptor-related polypeptide is defined as an amino acid sequence that is altered by one or more amino acids. The variant can have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. More rarely, a variant can have "nonconservative" changes, e.g., replacement of a glycine with a tryptophan. Similar minor variations can also include amino acid deletions or insertions, or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted without abolishing biological or immunological activity can be found using computer programs well known  
10 in the art, for example, DNASTar software.

A "deletion" is defined as a change in either amino acid or nucleotide sequence in which one or more amino acid or nucleotide residues, respectively, are absent as compared to an amino acid sequence or nucleotide sequence of a naturally occurring capsaicin receptor or capsaicin receptor-related polypeptide.

15 An "insertion" or "addition" is that change in an amino acid or nucleotide sequence which has resulted in the addition of one or more amino acid or nucleotide residues, respectively, as compared to an amino acid sequence or nucleotide sequence of a naturally occurring capsaicin receptor or capsaicin receptor-related polypeptide.

20 A "substitution" results from the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively as compared to an amino acid sequence or nucleotide sequence of a naturally occurring capsaicin receptor or capsaicin receptor-related polypeptide.

The term "biologically active" refers to capsaicin receptor polypeptide or capsaicin receptor-related polypeptide having structural, regulatory, or biochemical functions of a naturally occurring capsaicin receptor polypeptide or capsaicin receptor-related polypeptide, respectively. Likewise, "immunologically  
25 active" defines the capability of the natural, recombinant or synthetic capsaicin receptor (or capsaicin receptor-related polypeptide), or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "derivative" as used herein refers to the chemical modification of a nucleic acid encoding a capsaicin receptor or a capsaicin receptor-related polypeptide. Illustrative of such modifications would  
30 be replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would encode a polypeptide which retains essential biological characteristics of a natural capsaicin receptor or capsaicin receptor-related polypeptide.

As used herein the term "isolated" is meant to describe a compound of interest (e.g., either a polynucleotide or a polypeptide) that is in an environment different from that in which the compound  
35 naturally occurs. "Isolated" is meant to include compounds that are within samples that are substantially enriched for the compound of interest and/or in which the compound of interest is partially or substantially purified.

As used herein, the term "substantially purified" refers to a compound (e.g., either a polynucleotide or a polypeptide) that is removed from its natural environment and is at least 60% free, preferably 75% free, and most preferably 90% free from other components with which it is naturally associated.

5 "Stringency" typically occurs in a range from about  $T_m - 5^\circ\text{C}$  ( $5^\circ\text{C}$  below the  $T_m$  of the probe) to about  $20^\circ\text{C}$  to  $25^\circ\text{C}$  below  $T_m$ . As will be understood by those of skill in the art, a stringency hybridization can be used to identify or detect identical polynucleotide sequences or to identify or detect similar or related polynucleotide sequences.

10 The term "hybridization" as used herein shall include "any process by which a strand of nucleic acid joins with a complementary strand through base pairing" (Coombs 1994 Dictionary of Biotechnology, Stockton Press, New York NY). Amplification as carried out in the polymerase chain reaction technologies is described in Dieffenbach et al. 1995, PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview NY.

15 "Sequence identity" as used herein generally refers to a sequence identity of nucleotide or amino acid sequence, where the sequence identity is generally at least about 65%, preferably at least about 75%, more preferably at least about 85%, and can be greater than at least about 90% or more as determined by the Smith-Waterman homology search algorithm as implemented in MPSRCH program (Oxford Molecular). For the purposes of this invention, a preferred method of calculating percent identity is the Smith-Waterman algorithm, using the following. Global DNA sequence identity must be greater than 65% as determined by the Smith-Waterman homology search algorithm as implemented in MPSRCH program  
20 (Oxford Molecular) using an affine gap search with the following search parameters: gap open penalty, 12; and gap extension penalty, 1.

The term "transgene" is used herein to describe genetic material which has been or is about to be artificially inserted into the genome of a mammalian, particularly a mammalian cell of a living animal.

25 By "transgenic animal" is meant a non-human animal, usually a mammal, having a non-endogenous (*i.e.*, heterologous) nucleic acid sequence present as an extrachromosomal element in a portion of its cells or stably integrated into its germ line DNA (*i.e.*, in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

30 A "knock-out" of a target gene means an alteration in the sequence of the gene that results in a decrease of function of the target gene, preferably such that target gene expression is undetectable or insignificant. For example, a knock-out of a capsaicin receptor gene means that function of the capsaicin receptor has been substantially decreased so that capsaicin receptor expression is not detectable or only present at insignificant levels. "Knock-out" transgenics of the invention can be transgenic animals having a heterozygous or homozygous knock-out of the capsaicin receptor gene or capsaicin receptor-related  
35 polypeptide encoding sequence. "Knock-outs" also include conditional knock-outs, where alteration of the target gene can occur upon, for example, exposure of the animal to a substance that promotes target gene alteration, introduction of an enzyme that promotes recombination at the target gene site (e.g., Cre in the Cre-lox system), or other method for directing the target gene alteration postnatally.

- A "knock-in" of a target gene means an alteration in a host cell genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the target gene, e.g., by introduction of an additional copy of the target gene, or by operatively inserting a regulatory sequence that provides for enhanced expression of an endogenous copy of the target gene. For example, "knock-in" transgenics of the invention can be transgenic animals having a heterozygous or homozygous knock-in of the capsaicin receptor gene. "Knock-ins" also encompass conditional knock-ins.

The major genetic sequences provided herein are as follows:

SEQ ID NO	Sequence
1	Rat VR1 cDNA sequence
2	Rat VR1 amino acid sequence
3	Rat VRRP-1 (VR2) cDNA sequence
4	Rat VRRP-1 (VR2) amino acid sequence
5	Human VRRP-1 consensus sequence, region A
6	Human VRRP-1 consensus sequence, region B
7	Human VRRP-1 consensus sequence, region C
8	EST AA321554 DNA sequence
9	EST AA321554 amino acid sequence
10	mouse VR1 cDNA sequence
11	mouse VR1 cDNA
12	primer
13	primer
14	Rat VR1 amino acid sequence
15	Human T11251 amino acid sequence
16	Caliphora z80230 amino acid sequence
17	Drosophila TRP amino acid sequence
18	Bovine x99792 amino acid sequence
19	C. elegans z72508 amino acid sequence
20	Human VRRP-1 (VR2) DNA sequence
21	Human VRRP-1 (VR2) DNA sequence
22	Human VRRP-1 (VR2) DNA sequence
23	Human VRRP-1 (VR2) amino acid sequence
24	Chicken VR1 cDNA sequence
25	Chicken VR1 amino acid sequence
26	Human VR1 cDNA sequence
27	Human VR1 amino acid sequence
33	Human VR1 cDNA sequence
34	Human VR1 amino acid sequence
35	Human VR2 cDNA sequence

SEQ ID NO	Sequence
36	Human VR2 amino acid sequence

#### Overview of the Invention

The present invention is based upon the identification and isolation of a polynucleotide sequence encoding a capsaicin receptor polypeptide (e.g., the vanilloid receptor subtype 1 (VR1) polypeptide described herein) and a capsaicin receptor-related polypeptide (e.g., the vanilloid receptor-related polypeptide 1 (VRRP-1; or VR2) described herein). The corresponding genetic sequences are provided in the Seqlist, and are listed in the table provided above. Accordingly, the present invention encompasses such polynucleotides encoding capsaicin receptor and/or capsaicin receptor-related polypeptides, as well as the capsaicin receptor and capsaicin receptor-related polypeptides encoded by such polynucleotides.

A capsaicin receptor polypeptide-encoding polynucleotide was first isolated by virtue of the capsaicin receptor polypeptide-encoding sequence to facilitate expression of a functional capsaicin receptor in a cellular assay. In short, the capsaicin receptor polypeptide-encoding polynucleotide, when expressed in a mammalian or amphibian host cell, facilitated an increase in intracellular calcium concentration in the host cell upon exposure to the agonist capsaicin. This work lead to identification and isolation of a polynucleotide sequence encoding a capsaicin receptor referred to herein as a vanilloid receptor subtype 1 (VR1). The capsaicin receptor-encoding VR1 sequence was then used to isolate by PCR amplification a sequences encoding related polypeptides, resulting in isolation and identification of sequences encoding a capsaicin receptor-related polypeptide, of which VRRP-1 (VR2) is exemplary.

The invention also encompasses use of capsaicin receptor and capsaicin receptor-related polypeptide nucleic acid and amino acid sequences in the identification of capsaicin receptor-binding compounds, particularly capsaicin receptor-binding compounds having capsaicin receptor agonist or antagonist activity. The invention further encompasses the use of the polynucleotides disclosed herein to facilitate identification and isolation of polynucleotide and polypeptide sequences having homology to a capsaicin receptor and/or capsaicin receptor-related polypeptide of the invention; as well as the diagnosis, prevention and treatment of disease and/or pain syndromes associated with capsaicin receptor biological activity.

The polynucleotides of the invention can also be used as a molecular probe with which to determine the structure, location, and expression of capsaicin receptor, receptor subtypes, and capsaicin receptor-related polypeptides in mammals (including humans) and to investigate potential associations between disease states or clinical disorders (particularly those involving acute and chronic pain or inflammation) and defects or alterations in receptor structure, expression, or function.

#### Capsaicin Receptor and Capsaicin Receptor-Related Polypeptide Coding Sequences

In accordance with the invention, any nucleic acid sequence that encodes an amino acid sequence of a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide can be used to generate recombinant molecules which express a capsaicin receptor polypeptide or capsaicin receptor-related



polypeptide, respectively. The nucleic acid compositions used in the subject invention may encode all or a part of a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide of the invention as appropriate. Fragments may be obtained of the DNA sequence by chemically synthesizing oligonucleotides in accordance with conventional methods, by restriction enzyme digestion, by PCR  
5 amplification, etc. For the most part, DNA fragments will be of at least about ten contiguous nucleotides, usually at least about 15 nt, more usually at least about 18 nt to about 20 nt, more usually at least about 25 nt to about 50 nt. Such small DNA fragments are useful as primers for PCR, hybridization screening, etc. Larger DNA fragments, *i.e.* greater than 100 nt are useful for production of the encoded polypeptide.

The nucleic acid and deduced amino acid sequences of rat capsaicin receptor (subtype VR1) are  
10 provided as SEQ ID NOS:1 and 2. Nucleic acid and deduced amino acid sequences of a human capsaicin receptor (subtype VR1) are provided as SEQ ID NOS:33 and 34. A nucleotide sequence encoding murine capsaicin receptor subtype VR1 comprises the sequences of SEQ ID NOS:10 and 11. The chicken capsaicin receptor subtype VR1 is provided as SEQ ID NO:24 and 25.

The nucleic acid and deduced amino acid sequence of rat capsaicin receptor-related polypeptide  
15 1 (VRRP-1; or subtype VR2) are provided as SEQ ID NO:3 and 4, respectively. A sequence encoding a human capsaicin receptor-related polypeptide (referred to as human VR2) comprises the nucleotide sequence SEQ ID NOS:35 and 36.

The present invention also encompasses variants of capsaicin receptor and capsaicin receptor-related polypeptides. A preferred variant is one having at least 80% amino acid sequence similarity, more  
20 preferably at least 90% amino acid sequence similarity, still more preferably at least 95% amino acid sequence similarity to an amino acid sequence of a capsaicin receptor, subtype VR1 or VR2.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of degenerate variants of nucleotide sequences encoding capsaicin receptor and capsaicin receptor-related polypeptides, some bearing minimal homology to the nucleotide sequences of  
25 any known and naturally occurring gene, can be produced. The invention contemplates each and every possible variation of nucleotide sequence that can be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring capsaicin receptor or capsaicin receptor-related polypeptide, and all such variations are to be considered as being specifically disclosed herein.

30 Although nucleotide sequences that encode capsaicin receptor polypeptides, and their variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring polypeptides under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding receptors or their derivatives possessing a substantially different codon usage. Codons can be selected to increase the rate at which expression of the polypeptide occurs in a particular  
35 prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding capsaicin receptor, capsaicin receptor-related polypeptide, and their derivatives without altering the

encoded amino acid sequences include the production of RNA transcripts having more desirable properties (e.g., increased half-life) than transcripts produced from the naturally occurring sequence.

Nucleotide sequences encoding a capsaicin receptor polypeptide, capsaicin receptor-related polypeptide, and/or their derivatives can be synthesized entirely by synthetic chemistry, after which the  
5 synthetic gene can be inserted into any of the many available DNA vectors and expression systems using reagents that are well known in the art at the time of the filing of this application. Moreover, synthetic chemistry can be used to introduce mutations into a sequence encoding a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide.

Also included within the scope of the present invention are polynucleotide sequences that are  
10 capable of hybridizing to the nucleotide sequence of any of the provided nucleic acid sequences of capsaicin receptors, subtypes VR1 or VR2. Of particular interest are polynucleotide sequence capable of hybridizing under various conditions of stringency to the coding sequence for capsaicin receptor or capsaicin receptor-related polypeptide (e.g., nucleotides 81-2594 of SEQ ID NO:1, or nucleotides 14-2530 of SEQ ID NO:33), to a region of a capsaicin receptor-encoding sequence or capsaicin receptor-related  
15 polypeptide-encoding sequence that shares homology with other such sequences (e.g., a sequence encoding a contiguous stretch of amino acid residues present in SEQ ID NO:2 (e.g., amino acid residues 636 to 706 of SEQ ID NO:2), a sequence encoding a contiguous stretch of amino acid residues present in SEQ ID NO:33 (e.g., amino acid residues 636 to 706 of SEQ ID NO:33), and other sequences representing areas of homology with other capsaicin receptor-encoding sequences and/or capsaicin  
20 receptor-related polypeptides-encoding sequences, as well as sequences that uniquely identify capsaicin receptor-encoding sequences or capsaicin receptor-related polypeptide-encoding sequences of various species. Of particular interest are capsaicin receptor VR1 or VR2 polynucleotide sequences encoding a human capsaicin receptor polypeptide or human capsaicin receptor-related polypeptide. Hybridization conditions are based on the melting temperature ( $T_m$ ) of the nucleic acid binding complex or probe, as  
25 taught in Berger et al. 1987 Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA incorporated herein by reference, and can be used at a defined stringency.

Altered nucleic acid sequences encoding capsaicin receptor or capsaicin receptor-related polypeptide that can be used in accordance with the invention include deletions, insertions or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent  
30 capsaicin receptor or capsaicin receptor-related polypeptide. The protein can also comprise deletions, insertions or substitutions of amino acid residues that result in a polypeptide that is functionally equivalent to capsaicin receptor or capsaicin receptor-related polypeptide. Deliberate amino acid substitutions can be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues with the proviso that biological activity of capsaicin receptor is retained.  
35 For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine; glycine, alanine; asparagine, glutamine; serine, threonine phenylalanine, and tyrosine.

Alleles of capsaicin receptor, as well as alleles of capsaicin receptor-related polypeptide, are also encompassed by the present invention. As used herein, an "allele" or "allelic sequence" is an alternative form of a capsaicin receptor or capsaicin receptor-related polypeptide. Alleles result from a mutation (*i.e.*, an alteration in the nucleic acid sequence) and generally produce altered mRNAs and/or polypeptides that may or may not have an altered structure or function relative to naturally-occurring capsaicin receptor or capsaicin receptor-related polypeptide. Any given gene may have none, one, or many allelic forms. Common mutational changes that give rise to alleles are generally ascribed to natural deletions, additions or substitutions of amino acids. Each of these types of changes may occur alone or in combination with the other changes, and may occur once or multiple times in a given sequence.

Isolating Capsaicin Receptor-Encoding and Capsaicin Receptor-Related Polypeptide-Encoding Polynucleotides from Other Species

Capsaicin receptor polypeptide-encoding polynucleotides, capsaicin receptor-related polypeptide-encoding polynucleotides, or portions thereof can be used as probes for identifying and cloning homologs of the capsaicin receptor and capsaicin receptor-related polypeptide sequences disclosed herein. Of particular interest are mammalian homologs (especially the human homology of the disclosed rat capsaicin receptor-encoding and capsaicin receptor-related polypeptide-encoding sequences), where the homologs have substantial sequence similarity to one another, *i.e.* at least 40%, usually at least 60%, more usually at least 75%, usually at least 90%, more usually at least 95% sequence similarity. Mammalian homologs of capsaicin receptor may also share a high degree of similarity to the capsaicin receptor disclosed herein in the vicinity of the predicted pore-loop and sixth transmembrane domains. At these regions the capsaicin receptor homologs may exhibit high sequence similarity, *e.g.*, at least about 40% amino acid sequence identity, usually at least about 60% to 75% amino acid sequence identity, with at least about 40% nucleotide sequence similarity, usually at least about 60% to 90% nucleotide sequence similarity.

Sequence similarity is calculated based on a reference sequence, which may be a subset of a larger sequence, such as a conserved motif, coding region, flanking region, *etc.* A reference sequence will usually be at least about 18 nt long, more usually at least about 30 nt long, and may extend to the complete sequence that is being compared. Algorithms for sequence analysis are known in the art, such as BLAST, described in Altschul *et al.* (1990) J Mol Biol 215:403-10. The specificity of the probe, whether it is made from a highly specific region, *e.g.*, 10 unique nucleotides in the 5' regulatory region, or a less specific region, *e.g.*, especially in the 3' region, and the stringency of the hybridization or amplification (maximal, high, intermediate or low) will determine whether the probe identifies only naturally occurring sequences encoding capsaicin receptor, alleles or related sequences.

Where the probes of the invention are used in the detection of related sequences, the probes preferably comprise at least 30%, more preferably at least 50% of the nucleotides from any of the capsaicin receptor polypeptide-encoding sequences or the capsaicin receptor-related polypeptide-encoding described herein. The hybridization probes of the subject invention can be derived from the provided VR1 and VR2 nucleotide sequences, or from their corresponding genomic sequences including promoters, enhancer elements and introns of the naturally occurring capsaicin receptor-encoding sequence.

Hybridization probes can be detectably labeled with a variety of reporter molecules, including radionuclides (e.g.,  $^{32}\text{P}$  or  $^{35}\text{S}$ ), or enzymatic labels (e.g., alkaline phosphatase coupled to the probe via avidin/biotin coupling systems), and the like.

5 Specific hybridization probes can also be produced by cloning the provided nucleic acid sequences into vectors for production of mRNA probes. Such vectors, which are known in the art and are commercially available, can be used to synthesize RNA probes in vitro using an appropriate RNA polymerase (e.g., T7 or SP6 RNA polymerase) and appropriate radioactively labeled nucleotides.

10 Nucleic acids having sequence similarity are detected by hybridization under low stringency conditions, for example, at 50°C and 10XSSC (0.9 M saline/0.09 M sodium citrate) and remain bound when subjected to washing at 55°C in 1XSSC. Sequence identity may be determined by hybridization under stringent conditions, for example, at 50°C or higher and 0.1XSSC (9 mM saline/0.9 mM sodium citrate).

By using probes, particularly labeled probes of DNA sequences, one can isolate homologous or related genes. The source of homologous genes may be any species, e.g. primate species, particularly human; rodents, such as rats and mice, canines, felines, bovines, ovines, equines, yeast, *Drosophila*,  
15 *Caenorhabditis*, etc. Of particular interest is the identification and isolation of human capsaicin receptor polypeptide-encoding polynucleotides and human capsaicin receptor-related polypeptide-encoding polynucleotides.

The capsaicin receptor and capsaicin receptor-related polypeptide nucleic acid sequences can also be used to generate hybridization probes for mapping a naturally occurring genomic sequence. The  
20 sequence can be mapped to a particular chromosome or to a specific region of the chromosome using well known techniques. These include in situ hybridization to chromosomal spreads, flow-sorted chromosomal preparations, or artificial chromosome constructions such as yeast artificial chromosomes, bacterial artificial chromosomes, bacterial P1 constructions or single chromosome cDNA libraries as reviewed in Price 1993; Blood Rev 7:127-34 and Trask 1991; Trends Genet 7:149-54. Fluorescent in situ hybridization  
25 of chromosome spreads is described in, for example, Verma et al 1988 Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York NY.

Information from chromosomal mapping of sequences encoding capsaicin receptor or capsaicin receptor-related polypeptide can be correlated with additional genetic map data. Correlation between the location of a capsaicin receptor-encoding sequence, or a capsaicin receptor-related polypeptide-encoding  
30 sequence, on a physical chromosomal map and a specific disease (or predisposition to a specific disease) can help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention can be used to detect differences in gene sequences (e.g., differences in the chromosomal location due to translocation, inversion, etc. or other differences in the capsaicin receptor-encoding region due to insertional mutation(s) or deletion of capsaicin receptor- or capsaicin receptor-  
35 related polypeptide-encoding sequences) between normal, carrier, or affected individuals. Exemplary disorders that may benefit from such information include, but are not necessarily limited to, complex regional pain syndromes, reflex sympathetic dystrophies, postherpetic neuralgia, psoriasis, reactive airway diseases (e.g., asthma, chronic obstructive pulmonary disease), osteoarthritis, rheumatoid arthritis, diabetic

neuropathy, AIDS-associated neuropathies, and hereditary neuropathies (e.g, associated with capsaicin receptor dysfunction).

Extending the Capsaicin Receptor-Encoding Polynucleotide Sequence

5 The polynucleotide sequence encoding capsaicin receptor or capsaicin receptor-related polypeptide can be extended utilizing partial nucleotide sequence and various methods known in the art to detect upstream sequences such as promoters and regulatory elements. Gobinda et al 1993; PCR Methods Applic 2:318-22 disclose "restriction-site" polymerase chain reaction (PCR) as a direct method which uses universal primers to retrieve unknown sequence adjacent to a known locus. First, genomic DNA is amplified in the presence of primer to a linker sequence and a primer specific to the known region.

10 The amplified sequences are subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR can be used to amplify or extend sequences using divergent primers based on a known region (Triglia et al 1988 Nucleic Acids Res 16:8186). The primers can be designed using OLIGO®

15 4.06 Primer Analysis Software (1992; National Biosciences Inc, Plymouth MN), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. This method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

20 Capture PCR (Lagerstrom et al 1991 PCR Methods Applic 1:111-19) is a method for PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. Capture PCR also requires multiple restriction enzyme digestions and ligations to place an engineered double-stranded sequence into an unknown portion of the DNA molecule before PCR.

Another method that can be used to retrieve unknown sequences is that of Parker et al 1991; Nucleic Acids Res 19:3055-60. Additionally, one can use PCR, nested primers, and PromoterFinder libraries to "walk in" genomic DNA (PromoterFinder™ Clontech (Palo Alto CA). This process avoids the need to screen libraries and is useful in finding intron/exon junctions. Preferably, the libraries used to identify full length cDNAs have been size-selected to include larger cDNAs. More preferably, the cDNA libraries used to identify full-length cDNAs are those generated using random primers, in that such libraries

30 will contain more sequences comprising regions 5' of the sequence(s) of interest. A randomly primed library can be particularly useful where oligo d(T) libraries do not yield a full-length cDNA. Genomic libraries are preferred for identification and isolation of 5' nontranslated regulatory regions of a sequence(s) of interest.

Capillary electrophoresis can be used to analyze the size of, or confirm the nucleotide sequence

35 of, sequencing or PCR products. Systems for rapid sequencing are available from Perkin Elmer, Beckman Instruments (Fullerton CA), and other companies. Capillary sequencing can employ flowable polymers for electrophoretic separation, four different, laser-activatable fluorescent dyes (one for each nucleotide), and a charge coupled device camera for detection of the wavelengths emitted by the fluorescent dyes.

Output/light intensity is converted to electrical signal using appropriate software (e.g. Genotyper™ and Sequence Navigator™ from Perkin Elmer). The entire process from loading of the samples to computer analysis and electronic data display is computer controlled. Capillary electrophoresis is particularly suited to the sequencing of small pieces of DNA that might be present in limited amounts in a particular sample.

- 5 Capillary electrophoresis provides reproducible sequencing of up to 350 bp of M13 phage DNA in 30 min (Ruiz-Martinez et al 1993 Anal Chem 65:2851-2858).

#### Production of Polynucleotides Encoding Capsaicin Receptor or Capsaicin Receptor-Related Polypeptides

- 10 In accordance with the present invention, polynucleotide sequences that encode capsaicin receptor polypeptides or capsaicin receptor-related polypeptides (which capsaicin receptor polypeptides and capsaicin receptor-related polypeptides include fragments of the naturally-occurring polypeptide, fusion proteins, and functional equivalents thereof) can be used in recombinant DNA molecules that direct the expression of capsaicin receptor or capsaicin receptor-related polypeptides in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same
- 15 or a functionally equivalent amino acid sequence, can be used to clone and express capsaicin receptor or capsaicin receptor-related polypeptide. As will be understood by those of skill in the art, it may be advantageous to produce capsaicin receptor-encoding nucleotide sequences and capsaicin receptor-related polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. Codons preferred by a particular prokaryotic or eukaryotic host (Murray et al 1989 Nuc Acids Res 17:477-508) can
- 20 be selected, for example, to increase the rate of expression or to produce recombinant RNA transcripts having a desirable characteristic(s) (e.g., longer half-life than transcripts produced from naturally occurring sequence).

- The nucleotide sequences of the present invention can be engineered in order to alter an capsaicin receptor-encoding sequence or a capsaicin receptor-related polypeptide-encoding sequence for
- 25 a variety of reasons, including but not limited to, alterations that facilitate the cloning, processing and/or expression of the gene product. For example, mutations can be introduced using techniques that are well known in the art, e.g., site-directed mutagenesis to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, etc.

- In another embodiment of the invention, a natural, modified, or recombinant polynucleotide
- 30 encoding a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide can be ligated to a heterologous sequence to encode a fusion protein. Such fusion proteins can also be engineered to contain a cleavage site located between a capsaicin receptor polypeptide-encoding sequence (or capsaicin receptor-related polypeptide-encoding sequence) and a heterologous polypeptide sequence, such that the heterologous polypeptide sequence can be cleaved and purified away from the capsaicin receptor
- 35 polypeptide or capsaicin receptor-related polypeptide.

In an alternative embodiment of the invention, a nucleotide sequence encoding a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide can be synthesized, in whole or in part, using chemical methods well known in the art (see, e.g., Caruthers et al 1980 Nuc Acids Res Symp Ser

215-23, Horn et al (1980) Nuc Acids Res Symp Ser 225-32). Alternatively, the polypeptide itself can be produced using chemical methods to synthesize an amino acid sequence of a capsaicin receptor or capsaicin receptor-related polypeptide, in whole or in part. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge et al 1995 Science 269:202-204) and automated synthesis can be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer) in accordance with the instructions provided by the manufacturer.

The newly synthesized polypeptide can be substantially purified by preparative high performance liquid chromatography (e.g., Creighton 1983 Proteins, Structures and Molecular Principles, WH Freeman and Co, New York NY). The composition of the synthetic polypeptides can be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; Creighton, supra). Additionally the amino acid sequence of capsaicin receptor, capsaicin receptor-related polypeptide, or any part thereof, can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

#### 15 Capsaicin Receptor and Capsaicin Receptor-Related Polypeptide Expression Systems

The invention encompasses expression of capsaicin receptor polypeptides and capsaicin receptor-related polypeptides individually or in combination (e.g., co-expression). In order to express a biologically active capsaicin receptor polypeptide and/or capsaicin receptor-related polypeptide, the nucleotide sequence encoding a capsaicin receptor polypeptide, a capsaicin receptor-related polypeptide, and/or a functional equivalent of either, is inserted into an appropriate expression vector, i.e., a vector having the necessary elements for the transcription and translation of the inserted coding sequence. Methods well known to those skilled in the art can be used to construct expression vectors comprising a desired polypeptide-encoding sequence and appropriate transcriptional or translational controls. These methods include in vitro recombinant DNA techniques, synthetic techniques and in vivo recombination or genetic recombination. Such techniques are described in Sambrook et al 1989 Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY and Ausubel et al 1989 Current Protocols in Molecular Biology, John Wiley & Sons, New York NY.

A variety of expression vector/host cell systems can be utilized to express a capsaicin receptor polypeptide- and capsaicin receptor-related polypeptide-encoding sequence. These include, but are not limited to, amphibian oocytes (e.g., *Xenopus* oocytes); microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transfected with viral expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with bacterial expression vectors (e.g., Ti or pBR322 plasmid); or animal (e.g., mammalian) cell systems. Preferably, the sequences of the present invention, particularly capsaicin receptor-encoding sequences, are expressed in a mammalian cell system (e.g., human embryonic kidney cells (e.g., HEK 293), an amphibian oocyte (e.g., by injecting *Xenopus* oocytes with complementary capsaicin receptor-encoding RNA), or other host cell that is easily propagated in culture and can be

transformed or transfected to either transiently or stably express, preferably stably express, a capsaicin receptor-encoding sequence and/or capsaicin receptor-related polypeptide-encoding sequence).

Host cells can be selected for capsaicin receptor polypeptide and/or capsaicin receptor-related polypeptide expression according to the ability of the cell to modulate the expression of the inserted sequences or to process the expressed protein in a desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation. Post-translational processing that involves cleavage of a "prepro" form of the protein may also be important for correct polypeptide folding, membrane insertion, and/or function. Host cells such as HEK 293, CHO, HeLa, MDCK, WI38, *Xenopus* oocytes, and others have specific cellular machinery and characteristic mechanisms for such post-translational activities and may be chosen to ensure the correct modification and processing of the introduced, foreign polypeptide.

The vector(s) used for expression of a capsaicin receptor polypeptide and/or capsaicin receptor-related polypeptide will vary with a variety of factors including the host cell in which the capsaicin receptor polypeptide is to be expressed, whether capsaicin receptor polypeptide- and capsaicin receptor-related polypeptide sequences are to be co-expressed either from a single construct or from separate constructs, and the intended use for the polypeptide produced. For example, when large quantities of a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide are required (e.g., for the antibody production), vectors that direct high-level expression of fusion proteins that can be readily purified may be desirable. Such vectors include, for example, bacterial expression vectors, including multifunctional *E. coli* cloning and expression vectors such as Bluescript® (Stratagene; which provides for production of polypeptide- $\beta$ -galactosidase hybrid proteins); and pGEX vectors (Promega, Madison WI; which provides for production of glutathione S-transferase (GST) fusion proteins. Where the host cell is yeast (e.g., *Saccharomyces cerevisiae*) a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH can be used. For reviews, see Ausubel et al (supra) and Grant et al 1987 Methods in Enzymology 153:516-544.

Where the host cell is a mammalian cells, a number of expression systems can be used. For example, the expression vector can be derived from a viral-based expression system, such as an expression system derived from an adenovirus, SV40, CMV, or RSV nucleotide sequence. Expression efficiency can be enhanced by including enhancers appropriate to the cell system in use (Scharf et al 1994 Results Probl Cell Differ 20:125-62; Bittner et al 1987 Methods in Enzymol 153:516-544) (e.g., the RSV enhancer can be used to increase expression in mammalian host cells).

The "control elements" or "regulatory sequences" of these systems, which vary in their strength and specificities, are those nontranslated regions of the vector, enhancers, promoters, and 3' untranslated regions that interact with host cellular proteins to facilitate transcription and translation of a nucleotide sequence of interest. Depending on the vector system and host utilized, any number of suitable transcriptional and translational elements, including constitutive and inducible promoters, can be used.

Such control elements or regulatory sequences are selected according to the host cell in which the capsaicin receptor-encoding polynucleotide and/or capsaicin receptor-related polypeptide-encoding



polynucleotide is to be expressed. For example, in mammalian cell systems, promoters from the mammalian genes or from mammalian viruses are most appropriate. Where it is desirable to generate a cell line containing multiple copies of a capsaicin receptor polypeptide-encoding sequence or a capsaicin receptor-related polypeptide-encoding sequence, vectors derived from SV40 or EBV can be used in conjunction with other optional vector elements, e.g., an appropriate selectable marker.

Specific initiation signals may also be required for efficient translation of a capsaicin receptor polypeptide- or capsaicin receptor-related polypeptide-encoding sequence, e.g., the ATG initiation codon and flanking sequences for bacterial expression. Where a native sequence, including its initiation codon and upstream sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, where only coding sequence, or a portion thereof, is inserted in an expression vector, exogenous transcriptional control signals including the ATG initiation codon must be provided. Furthermore, the initiation codon must be in the correct reading frame to ensure transcription of the entire insert. Exogenous transcriptional elements and initiation codons can be derived from various origins, and can be either natural or synthetic.

Where long-term, high-yield recombinant polypeptide production is desired, stable expression is preferred. For example, cell lines that stably express capsaicin receptor and/or capsaicin receptor-related polypeptide can be transformed using expression vectors containing viral origins of replication or endogenous expression elements and a selectable marker gene. After introduction of the vector, cells can be grown in an enriched media before they are exposed to selective media. The selectable marker, which confers resistance to the selective media, allows growth and recovery of cells that successfully express the introduced sequences. Resistant, stably transformed cells can be proliferated using tissue culture techniques appropriate to the host cell type.

Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler et al 1977 Cell 11:223-32) and adenine phosphoribosyltransferase (Lowy et al 1980 Cell 22:817-23) genes which can be employed in tk- or aprt- cells, respectively. Also, antimetabolite or antibiotic resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler et al 1980 Proc Natl Acad Sci 77:3567-70); and npt, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin et al 1981 J Mol Biol 150:1-14). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman et al. 1988 Proc Natl Acad Sci 85:8047-51). Selectable markers also include visible markers such as anthocyanins,  $\beta$ -glucuronidase and its substrate, GUS, and luciferase and its substrate, luciferin. Such visible markers are useful to both identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes et al 1995 Methods Mol Biol 55:121-131).

Alternatively, host cells that contain the coding sequence for and express capsaicin receptor polypeptides and/or capsaicin receptor-related polypeptides can be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA

hybridization and protein bioassay or immunoassay techniques for the detection and/or quantitation of the nucleic acid or protein.

The presence of polynucleotide sequences encoding a capsaicin receptor and/or capsaicin receptor-related polypeptide can be detected by DNA-DNA or DNA-RNA hybridization or PCR amplification using probes, portions or fragments of polynucleotides encoding capsaicin receptor and/or capsaicin receptor-related polypeptide. Nucleic acid amplification-based assays involve the use of oligonucleotides or oligomers based on a sequence encoding a capsaicin receptor or capsaicin receptor-related polypeptide to detect transformants containing the desired DNA or RNA. As used herein "oligonucleotides" or "oligomers" refer to a nucleic acid sequence of at least about 10 nucleotides and as many as about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20-25 nucleotides which can be used as a probe or amplifier.

A variety of immunoassays for detecting and measuring the expression of a specific protein, using either protein-specific polyclonal or monoclonal antibodies are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). These and other assays are described in, e.g., Hampton et al 1990, Serological Methods, A Laboratory Manual, APS Press, St Paul MN and Maddox et al 1983, J Exp Med 158:1211.

A wide variety of detectable labels and conjugation techniques are known the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to sequences encoding a capsaicin receptor or capsaicin receptor-related polypeptide include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, a nucleotide sequence encoding a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide can be cloned into a vector for the production of an mRNA probe. Vectors and methods for production of mRNA probes are well known in the art. Suitable reporter molecules or labels include those radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles and the like, as described in U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149 and 4,366,241, each of which are incorporated herein by reference.

In a preferred embodiment, host cells expressing capsaicin receptor are screened and selected using a functional assay for capsaicin receptor activity. For example, host cells expressing functional capsaicin receptor can be screened for alterations in intracellular calcium concentrations upon exposure to a capsaicin receptor binding compound (e.g., capsaicin or resiniferatoxin). Where the capsaicin receptor binding compound is a capsaicin receptor agonist, binding of the agonist compound to the capsaicin receptor result in increased levels of intracellular calcium in the host cell expressing capsaicin receptor-encoding nucleic acid. Methods and compositions (e.g., fura-2) for monitoring intracellular calcium concentration are well known in the art.

#### Purification of Capsaicin Receptor Polypeptides and Capsaicin Receptor-Related Polypeptides

Methods for production of a polypeptide after identification of its encoding polynucleotide are well known in the art. Host cells transformed with a nucleotide sequence(s) encoding a capsaicin receptor

polypeptide and/or capsaicin receptor-related polypeptide-can be cultured under conditions suitable for the expression and recovery of the encoded polypeptide from cell culture. The polypeptide produced by a recombinant cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides  
5 encoding capsaicin receptor polypeptides or capsaicin receptor-related polypeptides can be designed with signal sequences that direct secretion of the encoded polypeptide(s) through a prokaryotic or eukaryotic cell membrane.

Purification of capsaicin receptor polypeptides and capsaicin receptor-related polypeptides can be facilitated by using a recombinant construct that includes a nucleotide sequence(s) encoding one or  
10 more polypeptide domains that, when expressed in-frame with the sequence encoding the capsaicin receptor or capsaicin receptor-related polypeptide, provides a fusion protein having a purification-facilitating domain (Kroll et al 1993 DNA Cell Biol 12:441-53). A cleavable linker sequences(s) between the purification domain and the capsaicin receptor polypeptide- or capsaicin receptor-related polypeptide-encoding sequence can be included to further facilitate purification.

15 Capsaicin receptor polypeptides and capsaicin receptor-related polypeptides (each of which polypeptides encompass polypeptides having a portion of the native amino acid sequence) can also be produced by direct peptide synthesis using solid-phase techniques (see, e.g., Stewart et al: 1969 Solid-Phase Peptide Synthesis, WH Freeman Co, San Francisco; Merrifield 1963 J Am Chem Soc 85:2149-2154). Various fragments of capsaicin receptor or capsaicin receptor-related polypeptide can be  
20 chemically synthesized separately and combined using chemical methods to produce the full length molecule.

Methods for purifying a desired polypeptide following either artificial synthesis or recombinant production are routine and well known in the art.

#### Uses of Capsaicin Receptor Polypeptides, Capsaicin Receptor-Related Polypeptides, and Nucleic Acid

##### 25 Encoding Capsaicin Receptor Polypeptides or Capsaicin Receptor-Related Polypeptides

In addition to the uses described above, the nucleotide and polypeptide sequences disclosed herein can be used in a variety of ways, including production of antibodies, identification of capsaicin receptor-binding compounds and capsaicin receptor-related polypeptide-binding compounds that affect capsaicin receptor function (e.g., in a drug screening assay), and in the identification of other  
30 polynucleotide sequences encoding capsaicin receptor polypeptides and capsaicin receptor-related polypeptides. In addition, sequences encoding capsaicin receptor polypeptides and capsaicin receptor-related polypeptides can be used in diagnostic assays (e.g., prenatal or postnatal diagnosis). Furthermore, capsaicin receptor-encoding sequences and their encoded polypeptides can also be used in assays to assess the capsaicin content of a sample (e.g., from a natural product, e.g, a chili pepper extract) or the  
35 capsaicin-promoting effects of an agent (e.g., a candidate agent for use a flavor enhancing additive to foods).

These and other applications of the sequences of the invention are described in more detail below.

Screening for Capsaicin Receptor- and Capsaicin Receptor-Related Polypeptide Binding Compounds

5 Capsaicin receptor polypeptides and capsaicin receptor-related polypeptides, each of which encompasses biologically active or immunogenic fragments or oligopeptides thereof, can be used for screening compounds that affect capsaicin receptor activity by, for example, specifically binding capsaicin receptor and affecting its function or specifically binding capsaicin receptor-related polypeptide and affecting its interaction with capsaicin receptor, thereby affecting capsaicin receptor activity. Identification of such compounds can be accomplished using any of a variety of drug screening techniques. Of particular interest is the identification of agents that have activity in affecting capsaicin receptor function.

10 Such agents are candidates for development of treatments for, inflammatory conditions associated at least in part with capsaicin receptor activity (e.g., psoriasis, reactive airway diseases (e.g., asthma, chronic obstructive pulmonary disease)), arthritis (e.g., osteoarthritis, rheumatoid arthritis), and for use as analgesics. Of particular interest are screening assays for agents that have a low toxicity for human cells. The polypeptide employed in such a test can be free in solution, affixed to a solid support, present on a cell surface, or located intracellularly. The screening assays of the invention are generally based upon the ability of the agent to bind to a capsaicin receptor polypeptide, bind to a capsaicin receptor-related polypeptide, and/or elicit or inhibit a capsaicin receptor-associated or capsaicin receptor-related polypeptide-associated biological activity (*i.e.*, a functional assay or an assay using radioligand binding assays).

20 The term "agent" as used herein describes any molecule, e.g. protein or pharmaceutical, with the capability of altering (*i.e.*, eliciting or inhibiting) or mimicking a desired physiological function of capsaicin receptor or capsaicin receptor-related polypeptide. Generally a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, *i.e.* at zero concentration or below the level of detection.

25 Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including, but not limited to: peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

35 Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial,

5 fungal, plant and animal extracts (including extracts from human tissue to identify endogenous factors affecting capsaicin receptor or capsaicin receptor-related polypeptide activity) are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce  
5 combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, *etc.* to produce structural analogs.

Preferably, the drug screening technique used provides for high throughput screening of compounds having suitable binding affinity to the capsaicin receptor, capsaicin receptor-related polypeptide, and/or eliciting a desired capsaicin receptor-associated or capsaicin receptor-related  
10 polypeptide-associated response. For example, large numbers of different small peptide test compounds can be synthesized on a solid substrate, such as plastic pins or some other surface (see, *e.g.*; Geysen WO Application 84/03564, published on September 13, 1984), the peptide test compounds contacted with capsaicin receptor polypeptides (or capsaicin receptor-related polypeptides), unreacted materials washed away, and bound capsaicin receptor (or bound capsaicin receptor-related polypeptide) detected by virtue  
15 of a detectable label or detection of a biological activity associated with capsaicin receptor activity (or capsaicin receptor-related polypeptide activity). Purified capsaicin receptor or purified capsaicin receptor-related polypeptide can also be coated directly onto plates for use in such in vitro drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the polypeptide and immobilize it on a solid support.

20 The invention also contemplates the use of competitive drug screening assays in which capsaicin receptor-specific neutralizing antibodies or capsaicin receptor-related polypeptide-specific neutralizing antibodies compete with a test compound for binding of capsaicin receptor polypeptide or capsaicin receptor-related polypeptide. In this manner, the antibodies can be used to detect the presence of any polypeptide that shares one or more antigenic determinants with a capsaicin receptor polypeptide or  
25 capsaicin receptor-related polypeptide.

#### Screening of Candidate Agents

A wide variety of assays may be used for identification of capsaicin receptor polypeptide and/or capsaicin receptor-related polypeptide binding agents, including labeled *in vitro* binding assays, immunoassays for protein binding, and the like. For example, by providing for the production of large  
30 amounts of capsaicin receptor polypeptides or capsaicin receptor-related polypeptides, one can identify ligands or substrates that bind to, modulate or mimic the action of the proteins. The purified protein may also be used for determination of three-dimensional crystal structure, which can be used for modeling intermolecular interactions.

The screening assay can be a binding assay, wherein one or more of the molecules may be joined  
35 to a label, and the label directly or indirectly provide a detectable signal. Various labels include radioisotopes, fluorescers, chemilumescers, enzymes, specific binding molecules, particles, *e.g.* magnetic particles, and the like. Specific binding molecules include pairs, such as biotin and streptavidin,

digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule that provides for detection, in accordance with known procedures.

A variety of other reagents may be included in the screening assays described herein. Where the assay is a binding assay, these include reagents like salts, neutral proteins, *e.g.* albumin, detergents, etc that are used to facilitate optimal protein-protein binding, protein-DNA binding, and/or reduce non-specific or background interactions. Reagents that improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, *etc.* may be used. The mixture of components are added in any order that provides for the requisite binding. Incubations are performed at any suitable temperature, typically between 4 and 40°C. Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high-throughput screening. Typically between 0.1 and 1 hours will be sufficient.

#### Functional Capsaicin Receptor and Capsaicin Receptor-Related Polypeptide Screening Assays

Preferably, capsaicin receptor-binding compounds are screened for agonistic or antagonist action in a functional assay that monitors a biological activity associated with capsaicin receptor function such as effects upon intracellular levels of cations in a capsaicin receptor-expressing host cell (*e.g.*, calcium, magnesium, guanidinium, cobalt, potassium, cesium, sodium, and choline, preferably calcium), ligand-activated conductances, cell death (*i.e.*, receptor-mediated cell death which can be monitored using, *e.g.*, morphological assays, chemical assays, or immunological assays), depolarization of the capsaicin receptor-expressing cells (*e.g.*, using fluorescent voltage-sensitive dyes), second messenger production (*e.g.*, through detection of changes in cyclic GMP levels (see, *e.g.*, Wood et al. 1989 J. Neurochem. 53:1203-1211), which can be detected by radioimmunoassay or ELISA), calcium-induced reporter gene expression (see, *e.g.*, Ginty 1997 Neuron 18:183-186), or other readily assayable biological activity associated with capsaicin receptor activity or inhibition of capsaicin receptor activity. Preferably, the functional assay is based upon detection of a biological activity of capsaicin receptor that can be assayed using high-throughput screening of multiple samples simultaneously, *e.g.*, a functional assay based upon detection of a change in fluorescence which in turn is associated with a change in capsaicin receptor activity. Such functional assays can be used to screen candidate agents for activity as either capsaicin receptor agonists or antagonists.

In a preferred embodiment, capsaicin receptor-expressing cells (preferably recombinant capsaicin receptor-expressing cells) are pre-loaded with fluorescently-labeled calcium (*e.g.* fura-2). The capsaicin receptor-expressing cells are then exposed to a candidate capsaicin receptor-binding compound and the effect of exposure to the compound monitored. Candidate compounds that have capsaicin receptor agonist activity are those that, when contacted with the capsaicin receptor-expressing cells, elicit a capsaicin receptor-mediated increase in intracellular calcium relative to control cells (*e.g.*, capsaicin receptor-expressing cells in the absence of the candidate compound, host cells without capsaicin receptor-encoding nucleic acid, capsaicin receptor-expressing cells exposed to a known capsaicin receptor agonist). Similarly, functional capsaicin receptor assays can be used to identify candidate compounds that block activity of a known capsaicin receptor agonist (*e.g.*, block the activity of or compete with capsaicin or

resiniferatoxin), block activity of a known capsaicin receptor antagonist (e.g., block the activity of or compete with capsazepine), and/or have activity as capsaicin receptor antagonists.

In another embodiment, the invention includes a method for identifying compounds that bind capsaicin receptor-related polypeptide, thereby eliciting an agonistic or antagonistic effect on capsaicin receptor-associated function as detected by e.g., intracellular levels of cations in the host cell. To this end, the functional assay involves contacting host cells expressing a capsaicin receptor alone (e.g., VR1) and with host cell co-expressing a capsaicin receptor and a capsaicin receptor-related polypeptide (e.g., VR1 and VRRP-1). Compounds that affect capsaicin receptor activity by affecting function of a capsaicin receptor-related polypeptide are those that affect a capsaicin receptor-associated activity in cells that co-express capsaicin receptor and capsaicin receptor-related polypeptide, but do not significantly affect capsaicin receptor-associated activity in host cells that express capsaicin receptor alone. For example, compounds that elicit a capsaicin receptor-mediated increase in intracellular calcium in cells co-expressing capsaicin receptor and capsaicin receptor-related polypeptide, but not in cells expressing capsaicin receptor alone, are identified as compounds that elicit capsaicin receptor agonist activity via interaction with a capsaicin receptor-related polypeptide.

Pharmaceutical Compositions and Other Compositions Comprising Agents Affecting Capsaicin Receptor Activity Identified by the Screening Assay of the Invention

Capsaicin receptor-binding compounds and capsaicin receptor-related polypeptide-binding compounds are useful in eliciting or inhibiting capsaicin receptor-mediated physiological responses, and can be particularly useful in a pharmaceutical composition for ameliorating symptoms associated with chronic pain, inflammation, and other physiological responses associated with capsaicin receptor-mediated activity.

The compounds having the desired pharmacological activity may be administered in a physiologically acceptable carrier to a host for treatment of a condition attributable to capsaicin receptor activity. Alternatively, the identified compounds may be used to enhance, regulate, or otherwise manipulate capsaicin receptor function. The therapeutic agents may be administered in a variety of ways, topically, subcutaneously, intraperitoneally, intravascularly, orally, intrathecally, epidermally, intravesicularly (e.g., as in bladder irrigation to treat neurogenic bladder syndromes), parenterally, etc. Inhaled treatments are of particular interest for the treatment of capsaicin receptor-associated inflammation associated with such conditions as asthma.

Depending upon the manner of introduction, the compounds may be formulated in a variety of ways. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt.%. The pharmaceutical compositions can be prepared in various forms, such as granules, tablets, pills, capsules, suspensions, salves, lotions and the like. Pharmaceutical grade organic or inorganic carriers and/or diluents suitable for the selected route of administration can be used to make up compositions containing the therapeutically-active compounds. Diluents known to the art include aqueous media, vegetable and animal oils and fats. Stabilizing agents, wetting and emulsifying agents, salts for

varying the osmotic pressure or buffers for securing an adequate pH value, and skin penetration enhancers can be used as auxiliary agents.

In addition, compositions comprising agents affecting capsaicin receptor activity (e.g., by binding capsaicin receptor or by binding a capsaicin receptor-related polypeptide) are useful in other applications, including use in defensive sprays (e.g., "pepper sprays") or as antidotes for such sprays. The screening methods of the invention can be used in a variety of ways to this end, including, for example, identification of drugs that have capsaicin-like activity, but lack or are substantially diminished in one or more of the undesirable side effects associated with capsaicin. For example, while capsaicin is effective in spray deterrents, exposure to capsaicin can be lethal. The screening method of the invention can thus be used to identify compounds that have the desired deterrent effect, but would not likely cause death upon exposure to amounts normally used in defensive sprays. Moreover, the screening method of the invention could be used to identify compounds that differentially affect capsaicin receptors of different mammalian species, thus enabling identification and design of capsaicin receptor agonists and antagonists that substantially affect capsaicin receptors with genus- or species-specificity. Thus, for example, the method of the invention can allow for identification of capsaicin receptor agonists for canine or bear capsaicin receptors, but that do not substantially stimulate human capsaicin receptors. This could be accomplished by screening for compounds that elicit a capsaicin receptor-associated biological activity in host cells expressing a canine capsaicin receptor, but relatively little or no biological activity in host cells expressing human capsaicin receptor.

#### Therapeutically Effective Dosages

The determination of an effective dose is well within the capability of those skilled in the art. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., using host cells expressing recombinant capsaicin receptor, or in animal models, usually rats, mice, rabbits, dogs, or pigs. The animal model is also used to achieve a desirable concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans. A therapeutically effective dose refers to that amount of an agent (e.g., a compound having activity as capsaicin receptor agonist or antagonist), polypeptide, or anti-polypeptide antibody, that provide the desired physiological effect (e.g., to ameliorate symptoms associated with capsaicin receptor-mediated inflammation or pain, or provide loss of temperature sensation).

Therapeutic efficacy and toxicity of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and expressed as the ratio LD50/ED50. Pharmaceutical compositions that exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The actual dosage can vary within this range depending upon, for example, the dosage form employed, sensitivity of the patient, and the route of administration.



The exact dosage is chosen by the individual physician in view of the patient to be treated. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Additional factors that may be taken into account include the severity of the disease state, location of the site to be treated; age, weight and gender of the patient; diet, time and frequency of administration; drug combination(s); reaction sensitivities; and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Use of capsaicin and capsaicin analogues in clinical applications and their methods of administration (e.g., formulations, dosages, routes of administration, etc.) are well known in the art (see, e.g., Campbell et al. 1993 "Clinical Applications of Capsaicin and Its Analogues," in Capsaicin in the Study of Pain, pgs. 255-272; USPN 5,569,910 (topical anti-inflammatory compositions comprising capsaicin); USPN 5,296,225 (topical composition comprising capsaicin for treating orofacial pain); USPN 5,290,816 (topical cream containing resiniferatoxin for desensitization of neurogenic inflammation); USPN 4,997,853 (topical composition containing capsaicin for treating superficial pain); USPN 5,403,868 (capsaicin derivatives useful as analgesic and anti-inflammatory agents); USPN 4,939,149 (administration of resiniferatoxin to cause sensory afferent C-fibre and thermo-regulatory desensitization); USPN 4,536,404 (topical treatment of post herpetic neuralgia by application of capsaicin), each of which is incorporated by reference in its entirety. Further general guidance on administration of capsaicin receptor agonist and antagonists can be found in, e.g., United States Pharmacopeia (USP), 17<sup>th</sup> Ed., pgs. 710-711; and Physician's Desk Reference 1996, Medical Economics Com., Montvale, NJ (see particularly Dolorac<sup>TM</sup> at 1054, Zostrix<sup>TM</sup> at 1056, and Zostrix-HP<sup>TM</sup> topical analgesic cream at 1056, each of which contain capsaicin); and the latest edition of Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton PA.

Use of Capsaicin Receptor-Encoding Polynucleotides in Assays for Quantitating the Capsaicin Content of a Sample or Determining the Capsaicin Activity of a Candidate Food Additive

Capsaicin receptor polypeptide-encoding polynucleotides and capsaicin receptor polypeptides can be used in an assay to determine, either qualitatively or quantitatively, to detect capsaicin or an agent having capsaicin activity, in a sample, where the sample is derived from a food product or contains a candidate agent for use as a flavoring agent (e.g., for use as a spice in food or food products). This assay takes advantage of the fact that, in addition to its analgesic effects upon afferent neurons, capsaicin is a member of the vanilloid family of compounds, which are responsible for making foods "spicy hot." For example, capsaicin is present in peppers (e.g., Thai green poblano verde, habenero, and guero peppers). Conventional assays for determining the amount of capsaicin in a pepper extract involve tedious extraction of the compound from pepper samples and quantitation by high pressure liquid chromatography (HPLC) (see, e.g., Woodbury 1980 J. Assoc. Off. Anal. Chem. 63:556-558). The amount of capsaicin is then correlated with number of Scoville heat units, a measure of "hotness."

The assay of the invention uses an isolated capsaicin receptor polypeptide to detect the amount of capsaicin in a sample, thus avoiding the chemical extraction technique employed in the conditional assay. The capsaicin receptor polypeptide used may be either bound to a solid support, present in solution, or present on the surface of a recombinant host cell. Binding of capsaicin to the capsaicin receptor polypeptide is detected as described in the screening assays described above.

Preferably, the assay for capsaicin or a compound having capsaicin activity in a sample is performed using a functional assay described above. More preferably, the functional assay uses capsaicin receptor-expressing recombinant eukaryotic cells (preferably mammalian cells or amphibian oocytes) that are preloaded with a calcium-sensitive fluorescent dye (e.g., fura-2, indo-1, fluo-3). The presence and/or amount of capsaicin or capsaicinoid compound in the sample is then determined by measuring a capsaicin receptor-mediated cellular effect, e.g., an alteration in voltage-activated conductances across the cellular membrane or an alteration in the intracellular levels of the detectably labeled cation. For example, where the detectably labeled cation is fluorescently labeled calcium, exposure of the pre-loaded host cells to a capsaicin-containing sample results in binding of the capsaicin to the capsaicin receptor polypeptide and the capsaicin receptor-mediated increase in intracellular calcium, which can be readily detected and quantitated. For example, the level of intracellular calcium influx mediated by the test sample is compared to the intracellular calcium influx associated with a control sample (e.g., with a sample having a known amount of capsaicin). The extent of the change in current, intracellular calcium concentration, or other capsaicin receptor-mediated phenomenon is then correlated with a concentration of capsaicin, which in turn can be assigned a Scoville heat unit.

Similarly, candidate agents for use as food additives to make a food or food product "hot" can be screened for their ability to elicit a capsaicin receptor-mediated cellular response (e.g., change in voltage-activated conductances or intracellular cation concentration). The assay has the advantage that the measure of hotness can be determined objectively, e.g., based upon the responses elicited by exposure to the capsaicin receptor.

Diagnostic Uses of Polynucleotides Encoding Capsaicin Receptor or Capsaicin Receptor-Related Polypeptides to Detect Capsaicin Receptor-Encoding Sequences

Polynucleotide sequences encoding capsaicin receptor polypeptide or capsaicin receptor-related polypeptide can be used in the diagnosis (e.g., prenatal or post-natal diagnosis) of conditions or diseases associated with, for example, capsaicin receptor expression, with a particular capsaicin receptor polymorphism or mutation, and/or with capsaicin receptor-related polypeptide expression. For example, polynucleotide sequences encoding capsaicin receptor or capsaicin receptor-related polypeptide can be used in hybridization or PCR assays of fluids or tissues from biopsies to detect capsaicin receptor or capsaicin receptor-related polypeptide expression, respectively. Suitable qualitative or quantitative methods include Southern or northern analysis, dot blot or other membrane-based technologies; PCR technologies; dip stick, pIN, chip and ELISA technologies. All of these techniques are well known in the art and are the basis of many commercially available diagnostic kits. Once disease is established, a

therapeutic agent is administered or other intervention or precautions initiated as appropriate for the capsaicin receptor-associated disorder.

Oligonucleotides based upon capsaicin receptor or capsaicin receptor-related polypeptid sequences can be used in PCR-based techniques for assessing capsaicin receptor-polypeptide expression, detection of capsaicin receptor polymorphisms associated with disorders, and/or capsaicin receptor-related polypeptide expression. Methods for PCR amplification are described in U.S. Patent Nos. 4,683,195 and 4,965,188. Such oligomers are generally chemically synthesized, or produced enzymatically or by recombinantly. Oligomers generally comprise two nucleotide sequences, one with sense orientation (5'→3') and one with antisense (3'←5'), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers can be employed under less stringent conditions for detection and/or quantitation of closely related DNA or RNA sequences.

Additional methods for quantitation of expression of a particular molecule according to the invention include radiolabeling (Melby et al 1993 J Immunol Methods 159:235-44) or biotinylating (Duplaa C 1993 Anal Biochem 229-36) nucleotides, coamplification of a control nucleic acid, and interpolation of experimental results according to standard curves. Quantitation of multiple samples can be made more time efficient by running the assay in an ELISA format in which the oligomer of interest is presented in various dilutions and rapid quantitation is accomplished by spectrophotometric or colorimetric detection.

#### Therapeutic Uses of Capsaicin Receptor Polypeptides and Capsaicin Receptor Polypeptide-Encoding Nucleic Acid

Polypeptides of, as well as nucleotide sequence encoding, capsaicin receptor polypeptides and capsaicin receptor-related polypeptides may be useful in the treatment of conditions associated with capsaicin receptor dysfunction (e.g., capsaicin receptor activity that is increased relative to capsaicin receptor activity in an unaffected patient or capsaicin receptor activity that is decreased relative to capsaicin receptor activity in an unaffected patient). In addition, expression of dominant-negative capsaicin receptor-encoding sequences may be therapeutically useful in a condition associated with elevated levels of capsaicin receptor activity. Where interaction of capsaicin receptor and a capsaicin receptor-related polypeptide is associated with a condition, interaction of these polypeptides can be disrupted by, for example, introduction of a peptide corresponding to an interaction domain of capsaicin receptor and capsaicin receptor-related polypeptide. Moreover, expression of a wild-type capsaicin receptor sequence in tumor cells may render such tumor cells more susceptible to capsaicin receptor-mediated cell death.

Expression vectors derived from retroviruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids, can be used for delivery of nucleotide sequences to the targeted organ, tissue or cell population. Preferably the targeted cell for delivery and expression of capsaicin receptor polypeptide-encoding sequences is a neuronal cell, more preferably an afferent neuron in order to enhance capsaicin receptor activity in the neuronal cell. Recombinant vectors for expression of antisense capsaicin receptor polynucleotides can be constructed according to methods well known in the art (see, for example, the techniques described in Sambrook et al (supra) and Ausubel et al (supra)).

Alternatively, expression of genes encoding capsaicin receptor can be decreased by transfecting a cell or tissue with expression vectors that express high levels of a desired capsaicin receptor-encoding fragment. Such constructs can flood cells with untranslatable sense or antisense sequences. Even in the absence of integration into the DNA, such vectors can continue to transcribe RNA molecules until all copies are disabled by endogenous nucleases. Such an approach to regulation of capsaicin receptor expression and activity can be useful in treatment of pain syndromes and/or inflammatory conditions associated with capsaicin receptor activity.

Modifications of gene expression can be obtained by designing antisense molecules, DNA, RNA or PNA, to the control regions of gene encoding capsaicin receptor (*i.e.*, the promoters, enhancers, and introns). Oligonucleotides derived from the transcription initiation site, *e.g.*, between -10 and +10 regions of the leader sequence, are preferred. The antisense molecules can also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes. Similarly, inhibition of expression can be achieved using "triple helix" base-pairing methodology. Triple helix pairing compromises the ability of the double helix to open sufficiently for binding of polymerases, transcription factors, or regulatory molecules.

Recent therapeutic advances using triplex DNA were reviewed by Gee JE et al (In: Huber et al. 1994 Molecular and Immunologic Approaches, Futura Publishing Co, Mt Kisco NY). Antisense molecules of the invention can be prepared by methods known in the art for the synthesis of RNA molecules, including techniques for chemical oligonucleotide synthesis, *e.g.*, solid phase phosphoramidite chemical synthesis. Such DNA sequences can be incorporated into a wide variety of vectors with suitable RNA polymerase promoters (*e.g.*, T7 or SP6). Alternatively, antisense cDNA constructs useful in the constitutive or inducible synthesis of antisense RNA can be introduced into cell lines, cells, or tissues.

Particularly where RNA molecules are to be administered for antisense therapy, the RNA can be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine and wybutosine as well as acetyl-, methyl-, thio- and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine that are not as easily recognized by endogenous endonucleases.

Methods for introducing vectors into cells or tissues include those methods discussed *infra* and which are equally suitable for *in vivo* therapy.

In a preferred embodiment, capsaicin receptor polypeptide-encoding polynucleotides are introduced *in vivo* into a target tumor cell for which organochemotherapy is desired. This aspect of the invention takes advantage of the range of capsaicin receptor response to exposure to agonists (*e.g.*, capsaicin, resiniferatoxin) and/or to temperature. For example, low concentrations of capsaicin receptor agonists (*e.g.*, capsaicin receptor agonist concentrations in the nanomolar range, *e.g.*, from about 200 nM to about 800 nM) are associated with capsaicin receptor stimulation and intracellular calcium influx. Where the capsaicin receptor is expressed in a neuronal cell, capsaicin receptor stimulation by low concentrations

of capsaicin receptor agonist is followed by neuronal desensitization. However, high concentrations of capsaicin receptor agonists (e.g., capsaicin receptor agonist concentrations in the micromolar range, e.g., from about 1  $\mu$ M to about 10  $\mu$ M) mediate neuronal degeneration and cell death. By expressing capsaicin receptor polypeptides in tumor cells, the tumor cell death can be substantially selectively facilitated by local administration of high concentrations of a capsaicin receptor agonist, or by local exposure to heat stimuli, or both, where the agonist concentration and/or heat stimulus is sufficient to mediate cell death in the capsaicin receptor-expressing target tumor cell, but does not substantially affect normal capsaicin receptor-expressing cells or affects a minimal number of such normal cells.

Alternatively, the capsaicin receptor polypeptide introduced into the tumor cells can be engineered to provide more selectivity in the response to organochemotherapy (i.e., to provide for activation of the capsaicin receptor expressed in the tumor cells with no or little activation of endogenous, wild-type capsaicin receptor). For example, capsaicin receptor can be modified so as to bind a specific capsaicin receptor agonist analogue, which analogue is substantially reduced in its ability to bind wildtype capsaicin receptors. Therefore, target cells (e.g., tumor cells) expressing the altered capsaicin receptor can be selectively stimulated by administration of the agonist having specificity for the altered capsaicin receptor polypeptide without substantially affecting cells expressing wildtype capsaicin receptor. Alternatively, the tumor cells can be transformed in vivo with a sequence encoding a modified capsaicin receptor, where the modified capsaicin receptor is more responsive to agonists (e.g., is more responsive to agonist, has increased affinity to agonists relative to wild-type thereby allowing activation of the modified receptors with no or little activation of the endogenous capsaicin receptor, and/or is modified so as to be more responsive to heat stimuli than wild-type capsaicin receptor). These embodiments thus allow administration of high or higher concentrations of the altered capsaicin receptor-targeted organochemotherapeutic, thereby providing for more selective organochemotherapy.

#### Anti-Capsaicin Receptor and Anti-Capsaicin Receptor-Related Polypeptide Antibodies

Capsaicin receptor-specific antibodies and capsaicin receptor-related polypeptide-specific antibodies are useful for identification of cells expressing either naturally-occurring or recombinant capsaicin receptor polypeptides or capsaicin receptor-related polypeptides, respectively, as well as the diagnosis of conditions and diseases associated with expression and/or function of capsaicin receptor and/or capsaicin receptor-related polypeptides. For example, anti-capsaicin receptor antibodies and anti-capsaicin receptor-related polypeptide antibodies can be used to detect increased or decreased receptor protein levels, and/or aberrant protein processing or oligomerization.

Anti-capsaicin receptor polypeptide antibodies and anti-capsaicin receptor-related polypeptide antibodies of the invention include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments and fragments produced by a Fab expression library. Antibodies of particular interest include, for example, antibodies that stimulate capsaicin receptor function and/or block binding of capsaicin receptor-binding compounds to capsaicin receptor. Such antibodies may be useful in, for example,

regulation of pain in pain syndromes, in screening assays for capsaicin receptor-binding agents, and in measurement of capsaicin receptor-activating compounds in a sample.

Capsaicin receptor polypeptides and capsaicin receptor-related polypeptides suitable for production of antibodies need not be biologically active; rather, the polypeptide, or oligopeptide need only be antigenic. Polypeptides used to generate capsaicin receptor-specific antibodies and capsaicin receptor-related polypeptide antibodies generally have an amino acid sequence consisting of at least five amino acids, preferably at least 10 amino acids. Preferably, antigenic capsaicin receptor polypeptides and antigenic capsaicin receptor-related polypeptides mimic an epitope of the native capsaicin receptor or native capsaicin receptor-related polypeptide, respectively. Antibodies specific for short polypeptides can be generated by linking the capsaicin receptor polypeptide or capsaicin receptor-related polypeptide to a carrier, or fusing the capsaicin receptor polypeptide or capsaicin receptor-related polypeptide to another protein (e.g., keyhole limpet hemocyanin), and using the carrier-linked chimeric molecule as an antigen. In general, anti-capsaicin receptor antibodies and capsaicin receptor-related polypeptide antibodies can be produced according to methods well known in the art. Recombinant immunoglobulins can be produced as according to U.S. Patent No. 4,816,567, incorporated herein by reference.

Various hosts, generally mammalian or amphibian hosts, can be used to produce anti-capsaicin receptor antibodies and anti-capsaicin receptor-related polypeptide antibodies (e.g., goats, rabbits, rats, mice). In general, antibodies are produced by immunizing the host (e.g., by injection) with a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide that retains immunogenic properties (which encompasses any portion of the native polypeptide, fragment or oligopeptide). Depending on the host species, various adjuvants can be used to increase the host's immunological response. Such adjuvants include but are not limited to, Freund's, mineral gels (e.g., aluminum hydroxide), and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are potentially useful human adjuvants.

Monoclonal antibodies can be prepared using any technique that provides for the production of antibody molecules by immortalized cell lines in culture. These techniques include, but are not limited to, the hybridoma technique originally described by Koehler and Milstein (1975 Nature 256:495-497), the human B-cell hybridoma technique (Kosbor et al (1983) Immunol Today 4:72; Cote et al (1983) Proc Natl Acad Sci 80:2026-2030) and the EBV-hybridoma technique (Cole et al (1985) Monoclonal Antibodies and Cancer Therapy, Alan R Liss Inc, New York NY, pp 77-96).

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison et al 1984 Proc Natl Acad Sci 81:6851-6855; Neuberger et al 1984 Nature 312:604-608; Takeda et al 1985 Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies that are capsaicin receptor-specific or capsaicin receptor-related polypeptide-specific.

Antibodies can be produced in vivo or by screening recombinant immunoglobulin libraries or panels of highly specific binding reagents as disclosed in Orlandi et al (1989, Proc Natl Acad Sci 86: 3833-3837), and Winter et al.(1991; Nature 349:293-299).

Antibody fragments having specific binding sites for a capsaicin receptor polypeptide or capsaicin receptor-related polypeptide can also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments, which can be produced by pepsin digestion of the antibody molecule, and Fab fragments, which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries can be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse et al 1989 Science 256:1275-1281).

A variety of protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies having established antigen specificities are well known in the art. Such immunoassays typically involve, for example, the formation of complexes between a capsaicin receptor polypeptide and a specific anti-capsaicin receptor antibody, and the detection and quantitation of capsaicin receptor-antibody complex formation. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two noninterfering epitopes on a specific capsaicin receptor protein is preferred, but a competitive binding assay can also be employed. These assays are described in Maddox et.al 1983, J Exp Med 158:1211.

#### Diagnostic Assays Using Capsaicin Receptor-Specific or Capsaicin Receptor-Related Polypeptide-Specific Antibodies

Particular capsaicin receptor antibodies and capsaicin receptor-related polypeptide antibodies are useful for the diagnosis of conditions or diseases characterized by abnormal expression or function of capsaicin receptor (e.g., detection of capsaicin receptor expression in skin to detect neuropathies or in assays to monitor patients having a capsaicin receptor-associated disorder or condition and/or being treated with capsaicin receptor agonists, antagonists, or inhibitors). For example, aberrant carp function might result from over- or under-production of a capsaicin receptor-related polypeptide; thus anti-capsaicin receptor-related antibodies can be used to detect these changes qualitatively or quantitatively. Diagnostic assays for capsaicin receptor or capsaicin receptor-related polypeptide include methods using a detectably-labeled anti-capsaicin receptor antibody or anti-capsaicin receptor-related polypeptide to detect capsaicin receptor in samples (e.g., extracts of cells or tissues). The polypeptides and antibodies of the present invention can be used with or without modification. Frequently, the polypeptides and antibodies are labeled by covalent or noncovalent attachment to a reporter molecule. A wide variety of such suitable reporter molecules are known in the art. Methods for detecting and quantitating antibody binding are well known in the art.

#### Pharmaceutical Compositions Containing Capsaicin Receptor Polypeptides, Capsaicin Receptor-Related Polypeptides, and/or Antibodies Thereto

The present invention also encompasses pharmaceutical compositions that can comprise capsaicin receptor polypeptides, anti-capsaicin receptor polypeptide antibodies, capsaicin receptor-related

polypeptides, or anti-capsaicin receptor-related polypeptides, alone or in combination with at least one other agent, such as a stabilizing compound, which can be administered in any sterile, biocompatible pharmaceutical carrier. The pharmaceutical compositions of the invention can be administered to a patient alone or in combination with other agents, drugs or hormones, in pharmaceutical compositions where it is mixed with excipient(s), or with pharmaceutically acceptable carriers. In one embodiment of the present invention, the pharmaceutically acceptable carrier is pharmaceutically inert.

Capsaicin receptor polypeptides and/or capsaicin receptor-related polypeptides can be administered in order to mitigate the effects of, for example, an endogenous factor that acts as a capsaicin receptor agonist or antagonist or to block or regulate the effects of a capsaicin receptor agonist or antagonist administered to an individual. Anti-capsaicin receptor polypeptide antibodies and/or anti-capsaicin receptor-related polypeptide antibodies can be administered to stimulate a capsaicin receptor in a desired fashion or to block the effects of an endogenous or exogenous capsaicin receptor-binding agonist or antagonist, e.g., via competitive binding to the capsaicin receptor. Pharmaceutical formulations comprising capsaicin receptor polypeptides, capsaicin receptor-related polypeptides, anti-capsaicin receptor antibodies, and/or anti-capsaicin receptor-related polypeptide antibodies can be formulated according to methods known in the art.

#### Transgenic Animals Expressing Polynucleotides Encoding Capsaicin Receptor and/or Capsaicin Receptor-Related Polypeptide-

Nucleic acids encoding capsaicin receptor and/or nucleic acids encoding capsaicin receptor-related polypeptide can be used to generate genetically modified non-human animals or site specific gene modifications in cell lines. The term "transgenic" is intended to encompass genetically modified animals having, for example, a deletion or other knock-out of capsaicin receptor gene activity (and/or capsaicin receptor-related polypeptide activity), an exogenous capsaicin receptor gene (or capsaicin receptor-related polypeptide gene) that is stably transmitted in the host cells, a "knock-in" having altered capsaicin receptor (and/or capsaicin receptor-related polypeptide) gene expression, or an exogenous capsaicin receptor or capsaicin receptor-related polypeptide promoter operably linked to a reporter gene. Of particular interest are homozygous and heterozygous knock-outs and knock-ins of capsaicin receptor and/or capsaicin receptor-related polypeptide function.

Transgenic animals may be made through homologous recombination, where the endogenous capsaicin receptor locus (and/or capsaicin receptor-related polypeptide locus) is altered. Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include plasmids, retroviruses and other animal viruses, YACs, and the like. Of interest are transgenic mammals, preferably a mammal from a genus selected from the group consisting of *Mus* (e.g., mice), *Rattus* (e.g., rats), *Oryctolagus* (e.g., rabbits) and *Mesocricetus* (e.g., hamsters).

A "knock-out" animal is genetically manipulated to substantially reduce, or eliminate endogenous capsaicin receptor function (and/or capsaicin receptor-related polypeptide function). Different approaches may be used to achieve the "knock-out". For example, a chromosomal deletion of all or part of the native



capsaicin receptor homolog (or native capsaicin receptor-related polypeptide homolog) may be induced.

Deletions of the non-coding regions, particularly the promoter region, 3' regulatory sequences, enhancers, or deletions of gene that activate expression of the capsaicin receptor gene and/or the capsaicin receptor-related polypeptide gene. A functional knock-out may also be achieved by the introduction of an anti-sense construct that blocks expression of the native gene(s) (for example, see Li and Cohen (1996) Cell 85:319-329).

Conditional knock-outs of capsaicin receptor gene function (and/or capsaicin receptor-related polypeptide gene function) are also included within the present invention. Conditional knock-outs are transgenic animals that exhibit a defect in capsaicin receptor gene function (and/or capsaicin receptor-related polypeptide gene function) upon exposure of the animal to a substance that promotes target gene alteration, introduction of an enzyme that promotes recombination at the target gene site (e.g., Cre in the Cre-loxP system), or other method for directing the target gene alteration.

For example, a transgenic animal having a conditional knock-out of capsaicin receptor gene function can be produced using the Cre-loxP recombination system (see, e.g., Kilby et al. 1993 Trends Genet 9:413-421). Cre is an enzyme that excises the DNA between two recognition sequences, termed loxP. This system can be used in a variety of ways to create conditional knock-outs of capsaicin receptor. For example, two independent transgenic mice can be produced: one transgenic for an capsaicin receptor sequence flanked by loxP sites and a second transgenic for Cre. The Cre transgene can be under the control of an inducible or developmentally regulated promoter (Gu et al. 1993 Cell 73:1155-1164; Gu et al. 1994 Science 265:103-106), or under control of a tissue-specific or cell type-specific promoter (e.g., a neuron-specific promoter). The capsaicin receptor transgenic is then crossed with the Cre transgenic to produce progeny deficient for the capsaicin receptor gene only in those cells that expressed Cre during development.

Transgenic animals may be made having an exogenous capsaicin receptor gene and/or exogenous capsaicin receptor-related polypeptide gene. The exogenous gene is usually either from a different species than the animal host, or is otherwise altered in its coding or non-coding sequence. The introduced gene may be a wild-type gene, naturally occurring polymorphism, or a genetically manipulated sequence, for example those previously described with deletions, substitutions or insertions in the coding or non-coding regions. The introduced sequence may encode an capsaicin receptor polypeptide and/or capsaicin receptor-related polypeptide. Where the introduced gene is a coding sequence, it is usually operably linked to a promoter, which may be constitutive or inducible, and other regulatory sequences required for expression in the host animal.

Specific constructs of interest include, but are not limited to, anti-sense polynucleotides encoding capsaicin receptor or capsaicin receptor-related polypeptide, or a ribozyme based on a capsaicin receptor or capsaicin receptor-related polypeptide sequence, which will block capsaicin receptor expression or capsaicin receptor-related polypeptide expression, respectively. Such anti-sense polynucleotides or ribozymes will also block expression of dominant negative mutations and over-expression of the corresponding gene. Also of interest is the expression of constructs encoding capsaicin receptor or

capsaicin receptor-related polypeptides in a host where the capsaicin receptor and/or capsaicin receptor-related polypeptide encoded by the construct is derived from a different species than the species of the host in which it is expressed (e.g., expression of human capsaicin receptor in a transgenic mouse). A detectable marker, such as *lac Z* may be introduced into the capsaicin receptor or capsaicin receptor-related polypeptide locus, where upregulation of expression of the corresponding gene will result in an easily detected change in phenotype. Constructs utilizing a promoter region of the capsaicin receptor gene or capsaicin receptor-related polypeptide gene in combination with a reporter gene are also of interest. Constructs having a sequence encoding a truncated or altered (e.g., mutated) capsaicin receptor or capsaicin receptor-related polypeptide are also of interest.

The modified cells or animals are useful in the study of function and regulation of capsaicin receptor and capsaicin receptor-related polypeptide. Such modified cells or animals are also useful in, for example, the study of the function of capsaicin receptor and capsaicin receptor-related polypeptides, as well as the study of the development of nociceptive neurons. Animals may also be used in functional studies, drug screening, etc., e.g. to determine the effect of a candidate drug on capsaicin receptor function or on symptoms associated with disease or conditions associated with capsaicin receptor function (e.g., capsaicin receptor defects or other altered capsaicin receptor activity). By providing expression of capsaicin receptor polypeptide and/or capsaicin receptor-related polypeptide in cells in which it is otherwise not normally produced (e.g., ectopic expression), one can induce changes in cell behavior.

DNA constructs for homologous recombination will comprise at least a portion of the capsaicin receptor gene (or capsaicin receptor-related polypeptide gene) with the desired genetic modification, and will include regions of homology to the target locus. DNA constructs for random integration need not include regions of homology to mediate recombination. Conveniently, markers for positive and negative selection are included. Methods for generating cells having targeted gene modifications through homologous recombination are known in the art. For various techniques for transfecting mammalian cells, see Keown et al. 1990 *Methods in Enzymology* 185:527-537.

For embryonic stem (ES) cells, an ES cell line may be employed, or embryonic cells may be obtained freshly from a host, e.g. mouse, rat, guinea pig, etc. Such cells are grown on an appropriate fibroblast-feeder layer or grown in the presence of appropriate growth factors, such as leukemia inhibiting factor (LIF). When ES cells have been transformed, they may be used to produce transgenic animals. After transformation, the cells are plated onto a feeder layer in an appropriate medium. Cells containing the construct may be detected by employing a selective medium. After sufficient time for colonies to grow, they are picked and analyzed for the occurrence of homologous recombination or integration of the construct. Those colonies that are positive may then be used for embryo manipulation and blastocyst injection. Blastocysts are obtained from 4 to 6 week old superovulated females. The ES cells are trypsinized, and the modified cells are injected into the blastocoel of the blastocyst. After injection, the blastocysts are returned to each uterine horn of pseudopregnant females. Females are then allowed to go to term and the resulting litters screened for mutant cells having the construct. By providing for a different phenotype of the blastocyst and the ES cells, chimeric progeny can be readily detected.

The chimeric animals are screened for the presence of the modified gene. Chimeric animals having the modification (normally chimeric males) are mated with wildtype animals to produce heterozygotes, and the heterozygotes mated to produce homozygotes. If the gene alterations cause lethality at some point in development, tissues or organs can be maintained as allogeneic or congenic grafts or transplants, or in in vitro culture.

Investigation of genetic function may utilize non-mammalian models, particularly using those organisms that are biologically and genetically well-characterized, such as *C. elegans*, *D. melanogaster* and *S. cerevisiae*. For example, transposon (Tc1) insertions in the nematode homolog of a capsaicin receptor gene or a promoter region of a capsaicin receptor gene may be made. The capsaicin receptor gene sequences may be used to knock-out or to complement defined genetic lesions in order to determine the physiological and biochemical pathways involved in function of neuronal cells. It is well known that human genes can complement mutations in lower eukaryotic models.

#### Biosensor Membranes Having Capsaicin Receptor Polypeptides

Due to the responsiveness of capsaicin receptor polypeptides to heat, capsaicin receptor polypeptides can be used in a biosensor for detection of changes in temperature. The biosensor utilizes electrochemical measurement of an ion current across a lipid membrane (or other medium) having a capsaicin receptor polypeptide incorporated therein. Upon stimulation of the capsaicin receptor polypeptide by heat, the capsaicin receptor polypeptide facilitates movement of ions (e.g., calcium) across the membrane, which is then detected as a change in current across the lipid bilayer. The temperature and/or change in temperature can be correlated with the relative increase in conductance across the bilayer due to capsaicin receptor polypeptide activation.

It is well known that amphiphilic molecules can be caused to aggregate in solution to form two or three dimensional ordered arrays such as monolayers, micelles, black lipid membranes, and vesicles or liposomes, which vesicles may have a single compartment or may be of the multilamellar type having a plurality of compartments. The membrane may contain any suitable combination of lipids, long-chain (C12-C24) organic compounds, as well as plastic materials or like polymers for physical reinforcement. Methods and compositions for manufacture of lipid bilayers incorporating a protein of interest, as well as methods and compositions for manufacture of the electrical and mechanical components of biosensors are well known in the art (see, e.g., USPN 5,328,847 (thin membrane sensor with biochemical switch); USPN 5,443,955 (receptor membranes and ionophore gating); USPN 5,234,566 (sensitivity and selectivity of ion channel biosensor membranes); USPN 5,074,977 (digital biosensors and method of using same); and USPN 5,156,810 (biosensors employing electrical, optical, and mechanical signals), each of which are hereby incorporated by reference for manufacture and use of biosensors incorporating a receptor of interest (i.e., capsaicin receptor).

Biosensors according to the present invention comprise at least one lipid membrane, where the membrane includes at least one capsaicin receptor polypeptide. Because capsaicin receptor polypeptides can function when contacted with ligand (e.g., capsaicin) or other effector that mediates capsaicin receptor

activity (e.g., heat), the orientation of capsaicin receptor in the membrane is not substantially important for the function of the biosensor.

Conventional microelectronic configurations will serve adequately to supply power for the sensor, provide a constant direct current voltage across the bilayer prior to heat detection, and measure the ion current surge following capsaicin receptor activation elicited by a change in temperature. It may be additionally desirable to incorporate into the detection electronics a provision for membrane integrity determination, based on the electrical noise accompanying a triggered current signal.

In general, heat is detected using the biosensor of the invention by detecting a change in conductance across the capsaicin receptor polypeptide-containing bilayer. For example, the capsaicin receptor polypeptide lipid bilayer is provided so that a first face of the lipid bilayer (the "heat detection face") is in contact with an buffer solution of neutral pH and containing a selected cation that is capable of transport by the capsaicin receptor (e.g., any cation such as calcium or magnesium, preferably sodium), while a second face of the capsaicin receptor polypeptide lipid bilayer is in contact with a neutral pH buffer having a concentration of the selected cation that is significantly less than the concentration of selected cation in the buffer bathing the heat detection face of the bilayer. Upon exposure of the bilayer's heat detection face to a change in temperature, heat facilitates capsaicin receptor function to provide for transport of the selected cation across the bilayer, resulting in a change in conductance across the bilayer. The change in conductance is then correlated with a change in temperature.

The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

## EXAMPLES

### Example 1: Expression Cloning of Rat Capsaicin Receptor-Encoding DNA

While electrophysiological assays in *Xenopus* oocytes have been employed to obtain cDNAs encoding a variety of cell surface receptors and ion channels (Brake et al. 1995 Nature 371:519-523), this approach proved unsuccessful in identifying a capsaicin receptor clone. A mammalian cell expression cloning strategy based on the ability of capsaicin to trigger an influx of calcium ions into sensory neurons was developed. First, a rodent dorsal root ganglion plasmid cDNA library was constructed in pcDNA3 (Invitrogen) essentially as described (Brake et al., *supra*). A mixture of polyadenylated RNA from newborn (P1) rat and adult mouse dorsal root ganglia was used to generate first-strand cDNA using an oligo (dT) primer containing a Not1 restriction site. Following second strand synthesis and attachment of BstX1 adaptors, the cDNA was digested with Not1. cDNA and BstX1/Not1-linearized pcDNA3 were each purified on potassium acetate gradients, ligated together, and transformed in DH5 $\alpha$  bacteria by electroporation.

The resulting  $2.4 \times 10^6$  independent bacterial clones were divided into 144 pools and stored at -80°C.

HEK 293 (human embryonal kidney) cells constitutively expressing the SV40 large T antigen were maintained in Medium A (DMEM supplemented with 10% fetal bovine serum (Hyclone), penicillin, streptomycin, and L-glutamine) at 37°C, 5% CO<sub>2</sub>. Except where indicated, transient transfections were

carried out with nearly-confluent cells that were replated at  $3.2 \times 10^5$ /35mm tissue culture dish. After 24hrs, the medium was replaced with 1ml Medium B (DMEM supplemented with 10% dialyzed fetal calf serum, penicillin, streptomycin, and L-glutamine). After 2 hrs at 37°C, cells were transfected with 12 µg plasmid DNA using a calcium phosphate precipitation kit (Specialty Media). The following day, cells were rinsed  
5 once with phosphate buffered saline (PBS) containing 1mM EDTA, washed from the plates, collected by centrifugation (200 x g, 5min, 22°C), resuspended in Medium B, and re-plated onto polyornithine-coated coverslips. Under these conditions, each cell acquired plasmids encoding approximately 200 different cDNAs.

Between 6 and 24 hours later, cells were loaded with fura2 (30 min at 37°C) in CIB buffer(mM: 130  
10 NaCl, 3 KCl, 2.5 CaCl<sub>2</sub>, 0.6 MgCl<sub>2</sub>, 1.2 NaHCO<sub>3</sub>, 10 Glucose, 10 Hepes, pH 7.45) containing 10 µM fura-2 acetoxymethyl ester and 0.02% pleuronic acid (Molecular Probes), then rinsed twice with CIB. Ratiometric calcium imaging was performed using a Nikon Diaphot fluorescence microscope equipped with a variable filter wheel (Sutter Instruments) and an intensified CCD camera (Hamamatsu). Dual images (at 340 nm and 380 nm excitation, 510 nm emission) were collected and pseudocolor ratiometric images monitored  
15 during the experiment (Metafluor imaging software, Universal Imaging). Cells were initially imaged in 200 ml CIB, after which 200 ml CIB containing capsaicin at twice the desired concentration was added. Following stimuli, cells were observed for 60-120 s. For each library pool, one microscopic field (300-500 cells) was assayed in each of eight wells.

While cells transfected with most of the assayed pools or with pcDNA3 alone exhibited no  
20 capsaicin responsiveness, 1% to 5% of the cells transfected with one of the cDNA library pools exhibited a profound increase in cytoplasmic calcium concentrations upon capsaicin treatment. cDNA from this pool was further subdivided into smaller pools, and those subpools retransfected into HEK293 cells. In cell populations transfected with some of these subpools, an even larger fraction of cells responded to capsaicin, indicating that a capsaicin receptor-encoding cDNA had been enriched within the population.  
25 The process of pool subdivision and reassay was continued until a single plasmid was isolated that conferred capsaicin-responsiveness upon great than 70% of the transfected cells. The clone that conferred capsaicin-responsiveness contained a 3 kb cDNA insert.

#### Example 2: Sequencing and Characterization of Capsaicin Receptor-Encoding cDNA

30 The 3 kb cDNA insert was sequenced using an automated sequencer (ABI). Homology searches were performed against the nonredundant Genbank database and against an EST database (dbest) using blastn, blastx, and tblastx search programs. Hydrophilicity was calculated using the Hopp-Woods algorithm and a window size of ten 47. The insert was determined to be of rat origin by sequencing an independent cDNA isolated from a rat DRG library and a PCR product derived from mouse DRG cDNA. The sequence  
35 of the isolated rat capsaicin receptor-encoding polynucleotide (SEQ ID NO:1) and its corresponding amino acid sequence (SEQ ID NO:2) are shown in Figure 1. Because a vanilloid moiety constitutes an essential chemical component of capsaicin and resiniferatoxin structures, the proposed site of action of these compounds is more generally referred to as the vanilloid receptor (Szallasi 1994 Gen. Pharmac. 25:223-

243). Accordingly, the newly cloned cDNA was termed VR1, for vanilloid receptor subtype 1. The term "capsaicin receptor" as used here encompasses VR1, but is not limited to VR1.

The VR1-encoding cDNA contains a 2514-nucleotide open reading frame encoding a protein of 838 amino acids with a predicted molecular mass of 95 kD (Figs. 1 and 2A). Hydrophilicity analysis suggests that VR1 is a polytopic protein containing 6 transmembrane domains (noted as "TM" and shaded boxes in Fig. 1 and predicted to be mostly  $\beta$ -sheet (see Fig. 1B)) with an additional short hydrophobic stretch between transmembrane regions 5 and 6 (light shaded region). The amino-terminal hydrophilic segment (432 aa) contains a relatively proline-rich region followed by 3 ankyrin repeat domains (open boxes). The carboxyl-terminus (154 aa) contains no recognizable motifs.

A homology search of protein databases revealed significant similarities between VR1 and members of the family of putative store-operated calcium channels (SOCs) whose prototypical members include the drosophila retinal proteins TRP and TRPL 32, 33 (Fig. 1C). Members of this family have been proposed to mediate the entry of extracellular calcium into cells in response to the depletion of intracellular calcium stores 34. These proteins resemble VR1 with respect to their predicted topological organization and the presence of multiple amino-terminal ankyrin repeats 33. There is also striking amino acid sequence similarity between VR1 and SOCs within and adjacent to the sixth transmembrane region, including the short hydrophobic region between transmembrane domains 5 and 6 that may contribute to the ion permeation path 33. Outside these regions, VR1 shares little sequence similarity with SOCs, suggesting that VR1 is a distant relative of this family of channel proteins. Given the high permeability of VR1 to calcium ions, we nonetheless considered the possibility that it might function as a SOC.

An expressed sequence tag (EST) database homology search revealed several human clones bearing a high degree of similarity to VR1 at both the nucleotide and predicted amino acid levels (Fig. 1C).

Three of these partial cDNAs, independently isolated from different sources, encode sequences in the vicinity of the predicted VR1 pore-loop and sixth transmembrane domains. As shown in Figure 2C, the similarity of one of these clones (hVR, Genbank accession T12251) to the corresponding region of VR1 is extremely high (68% amino acid identity and 84% similarity within the region shown), suggesting that it is likely to be the human VR1 orthologue or a closely related subtype. Human EST clones corresponding to other domains of VR1 show comparable degrees of similarity (not shown) and could represent fragments of the same human transcript.

### Example 3: VR1 Does Not Function as a Store-Operated Calcium Channel (SOC)

The amino acid sequence similarities between VR1 and SOCs suggested that the capsaicin receptor might function as an SOC. To test this, calcium-dependent inward currents were examined in VR1-expressing, intracellular calcium-depleted oocytes according to methods well known in the art.

Briefly, cRNA transcripts were synthesized from NotI-linearized VR1 cDNA templates using T7 RNA polymerase 17. Defolliculated *Xenopus laevis* oocytes were injected with 0.5 - 5ng VR1 cRNA. Four to seven days after injection, two electrode voltage clamp recording was performed ( $E_{\text{hold}} = -60$  mV for  $I_{C_{50}}$  curve and thermal stimulation experiments, and -40 mV for all other experiments) using a Geneclamp 500

amplifier (Axon Instruments) and a MacLab A/D converter (MacLab). The recording chamber was perfused at a rate of 2 ml/min with frog ringer's solution containing (mM) 90 NaCl, 1.0 KCl, 2.4 NaHCO<sub>3</sub>, 0.1 BaCl<sub>2</sub>, 1.0 MgCl<sub>2</sub>, and 10 HEPES, pH 7.6, at room temperature. Prior to performing the store-operated current assays, oocytes were incubated for 1-2 hrs in calcium-free, barium-free frog ringer's solution containing  
5 1 mM EGTA and 1  $\mu$ M Thapsigargin. During voltage clamp recording, these oocytes were intermittently exposed to frog ringer's solution containing 2 mM Ca<sup>2+</sup> and no EGTA to detect calcium-dependent currents (15 second pulses at 2 minute intervals) (Petersen et al. 1995 Biochem J. 311:41-44).

In water-injected control oocytes, a clear depletion-induced current was seen, as previously described (Petersen et al., *supra*). In VR1-expressing oocytes, no quantitative or qualitative differences  
10 were observed in this response (not shown). Moreover, application of SKF 96365 (20  $\mu$ M), an inhibitor of depletion-stimulated calcium entry (Merritt et al. 1990 Biochem J. 271:515-522), had no effect on capsaicin-evoked currents in VR1-expressing oocytes. Thus, VR1 does not appear to be a functional SOC under these circumstances.

#### 15 Example 4: Sensory Neuron-Specific Expression of Capsaicin Receptor

The distribution of VR1 transcripts in neuronal and non-neural rat tissues was assessed by both Northern blot and in situ hybridization analyses. Adult Sprague-Dawley rats were euthanized by asphyxiation in CO<sub>2</sub> and tissues freshly dissected. Poly A<sup>+</sup> RNA was purified either by lysis in guanidinium isothiocyanate followed by purification on oligo-dT cellulose or with the FastTrack kit (Invitrogen).  
20 Approximately 2  $\mu$ g of each sample was electrophoresed through a 0.8% agarose-formaldehyde gel, transferred to a nylon membrane (Hybond N, Amersham), and hybridized with a <sup>32</sup>P-labeled probe representing the entire VR 1 cDNA. Blots were washed at high stringency and autoradiographed. After probing with capsaicin receptor cDNA, the same filters were reprobed with a radiolabeled cyclophilin cDNA fragment as a control (e.g., to correct for relative RNA loading between samples).

For in situ hybridization histochemistry, adult female Sprague-Dawley rats were anesthetized and perfused with 4% paraformaldehyde in PBS. Dorsal root ganglia, trigeminal ganglia and spinal cord were dissected, frozen in liquid N<sub>2</sub>, embedded in OCT mounting medium, and sectioned on a cryostat. Sections (15 micron) were processed and probed at 55°C overnight with a digoxigenin-labeled cRNA generated by in vitro transcription of a 1kb fragment of the VR1 cDNA (nt 1513-2482) using the Genius kit (Boehringer  
30 Mannheim). Sections were developed with alkaline phosphatase-conjugated anti-digoxigenin Fab fragments according to the manufacturer's instructions.

Both Northern blot analysis and in situ hybridization histochemistry indicated that VR1 transcripts are expressed selectively within dorsal root and trigeminal ganglia. An mRNA species of approximately 4 kb was prominently expressed in trigeminal and dorsal root sensory ganglia, both of which have been  
35 shown to contain capsaicin-sensitive neurons. This transcript was absent from all other tissues examined, including spinal cord and brain. A much smaller RNA species (approx. 1.5 kb) was detected in the kidney, but it is unclear whether this transcript could encode a functional VR1 protein.

In situ hybridization to assess the cellular pattern of VR1 expression within sensory ganglia showed that VR1 was expressed predominately within a subset of neurons with small diameters within both dorsal root and trigeminal ganglia. This is in keeping with the observation that most capsaicin-sensitive neurons have relatively small- to medium-sized cell bodies (Holzer 1991 Pharmacol. Rev. 43:143-201; Jansco et al. 1977 Nature 270:741-743). In contrast to the predominant expression of VR1 transcripts in neurons of the dorsal root ganglion, no visible signal was observed in the adjacent spinal cord dorsal horn. While binding sites for radiolabeled resiniferatoxin have been detected in the dorsal horn, these sites are believed to reside on presynaptic terminal that project from primary nociceptors located within adjacent dorsal root ganglia (Holzer 1991, *supra*). The results here support this interpretation.

#### Example 5: Assessment of VR1 Pharmacology in *Xenopus* Oocytes

To compare the pharmacological properties of the cloned capsaicin receptor to those of native vanilloid sites in sensory ganglia, VR1 was expressed in oocyte and used in whole-cell voltage clamp analysis to quantitatively examine its electrophysiological response to a variety of vanilloid agonists (capsaicin, resiniferatoxin) and antagonists (capsazepine). VR1 was expressed in *Xenopus* oocytes as described above (Example 3), except that  $\text{CaCl}_2$  (2 mM) was used in place of  $\text{BaCl}_2$  when generating the capsazepine inhibition curve. The agonists capsaicin and resiniferatoxin were applied sequentially to the same *Xenopus* oocyte expressing VR1. Membrane currents were recorded in the whole cell voltage clamp configuration ( $V_{\text{hold}} = -40\text{mV}$ ).

The results of the capsaicin and resiniferatoxin studies are shown in Figs. 10A-10B. Bars denote duration of agonist application. At negative holding potentials, exposure to capsaicin or resiniferatoxin produced dose-dependent inward current responses in VR1 expressing oocytes, but not in water-injected control cells. As observed in sensory neurons (Winter et al. 1990 Brain Res. 520:131-140; Liu et al. 1994 Proc. Natl. Acad. Sci. USA 91:738-741), capsaicin-evoked current responses returned rapidly to baseline following agonist removal, whereas resiniferatoxin responses often failed to recover, even after a prolonged washout period. Half-maximal effective concentrations for these agonists were within an order of magnitude of those reported for native vanilloid receptors (Oh et al., *supra*; Bevan et al. 1992 Br. J. Pharmacol. 107:544-552), with resiniferatoxin being approximately 20-fold more potent than capsaicin ( $\text{EC}_{50} = 39.1 \text{ nM}$  and  $711.9 \text{ nM}$ , respectively). Hill coefficients derived from these analyses (1.95 and 2.08, respectively) suggest that full activation of the receptor involves the binding of more than one agonist molecule, again consistent with previously described properties of native vanilloid receptors (Oh et al., *supra*; Szallasi 1994 Gen. Pharmac. 25:223-243).

As shown in Figs. 11A and 11B, capsaicin-evoked responses in VR1 expressing oocytes were blocked by the competitive vanilloid receptor antagonist capsazepine at concentrations ( $\text{IC}_{50} = 283.5 \text{ nM}$ ) that inhibit native receptors (Figs. 10A-10B). The current tracing shows that the block of capsaicin activity (cap;  $0.6 \mu\text{M}$ ) by capsazepine (cpz;  $10 \mu\text{M}$ ) is reversible. Another pharmacological signature of vanilloid receptors is their sensitivity to the non-competitive antagonist ruthenium red (RR;  $10 \mu\text{M}$ ), which also blocked capsaicin-evoked responses (cap;  $0.6 \mu\text{M}$ ) in a reversible manner (Figs. 11A). Responses to



resiniferatoxin (50 nM) were also reversibly antagonized by capsazepine (5  $\mu$ M) or ruthenium red (10  $\mu$ M) (not shown).

#### Example 6: Patch Clamp Analysis Of Recombinant Capsaicin Receptors Expressed In HEK293 Embryonal

##### 5 Kidney Cells

The recombinant capsaicin receptor cloned in Example 1 was further characterized in studies using patch clamp analysis in capsaicin receptor-expressing HEK293 cells. HEK293 cells transfected with a control vector (pcDNA3 without a capsaicin receptor-encoding sequence). Patch-clamp recordings were carried out with transiently transfected HEK293 cells at 22°C. Standard bath solution for whole-cell  
10 recordings contained (mM) 140 NaCl, 5 KCl, 2 MgCl<sub>2</sub>, 2 CaCl<sub>2</sub>, 10 HEPES, 10 glucose, pH 7.4 (adjusted with NaOH). In calcium-free bath solution, CaCl<sub>2</sub> was removed and 5 mM EGTA was added.

For monovalent cation substitution experiments, after the whole-cell configuration was obtained in control bath solution, the bath solution was changed to (mM): 140 NaCl (or KCl or CsCl), 10 glucose, and 10 HEPES (adjusted to pH 7.4 with NaOH, KOH or CsOH, respectively) and the reversal potential  
15 measured using voltage-ramps. For divalent cation permeability experiments, the bath solution was changed to (mM) 110 MgCl<sub>2</sub> (or CaCl<sub>2</sub>), 2 Mg(OH)<sub>2</sub> (or Ca(OH)<sub>2</sub>), 10 glucose, 10 HEPES, pH 7.4 (adjusted with HCl).

Bath solution for outside-out patch recordings and pipette solution for inside-out patch recordings contained (mM) 140 NaCl, 10 HEPES, pH 7.4 (adjusted with NaOH). Bath solution for inside-out patch  
20 recordings and pipette solutions for outside-out patch recordings and ion substitution experiments contained: (mM) 140 NaCl, 10 HEPES, 5 EGTA, pH 7.4 (adjusted with NaOH). Pipette solution for whole-cell recordings contained (mM) 140 CsCl (or 130 CsAspartate and 10 NaCl), 5 EGTA, 10 HEPES, pH 7.4 (adjusted with CsOH). Liquid junction potentials were measured directly in separate experiments; they did not exceed 3 mV with solutions used and no correction for this offset was made.

Whole-cell recording data were sampled at 20 kHz and filtered at 5 kHz for analysis (Axopatch 200  
25 amplifier with pCLAMP software, Axon Instruments). Single-channel recording data were sampled at 10 kHz and filtered at 1 kHz. Permeability ratios for monovalent cations to Na ( $P_X/P_{Na}$ ) were calculated as follows:  $P_X/P_{Na} = \exp(\Delta V_{rev}F/RT)$ , where  $V_{rev}$  is the reversal potential, F is Faraday's constant, R is the universal gas constant, and T is absolute temperature. For measurements of divalent permeability,  $P_V/P_{Na}$   
30 was calculated as follows:  $P_V/P_{Na} = [Na^+]_i \exp(\Delta V_{rev}F/RT) / (1 + \exp(\Delta V_{rev}F/RT)) / 4[Y^{2+}]_o$ . Ion activity coefficients of 0.75 (sodium) and 0.25 (calcium or magnesium) were used as correction factors.

Figure 3 show the results of whole cell voltage clamp analysis of capsaicin receptor-expressing HEK293 cells at -60mV using a CsAspartate-filled pipet. These data show an inward cation-specific current which is present only during capsaicin treatment (the time period during which capsaicin was present (1 $\mu$ M)  
35 is indicated by the bar above the plot), and which developed with a short latency upon bath application of capsaicin. No such currents were observed on control, mock-transfected cells. Figure 4 shows the voltage steps (400ms) from -100 mV to +40mV (vertical lines in Figure 3) on an expanded time scale. The currents in the absence of capsaicin (a) were subtracted from the currents obtained in the presence of capsaicin (b).

This analysis of the data revealed a time-independent, receptor-dependent current. In calcium-free medium, the capsaicin-evoked current was also time-independent both at a constant holding potential of -60 mV (Fig. 3) and during voltage steps from -100 to +40 mV (in 20mV increments, Fig. 4). This property enabled characterization of capsaicin-mediated currents under steady-states response conditions in subsequent experiments. Current-voltage relations derived from these data show that such responses exhibit prominent outward rectification resembling that observed in cultured dorsal root ganglion neurons (Fig. 4; Oh et al., *supra*). Because the bath solution used in these experiments consisted mainly of sodium chloride, whereas the patch pipet was filled with cesium aspartate, the observed reversal potential close to 0 mV ( $E_{rev} = 0.5 \pm 0.9$  mV;  $n=13$ ) indicates that the capsaicin-mediated response involves the opening of a cation-selective channel.

In sensory neurons, vanilloid-evoked currents are carried by a mixture of mono- and divalent cations (Bevan et al. 1990 Trends Pharmacol. Sci. 11:330-333; Oh et al., *supra*; Wood et al. 1988 J. Neuroscience 8:3208-3220). This phenomena was examined in VR1-expressing mammalian cells through a series of ion substitution experiments to examine the relative contributions of various cations to capsaicin-evoked currents in VR1-expressing cells. Current-voltage relations established for cells bathed in solutions of differing cationic compositions (Fig. 5; Na<sup>+</sup> (labeled a), K<sup>+</sup> (labeled b), Cs<sup>+</sup> (labeled c), Mg<sup>2+</sup> (labeled d), or Ca<sup>2+</sup> (labeled e) revealed that VR1 does not discriminate among monovalent cations, but exhibits a notable preference for divalent cations. Replacement of extracellular NaCl (140 mM) with equimolar KCl or CsCl did not significantly shift reversal potential ( $E_{rev} = -0.7 \pm 1.2$  mV,  $n=8$ ;  $-1.5$  mV,  $n=9$ ;  $-4.3 \pm 0.9$  mV,  $n=8$ , respectively.  $P_K/P_{Na} = 0.94$ ;  $P_{CS}/P_{Na} = 0.85$ ). Replacement of extracellular NaCl with isotonic (112 mM) MgCl<sub>2</sub> or CaCl<sub>2</sub> shifted  $E_{rev}$  to  $14.4 \pm 1.3$  mV ( $n=3$ ) or  $24.3 \pm 2.3$  mV ( $n=7$ ), respectively. As summarized in Fig. 5, the data thus revealed that the capsaicin receptor-expressing cell membranes had the following relative cation permeabilities for the capsaicin-activated current: Ca<sup>2+</sup> > Mg<sup>2+</sup> >> Na<sup>+</sup> ≈ K<sup>+</sup> ≈ Cs<sup>+</sup>. The very high relative permeability of VR1 to calcium ions ( $P_{Ca}/P_{Na} = 9.60$ ;  $P_{Mg}/P_{Na} = 4.99$ ) exceeds that observed for most non-selective cation channels and is comparable to values reported for NMDA-type glutamate receptors ( $P_{Ca}/P_{Na} = 10.6$ ) (Mayer et al. 1987 J. Physiol. 394:501-527), which are noted for this property. With all bath solutions examined, an outwardly rectifying current-voltage relation was observed, although this feature was somewhat less prominent in MgCl<sub>2</sub>- or CaCl<sub>2</sub>-containing bath solutions.

In cultured sensory neurons, electrophysiological analyses of vanilloid-evoked responses have shown them to be kinetically complex and to desensitize with continuous vanilloid exposure (Liu et al., *supra*; Yeats et al. 1992 J. Physiol. 446:390P). This electrophysiological desensitization (which might underlie aspects of physiological desensitization produced by vanilloids in vivo) appears to depend, in part, upon the presence of extracellular calcium (Yeats et al., *supra*; Holzer 1991 Pharmacol. Rev. 43:143-201). To test the calcium dependency of VR1-expressing cells ability to respond to capsaicin, whole-cell current responses were tested in both calcium-containing standard bath solution and in calcium-free solution (Fig. 6A-F). Capsaicin (1 μM) was applied every 5 min; CsCl was used as pipette solution. The ratios of current size at the end of the third application to the peak of the first application were  $95.3 \pm 2.6\%$  ( $n = 3$ ) in

calcium-free solution, and  $13.0 \pm 4.3\%$  ( $n=5$ ) in calcium-containing solution (T test;  $p < 0.00001$ ). Indeed, in the absence of extracellular calcium, capsaicin-evoked responses in VR1-transfected cells showed little or no desensitization during prolonged agonist application or with successive agonist challenges ( $4.7 \pm 2.3\%$  decrease between first and third applications,  $n=3$ ). In contrast, responses evoked in calcium-containing bath solution consisted of at least two distinct components: one desensitizing ( $87 \pm 4.3\%$  decrease between first and third applications,  $n=5$ ); and one relatively non-desensitizing. Thus, desensitization and multiphasic kinetics of vanilloid-evoked responses can be reproduced outside of a neuronal context and distinguished by their dependence on ambient calcium levels.

The behavior of the VR1 response was also examined in membrane patches excised from transfected cells. Inside-out (I/O) or outside-out (O/O) patches were excised from VR1-transfected cells and analyzed in symmetrical 140 mM NaCl at the indicated holding potentials with capsaicin ( $1 \mu\text{M}$ ) in the bath solution. Dashed lines in the data represented in Fig. 7 indicate closed channel state. In other patches, multiple simultaneous channel openings were observed. The large and well resolved currents of unitary amplitude observed only in the presence of capsaicin (Fig. 7) indicate the existence of capsaicin-gated ion channels within these patches, whose activation does not depend upon soluble cytoplasmic components. The current voltage curve of mean single channel amplitudes ( $\pm$  s.e.m.; Fig. 8), which was calculated from data shown in Fig. 7, also exhibits pronounced outward rectification. The current-voltage relation at the single-channel level was essentially identical to that established in whole-cell configuration, owing to its outward rectification and reversal potential near 0 mV under similar ionic conditions. Unitary conductances of 76.7 pS at positive potentials and 35.4 pS at negative potentials were observed with sodium as the sole charge carrier. These single channel properties mirror those previously described for native vanilloid receptors (Oh et al., *supra*; Forbes et al. 1988 Neurosci. Lett. Suppl. 32:S3).

It has been suggested that the site of vanilloid action may not be confined to the extracellular side of the plasma membrane, reflecting the lipophilic nature of these compounds (James et al. in Capsaicin in the Study of Pain (ed. Wood) Pgs. 83-104 (Academic Press, London, 1993). Interestingly, capsaicin was able to produce identical responses when added to either side of a patch excised from a VR1-expressing cell (Fig. 7), consistent with the notion that vanilloids can permeate or cross the lipid bilayer to mediate their effects. A less likely, but formally consistent explanation is that vanilloid receptors have functionally equivalent capsaicin binding sites on both sides of the plasma membrane.

These data show that the cloned capsaicin receptor behaves in patch clamp analysis in a manner very similar to that reported for wildtype capsaicin receptor.

#### Example 7: Use of Recombinant Capsaicin Receptor to Quantitate Vanilloid Concentrations

As has been recognized for years, the relative pungencies of pepper varieties span an enormously wide range, reflecting, in part, differences in vanilloid content. To-date, methods for rating peppers with respect to their relative "hotness" have relied on subjective psychophysical assays (in which values are reported in Scoville units) (Scoville 1912 J. Am. Pharm. Assoc. 1:453-454) or on biochemical determination of capsaicin content (Woodbury 1980 J. Assoc. Off. Anal. Chem. 63:556-558). To further explore whether

the electrophysiological response of the cloned vanilloid receptor to pepper extracts was in proportion to ability of these peppers to evoke pain.

Several different types of hot peppers (15 g; Thai green, poblano verde, habanero, and wax) were minced and extracted overnight at room temperature with 50 ml absolute ethanol. Soluble extracts were concentrated 15-fold by vacuum desiccation, then diluted 1000-fold in frog ringer's solution for electrophysiological assay. Equivalent fractions (normalized to pepper weight) were tested for their ability to activate the recombinant capsaicin receptor expressed in *Xenopus* oocytes. Capsaicin receptor activation was assessed using a two-electrode voltage-clamp assay to quantitatively measure currents elicited by each pepper extract. Responses were normalized to the response obtained with pure capsaicin (10  $\mu$ M set at 100). The data are shown in Fig. 11 (each value represents an average of four independent determinations, each from separate oocytes; 30 sec application). The relative response of the cloned receptor to the pepper samples and the capsaicin control are shown in the histogram with representative current traces shown to the right of each bar in the histogram. Extracts evoked no response in water-injected cells.

The relative responses of capsaicin receptor to the pepper extracts correlates with the relative hotness of each pepper as determined by conventional analyses and Scoville heat unit assignments. Moreover, the differential "hotness" of these pepper variants, as determined by subjective psychophysical ratings (Berkeley et al. Peppers: A Cookbook pgs. 1-120 (Simon and Schuster, New York, 1992), correlated with their rank order potencies as activators of VR1.

#### Example 8: Capsaicin Induces Death of Cells Expressing VR1

Capsaicin is widely recognized as a neurotoxin that selectively destroys primary afferent nociceptors in vivo and in vitro (Jansco et al. 1977 *Nature* 270:741-743; Wood et al. 1988 *J. Neuroscience* 8:3208-3220). In order to determine whether this selective toxicity solely is a reflection of the specificity of receptor expression, or whether it depends upon additional properties of sensory neurons or their milieu, the ability of capsaicin to kill non-neuronal cells expressing vanilloid receptors was examined in vitro. HEK293 cells were transiently transfected with either vector alone (pcDNA3), vanilloid receptor cDNA diluted 1:50 in pcDNA3 (VR1 1:50) or vanilloid receptor cDNA alone (VR1). Fourteen hours later, culture medium was replaced with medium containing capsaicin (3  $\mu$ M, filled bars) or vehicle (ethanol 0.3%, open bars) (Fig. 12). After seven hours at 37°C, the percentage of dead cells was determined using ethidium homodimer staining. Data represent the mean  $\pm$  s.e.m. of triplicate determinations from a representative experiment. Asterisks indicate a significant difference from ethanol-treated cells (T-test,  $P < 0.0001$ ). Similar results were obtained in three independent experiments.

As shown in Fig. 12, within several hours of continuous exposure to capsaicin, HEK293 cells transfected with VR1 exhibited rampant death, as determined morphologically and by the use of vital stains. In contrast, cells transfected with vector alone were not killed by this treatment. The cell death was characterized by prominent cytoplasmic swelling, coalescence of cytoplasmic contents, and eventual lysis. Thus, VR1 expression in a non-neuronal context can recapitulate the cytotoxicity observed in

vanilloid-treated sensory neurons. Staining with Hoechst dye 33342 revealed no evidence of the nuclear fragmentation often associated with apoptotic cell death 28 (not shown). Together, these observations are consistent with necrotic cell death resulting from excessive ion influx, as has been proposed for vanilloid-induced death of nociceptors (Bevan et al. 1990 Trends Pharmacol. Sci. 11:330-333), glutamate-induced excitotoxicity (Choi 1994 Prog. Brain Res. 199:47-51), and neurodegeneration caused by constitutively activating mutations of various ion channels (Hong et al. 1994 Nature 367:470-473; Hess 1996 Neuron 16:1073-1076).

#### Example 9: Hydrogen Ions Potentiate The Effect Of Capsaicin On VR1

10 A reduction in tissue pH resulting from infection, inflammation, or ischemia can produce pain in mammals. This effect has been attributed to the ability of protons to activate excitatory channels on the surface of nociceptive neurons. A subset of these responses have been reported to share properties in common with those elicited by vanilloids, including similar kinetics, ion selectivity, and antagonism by ruthenium red (Bevan et al. 1994 Trends Neurosci. 17:509-512). Moreover, subthreshold concentrations of hydrogen ions have been shown to potentiate the effects of low concentrations of capsaicin in sensory neurons (Petersen et al. 1993 Pain 54:37-42; Kress et al. 1996 Neurosci. Lett. 211:5-8). It has therefore been proposed that protons might act as endogenous activators or modulators of vanilloid receptors (Bevan et al. 1994 *supra*).

20 To address this possibility, the effects of hydrogen ions on the cloned vanilloid receptor were examined using the oocyte expression system. Capsaicin (0.3  $\mu$ M) was administered throughout the time period tested (spanned by the arrows in Fig. 13) ( $V_{\text{hold}} = -40$  mV). The pH of the bath solution was changed during the experiment (as indicated by the horizontal bars in Fig. 13). VR1-expressing oocytes exhibited no responses to pH 6.3 bath solution without capsaicin; water-injected control oocytes exhibited no responses to either capsaicin or pH 6.3 bath solution (not shown). The current responses obtained from 9 independent VR1-expressing oocytes are summarized in Fig. 14. The grey portion of each bar indicates peak current evoked by capsaicin at pH 7.6, while the black portion represents the additional current evoked by changing the pH to 6.3.

30 Abrupt reduction in bath solution pH (from 7.6 to 5.5) was not sufficient to activate VR1 in the absence of capsaicin (fewer than 10% of VR1-expressing oocytes treated in this way exhibited a large inward current (not shown)), suggesting that hydrogen ions alone cannot efficiently activate this protein. Next, the effect of both capsaicin and pH on VR1 activation was examined. VR1-expressing oocytes were treated with a submaximal concentration of capsaicin (500 nM) at pH 7.6 (Fig. 13). Once their current responses reached a relatively stable plateau, the oocytes were exposed to a solution containing the same concentration of capsaicin at pH 6.3. Under these conditions, the inward current rapidly increased to a new plateau up to five-fold greater in magnitude than the first. Upon returning to pH 7.6, the oocyte response subsided to its initial plateau, and upon the removal of capsaicin it returned to baseline. This potentiation was seen only with sub-saturating concentrations of agonist, as reduced pH did not augment responses

to 10  $\mu$ M capsaicin (not shown). These results suggest that while hydrogen ions alone are not sufficient to activate VR1, they can markedly potentiate capsaicin-evoked responses.

Example 10: The Vanilloid Receptor Is Activated By Noxious Heat

5           The effects of elevated temperature on VR1 activity in calcium influx, conductance, and capsaicin and RR responsivity were examined.

a) Effect of heat upon intracellular calcium

          The effects of heat upon VR1 activity in mediating calcium influx were examined using transfected HEK293 cells and the fluorescent calcium influx screening method described above. Cells were analyzed using microscopic fluorescent calcium imaging before and immediately after the addition of heated calcium imaging buffer (300 ml CIB at 65°C was applied to cells residing in 150 ml CIB at 22°C.) Under these conditions, cells were transiently exposed to a peak temperature of ~45°C.

          While cells transfected with vector alone exhibited only a mild, diffuse change in cytoplasmic free calcium, a large proportion of cells expressing VR1 exhibited a pronounced elevation of calcium levels within seconds of heat treatment. These responses subsided within a few minutes and a subsequent challenge with capsaicin produced a characteristic calcium response, suggesting that the response to heat is a specific signaling event and not a consequence of non-specific membrane perturbation or disruption to cell integrity.

b) Effect of heat upon conductance

20           Whole-cell patch-clamp analysis ( $V_{\text{hold}} = -60$  mV) of VR1-transfected HEK293 cells was performed to examine whether specific heat-evoked membrane currents are associated with this phenomenon. The temperature of the bath medium was raised from 22°C to 48°C (heat) and then restored to 22°C, after which capsaicin (0.5  $\mu$ M) was added to the bath. Ionic conditions were identical to those described for the data in Fig. 2. Voltage-ramps (-100 to +40 mV in 500 ms) were applied before, between, and during responses.

          Exposure of these cells to a rapid increase in temperature (22°C to 50°C in 25 seconds, monitored with an in-bath thermistor) produced large inward currents ( $791 \pm 235$  pA at -60 mV;  $n=9$ ) that were typically similar in amplitude to that evoked by a subsequent application of capsaicin at 500 nM (Fig. 15A). Both heat- and vanilloid-evoked responses showed outward rectification, suggesting that they are mediated by the same entity (Fig. 15B). By comparison, thermally-evoked responses of control, vector-transfected cells were much smaller and exhibited no rectification ( $131 \pm 23$  nA,  $n=8$ ). In addition, the heat response in VR1-transfected cells desensitized during stimulus application, whereas the small thermal response observed in control vector-transfected cells did not. These results suggest that VR1 is acting as a thermal transducer, either by itself or in conjunction with other cellular components.

35           c) Heat activation of VR1 in oocytes

          To determine whether VR1 could mediate similar responses to heat in a different cellular environment, heat activation of VR1 was tested in the oocyte system. Oocytes injected with either VR1 cRNA or water were subjected to two-electrode voltage-clamp ( $V_{\text{hold}} = -60$  mV) while the temperature of

the perfusing buffer was raised from 22.7°C to the level indicated, then held constant for 60 sec. The magnitudes of the resulting inward currents are shown in Fig. 16 as the mean  $\pm$  s.e.m. (VR1, n = 8; water, n=6 independent cells). The asterisk indicates a significant difference from water-injected oocytes (T-test,  $P < 0.0005$ ).

5 In control, water injected oocytes, acute elevation of perfusate temperature produced a small inward current that increased linearly up to 50°C (Fig. 16). VR1 expressing oocytes exhibited similar responses at temperatures up to 40°C, but above this threshold, their responses were significantly larger than those of controls. Thus, even in this non-mammalian context, the VR1-mediated temperature response profile is remarkably consistent with that reported for thermal nociceptors (Fields, *supra*).

#### 10 d) Inhibition of VR1 heat activation by ruthenium red

To further determine whether heat acts specifically through the capsaicin receptor, the ability of ruthenium red to inhibit the heat-mediated response was tested in VR1-expressing oocytes in the system described above. The current tracings shown in Fig. 17 were generated from representative VR1- or water-injected oocytes ( $V_{\text{hold}} = -60$  mV) during successive applications of the indicated stimuli.

15 VR1-injected oocytes exhibited the following mean inward current responses  $\pm$  s.e.m. (n=5): capsaicin (1  $\mu$ M),  $1221 \pm 148$  nA; heat (50 °C),  $2009 \pm 134$  nA; heat plus ruthenium red (10  $\mu$ M),  $243 \pm 47$  nA. Inhibition by ruthenium red was significant ( $88 \pm 2\%$ , n=5; Paired T-test,  $P < 0.0001$ ). No diminution in current response was observed when successive heat pulses were administered in the absence of ruthenium red. Water-injected oocytes showed no response to capsaicin and much smaller responses to heat ( $338 \pm 101$  nA, n=5). Ruthenium red inhibited these responses by only  $21 \pm 26\%$  (n=5; Paired T-test,  $P < 0.1$ ).

20 These data indicate that VR1 is directly involved in this thermal response; application of ruthenium red reduced significantly ( $88 \pm 2\%$ ; n=5) the response of VR1-expressing oocytes to heat, while the smaller response seen in control cells was reduced by only  $21 \pm 26\%$  (n=5) (Fig. 17). Taken together, these observations strongly support the hypothesis that VR1 is activated by noxious, but not innocuous heat.

#### Example 11: Chromosomal Localization And Isolation Of The Mouse VR1 Gene

30 The chromosomal localization of the mouse VR1 gene was determined using fluorescent in situ hybridization (FISH) according to methods well known in the art. Briefly, a bacterial artificial chromosome containing a 90-100 kb insert of mouse genomic DNA encoding portions of the mouse VR1 gene was isolated by PCR analysis and Southern hybridization using rat VR1-derived probes. This insert was labeled with digoxigenin and applied to metaphase spreads of mouse chromosomes. Fluorescently tagged anti-digoxigenin antibodies were then used to visualize the position on the chromosomes to which the VR1 gene hybridized.

35 The VR1 gene mapped to the B3 band of mouse chromosome 11, approximately 49% of the way from the heterochromatic-euchromatic boundary to the telomere of chromosome 11. This region of the mouse chromosome is syntenic to human chromosome 17, particularly the regions 17p11-13, 17pter, and

17qter. It is therefore probable that the human VR1 gene is located within those analogous regions on the human chromosome.

The mouse VR1 gene was sequenced according to methods well known in the art. The nucleotide (SEQ ID NO:10) and amino acid (SEQ ID NO:11) sequences of mouse VR1 are provided in the Sequence Listing below. The rat and mouse VR1 amino acid sequences are more than 95% identical.

#### Example 12: Identification Of Capsaicin Receptor-Related Polypeptide VRRP-1

A Genbank database search using VR1 revealed a number of human and mouse EST sequences similar VR1. Alignment of these EST sequences suggested that all of them, except one (see below) encode identical or very similar genes, suggesting that they represent fragments of the human and mouse versions of the VR1 gene. Over all regions analyzed, the predicted sequences of the encoded human and mouse proteins were highly identical to one another, but only about 50% identical to the rat VR1. Because mouse VR1 protein is more similar to the rat VR1 protein than 50%, we concluded that these EST sequences must encode human and mouse versions of a related protein, which we have termed VRRP-1. VRRP-1 is an example of the capsaicin receptor-related polypeptides encompassed by the present invention.

Portions of the VRRP-1 genes from mouse brain, rat brain, and human CCRF-CEM cells were cloned using PCR primers (GAC CAG CAA GTA CCT CAC (SEQ ID NO:12) and C TCC CAT GCA GCC CAG TTT ACT TCC TCC ACC CTG AAG CAC CAG CGC TCA (SEQ ID NO:13)), which were based on the consensus of human and mouse EST sequences. A full-length rat VRRP-1 cDNA was isolated from a rat brain cDNA library using the rat PCR product as a radiolabeled probe. The rat VRRP-1 cDNA (SEQ ID NO:3; amino acid sequence SEQ ID NO:4)) is approximately 49% identical to rat VR1 at the amino acid level (SEQ ID NOS:2 (rat VR1) and 4 (rat VRRP-1)) and 59% identical at the nucleotide level (SEQ ID NOS:1 (rat VR1) and 3 (rat VRRP-1)).

VRRP-1 does not appear to be activated by capsaicin or heat. Preliminary evidence suggests VRRP-1 may interact with VR1.

#### Example 13: Identification Of Human Capsaicin Receptor VR1

Comparison of VR1 with VRRP-1 and other sequences in the Genbank database suggested that VR1 and VRRP-1 are much more closely related to one another than to any other cloned sequences, with one exception. A single human EST sequence (accession number AA321554; SEQ ID NO:8) obtained from human CCRF-CEM lymphoid cells encodes an amino acid sequence (SEQ ID NO:9) that is at least 71% identical and at least 80% similar to the predicted extreme carboxy terminal domain of the rat VR1 (amino acid residues 774 to 838 of SEQ ID NO:2; see Region 2 of the schematic in Fig. 18). Moreover, rat VR1 (SEQ ID NO:1) and the human EST AA321554 (SEQ ID NO:8) are 75% identical at the nucleotide level. In addition, EST AA321554 contains a stop codon in the same position as the stop codon in rat VR1.

In contrast, although there is homology between a portion of EST AA321554 and the carboxy terminus of rat VRRP-1 (see Region 1 in Fig. 18), the rat VRRP-1 polypeptide is shorter than the rat VR1



polypeptide and the protein from which EST AA321554 appears to be derived. Moreover, even within Region 1, there is greater homology between rat VR1 and EST AA321554 than between either rat VR1 and rat VRRP-1 or than between rat VRRP-1 and EST AA321554. Therefore, the human EST sequence AA321554 represents the human version (ortholog) of rat VR1.

- 5 PCR primers based upon the human sequence were designed to amplify this fragment from cDNA isolated from CCRF-CEM cells or from human sensory ganglion cDNA. The resulting fragment is used as a probe to screen a human genomic DNA library to obtain a full-length human VR1 cDNA sequence from CCRF-CEM cell or human sensory ganglion cDNA. Using the resulting human VR1 genomic fragment, the chromosomal localization of the VR1 gene is confirmed by FISH. The function of the
- 10 polypeptide encoded by the human VR1 gene is confirmed using the functional assays described above.

#### Example 14: Identification Of Human Capsaicin Receptor-Related Polypeptide VRRP-1

- Rat VRRP-1 sequences were used to screen the Genbank database to identify capsaicin receptor-related polypeptides from other organisms. The screen identified several human and mouse EST
- 15 sequences having homology to three separate regions of rat VRRP-1, which regions are termed Regions A, B, and C. Regions A, B, and C represent portions of the VRRP-1 sequence within which the human and mouse VRRP-1-encoding ESTs are clustered, listed from 5' to 3' respectively. Region A encompasses from about residue 580 to about residue 850; Region B encompasses from about nucleotide residue 960 to about residue 1730; and Region C encompasses from about nucleotide residue 1820 to about residue
- 20 2505 in rat the VRRP-1 nucleotide sequence. A summary of the human and mouse EST sequences corresponding to each of these regions is provided in the table below.

**Table: Human and Mouse EST Sequences Corresponding to Rat VRRP-1**

	<b>Rat VRRP-1 Region A</b> (Genbank Accession Nos.)	<b>Rat VRRP-1 Region B</b> (Genbank Accession Nos.)	<b>Rat VRRP-1 Region C</b> (Genbank Accession Nos.)
<b>Human ESTs</b>	H20025, AA236416, H51393, AA236417, H27879, H50364, N21167, AA461295, N26729, H21490, H49060	AA281349, W44731, N23395, W38665, AA357145, N24224	W92895, T12251, AA304033, N35179, AA281348
<b>Mouse ESTs</b>	W82502, W53556	AA139413	AA476107, AA015295, AA274980

- 25 These human and mouse EST sequences can be used to determine a consensus nucleotide sequence for each of Regions A, B, and C. The consensus nucleotide sequence for human VRRP-1 for Region A (SEQ ID NO:5), Region B (SEQ ID NO:6), and Region C (SEQ ID NO:7) are provided in the Sequence Listing below. The consensus sequences can be used to design PCR probes to isolate a fragment encoding the full-length VRRP-1 using methods that are well known in the art.

Example 15. Cloning and Sequencing of a Human VRRP-1

The rat VR1 nucleotide and protein sequences were used to search the genbank databases for related entities. A number of expressed sequence tag (EST) sequences were found that exhibited homology to the rat VR1. These were from human and mouse sources. These sequences were aligned with each other and with rat VR1 and found that all but one of the human sequences appeared to encode the same protein and that this protein was highly homologous to the protein encoded by the mouse sequences. The predicted sequences of these human and mouse proteins were about 50% identical to the rat VR1 protein but much more highly related to one another.

Using the human and mouse EST sequences, PCR primers were designed that were then used on rat brain-derived cDNA to amplify a DNA fragment encoding most of this putative protein. This fragment was radiolabeled and used as a hybridization probe to isolate a full-length cDNA from a rat brain-derived cDNA library. The cDNA (rat VRRP-1) encodes a protein of 761 aa (SEQ ID NO:3) that is 49% identical to the rat VR1 protein and 74% identical to the human VRRP1 protein (SEQ ID NO:23) predicted from the available EST sequences. The rat VRRP-1 mRNA is expressed in sensory ganglia and in other tissues such as brain, spinal cord, spleen, lung, and large intestine.

The human and mouse EST sequences were used to design PCR primers that would allow amplification of the human VRRP-1 sequence from a human-derived cDNA source. Using cDNA derived from human CCRF-CEM cells, a fragment of the human VRRP-1 cDNA was amplified and sequenced, thereby confirming its identity with a subset of the reported EST sequences. Subsequently, PCR primers directed against the 5 prime and 3 prime ends of the predicted human VRRP-1 sequence were used to amplify from CCRF-CEM-derived cDNA a DNA band of approximately 2500bp, which is the correct size for the human cDNA, as predicted by the alignment of the human EST sequences with the rat VRRP-1. The human cDNA was then sequenced using standard methods well known in the art. The DNA sequence of human VRRP-1 (VR2) is provided as SEQ ID NO:35, with the deduced amino acid sequence provided as SEQ ID NO:36.

Example 16. Cloning Chicken VR1 Homologues

Degenerate oligonucleotides were designed based on the amino acid sequence of rat VR1. The oligonucleotides ODJ3885 and ODJ3887 corresponding to VR1 amino acid residues 638-644 and 676-682, respectively, were used as primers for polymerase chain reactions (PCR) with chick genomic DNA as template.

ODJ3885 (SEQ ID NO:28) - 5' TT (TC) AA (AG) TT (TC) AC (GATC) AT (ATC) GG (GATC) ATG

ODJ3887 (SEQ ID NO:29) 5' CAT (GATC) A (GA) (GATC) GC (GAT) AT (GATC) A (GA) CAT (AG) TT

Products of approximately 130 bp resulted, which were isolated and ligated into the vector pT-Adv (Clontech). The inserts in several of these plasmid clones were sequenced. The products from chick genomic DNA fell into two classes: one also corresponding to a very close homologue, and another corresponding to a somewhat more divergent homologue.

40 CVR-PCR1 (SEQ ID NO:30)

TTCAAGTTCACGATTGGGATGGGTGACCTGGATTTTCATGAACATGCCAGATTCAGATACTTTGTCATGCTTCTG  
CTGCTGCTTTTTGTGATCCTCACCTACATCCTTTTGCTCAACATGCTTATAGCCCTTATA

**CVR-PCR2** (SEQ ID NO:31)

- 5 TTCAAGTTCACGATTGGGATGGGAGACCTGGAGTTTACAGAGAACTACAGGTTCAAGTCTGTGTTTGTATCCTT  
TTGGTTCTCTATGTCATCCTTACGTACATCCTCTGCTCAATATGCTTATAGCCCTAATG

A 150 bp EcoRI fragment containing CVR-PCR2 was used as a hybridization probe to screen clones from a cDNA library derived from RNA isolated from chick embryonic dorsal root ganglia (DRG).

- 10 Several hybridizing plasmids were identified. Two of these correspond to a probable chick orthologue of rat VR1. The insert of one of these pCVR2 was sequenced in its entirety (SEQ ID NO:24). The deduced protein sequence (SEQ ID NO:25) shows an amino acid identity to rat VR1 of 67%. Nucleotide alignment of the coding regions of rat and chick VR1 cDNAs also shows 67% identity. Electrophysiological and calcium-imaging analysis of HEK293 cells transfected with pCVR2 indicate that the encoded protein  
15 responds to protons and to high doses of the vanilloid, resiniferatoxin, but not to capsaicin and to heating protocols which activate rat VR1.

**Example 17. Cloning of a Human Vanilloid Receptor**

- A PCR reaction using ODJ3885 and ODJ3887 with human genomic DNA as template produced  
20 a 130bp product. This band was purified and cloned into pT-Adv. The inserts of several clones were sequenced, which showed them all to encode a very close homologue or orthologue of rat VR1. The nucleotide sequence (SEQ ID NO:26) is 91% identical to the corresponding region of rat VR1. The deduced protein sequence over this 45 codon segment is identical to that of rat VR1. Using this new alignment, additional PCR primers were designed to allow amplification of larger segments  
25 of VR1-homologous sequences from human cDNA sources. Primers ODJ4018, corresponding to VR1 amino acid residues 423-429, and ODJ3767, which was derived from the sequence of human EST AA321554, were used in a PCR reaction using as template cDNA from human DRG.

**ODJ4018** (SEQ ID NO:32) 5' TA(TC) TT(TC) AA(TC) TT(TC) TT(TC) GT(GATC) TA 3'

**ODJ3767** (SEQ ID NO:33) 5' AAA AGG GGG ACC AGG GC 3'

- 30 The resulting products were separated by gel electrophoresis transferred to nylon membranes for hybridization with a 150bp probe derived from the previous PCR analysis. A hybridizing fragment of about 1100 bp was thus identified. This is the size expected to be produced by the position of these primers in the rat VR1 sequence.

- The fragment is cloned for sequence analysis. The resulting sequence data is used to design  
35 primers for cloning a full length human cDNA corresponding to this sequence. This will be accomplished using the RACE PCR cloning method [Frohman, M.A. (1993) Methods Enzymol, 218:340-358.]. This may also be carried out using primers derived from the sequence of the small PCR fragment HVR-PCR1.

**Example 18. Cloning and Sequencing of a Human Vanilloid Receptor (VR1)**

- 40 In order to obtain sequences corresponding to the human orthologue of rat VR1, a segment of human genomic DNA was identified which contained sequences present in hVR-PCR1. This genomic DNA was isolated from a library of BAC plasmid clones containing large segments of human genomic DNA

(Shizuya, et al 1992, *Proc Nat Acad of Sci USA*, 89:8794-8797) by Genomic Systems Inc., using oligonucleotide PCR primers derived from the sequence determined for hVR-PCR1. PCR reactions using oligonucleotides ODJ4079 (GGCGACCTGGAGTTCAGTGG (SEQ ID NO:37)) and ODJ4080 (GAGCAGGAGGATGTAGGTGAG (SEQ ID NO:38)) as primers, and human genomic DNA as template  
 5 resulted in the expected 92 bp product. A product of the same size was also obtained using as template cDNA from CCRF-CEM, a human cell line from which the EST sequence #24046 was obtained. This EST sequence appeared to correspond to a close homologue or orthologue of rat VR1. Using these primers to screen a human BAC library by PCR, Genomes Systems provided two BAC plasmid clones, 20614 and 20615.

10 These two clones were further analyzed by restriction digestion and Southern blotting, using the VR1 hybridization probes described above. Inspection of the pattern of restriction fragments using several different restriction endonucleases indicated that these two clones were probably identical. In order to confirm that these plasmids, in fact, contained VR1-related sequences, Southern blots from these digests were hybridized incubated with different VR1 hybridization probes. The blots were first hybridized to a <sup>32</sup>P  
 15 labeled 150 bp EcoRI fragment from hVR-PCR1. A single fragment from each digest hybridized with this probe.

In the case of a PstI digest the hybridizing fragment was approximately 250 bp. The products of PstI digestion of this BAC plasmid were ligated into PstI-digested pBluescriptSK+. The resulting ligation products were used to transform cells of the *E. coli* strain DH5α. Resulting transformants were screened  
 20 by hybridization with the same hVR-PCR1 probe. The insert of one of these clones, hVR1-P1 was sequenced. The results showed that it was highly similar to rat VR1 and corresponded to hVR-PCR1. Alignment of the exon portion of this insert with rVR1 cDNA is shown below.

25	hVR1-P1: 1	ctgcagcttcagatgttcttctgtctctcctgtgcatcttgttgacagtctcaccatgag
	rVR1: 2180	ctgcagcttcagatgttcttctgtctctcctgtgcaatcttgttgacaggtctcaccatgag
30	hVR1-P1: 61	ggcgatgagcatgttgagcaggaggatgtaggtgagaattacataggccagcagcaggat
	rVR1: 2120	agcaatgagcatgttgagcagaaggatgtaggtgagaatcacataggccagtaacaggat
35	hVR1-P1: 121	gatgaagacagccttgaagtcatagttctcagtgaactccaggctcgcccatgccgatggt
	rVR1: 2060	gatgaagacagccttgaagtcgtagttctcagtgaactccaggctcgcccatgccgatggt
40	hVR1-P1: 181	gaacttgaacagctccaggcaggtggagtacaggctgtttaggag (SEQ ID NO:39)
	rVR1: 2000	gaacttgaacagctccagacatgtggaatacaggctgtttaggag (SEQ ID NO:40)

In order to further localize VR1-related sequences on this BAC plasmid insert, Southern blots were performed using a 1008 bp NheI fragment from the rVR1 cDNA as a hybridization probe. This  
 45 fragment includes almost the entire 5' portion of the rVR1 coding sequence. In this case each digest produced one or more fragments that hybridized strongly with this probe. In particular, HindIII digestion

5

15

30

35

40

45

50

HVR1-B2: 14 MKKWSSTD LGAAADPLQKDTCPDPLDGD P NSRPPFAKPQLSTAKSRTLFGKGDSEEAFF  
 M++ +S D + P Q+++C DP D DPN +PPP KP + T +SRTLFGKGDSEEA P  
 rVR1: 1 MEQRASLDSEESPPQENSCLDPPORDPNCKPPPVKPHIFTTSRTRLFGKGDSEEA SP  
 HVR1-B2: 194 VDCPHEEGELDSCPTITVSPVITIQRPRXRP 286 (SEQ ID NO:45)  
 +DCP+EEG L SCP ITVS V+TIQRP P  
 rVR1: 61 LDCPYEEGGLASCPITVSSVLTIQRPGDGP 91 (SEQ ID NO:46)

Using this sequence information, two primers were designed to allow production of a human VR1 cDNA from using RT-PCR from polyA+ RNA isolated from the CCRF-CEM cell line. The two primers ODJ4157 (AGAAATGGAGCAGCACAGACTTGG (SEQ ID NO:47)) and ODJ4162 (TCACTTCTCCCCGGAAGCGGCAG (SEQ ID NO:48)) were used as primers in a PCR reaction with CCRF-CEM cDNA as template. A product of about 2500 bp resulted from this reaction. Southern blot analysis of this product using a hVR-PCR1 hybridization probe, indicated that this product was, in fact, VR1-related. The product was purified by preparative agarose gel electrophoresis and subcloned into the vector pT-Adv (Clontech). Several clones were isolated and four of these were subjected to DNA sequence analysis. The resulting DNA sequence of human VR1 is provided as SEQ ID NO:33, with the deduced amino acid sequence provided as SEQ ID NO:34.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Before the present nucleotide and polypeptide sequences are described, it is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors and reagents described as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a host cell" includes a plurality of such host cells and reference to "the antibody" includes reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs. Although any methods, devices and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, the preferred methods, devices and materials are now described.

- All publications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing, for example, the cell lines, vectors, and methodologies which are described in the publications which might be used in connection with the presently described invention. The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior invention.
- 5

## CLAIMS

1. An isolated capsaicin receptor polypeptide.
- 5 2. The capsaicin receptor polypeptide of Claim 1, wherein the polypeptide is a VR1 subtype.
3. The capsaicin receptor polypeptide of Claim 1, wherein the polypeptide is a VR2 subtype.
4. The capsaicin receptor polypeptide of Claim 1, wherein the polypeptide comprises an  
10 amino acid sequence selected from the group consisting of SEQ ID NO:2, 4, 9, 11, 23, 25, 27, 34, or 36.
5. An isolated polynucleotide sequence encoding a capsaicin receptor polypeptide of claim  
1.
- 15 6. The polynucleotide sequence of Claim 5, wherein the sequence encodes a VR1 subtype.
7. The polynucleotide sequence of Claim 5, wherein the sequence encodes a VR2 subtype.
8. The isolated polynucleotide sequence of Claim 5, wherein the sequence comprises a  
20 sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 6, 7, 8, 10, 20, 21, 22, 24, 26, 33, or 35.
9. A recombinant expression vector comprising the polynucleotide sequence of Claim 5.
- 25 10. A recombinant host cell containing the polynucleotide sequence of Claim 5.
11. A method for producing the capsaicin receptor polypeptide of Claim 1, the method comprising the steps of:  
30 a) culturing a recombinant host cell containing a capsaicin receptor polypeptide-encoding polynucleotide sequence under conditions suitable for the expression of the polypeptide; and  
b) recovering the polypeptide from the host cell culture.
12. An isolated antibody that specifically binds a capsaicin receptor polypeptide of claim 1.
- 35 13. A method for identifying compounds that bind a capsaicin receptor polypeptide, the method comprising the steps of:  
contacting a capsaicin receptor polypeptide with a test compound; and  
detecting specific binding of the test compound to the capsaicin receptor polypeptide.



14. The method of claim 13, wherein said detecting is by detecting of an alteration of intracellular calcium concentration in the capsaicin receptor-expressing host cell.

15. A method for detecting a vanilloid compound in a sample, wherein the vanilloid compound  
5 has activity in binding a capsaicin receptor polypeptide, the method comprising the steps of:  
contacting a sample suspected of containing a vanilloid compound with a eukaryotic host cell  
expressing a capsaicin receptor polypeptide;  
detecting an alteration of a cellular response associated with capsaicin receptor activity in the  
capsaicin receptor-expressing host cell.

10

16. The method of claim 15, wherein the cellular response associated with capsaicin receptor activity is an increase in intracellular calcium concentration.

17. A pharmaceutical composition comprising a substantially purified capsaicin receptor  
15 polypeptide and a suitable pharmaceutical carrier.

18. A non-human transgenic animal model for capsaicin receptor gene function wherein the transgenic animal is characterized by having an alteration in capsaicin receptor function relative to a normal animal of the same species.

20

19. A method of screening for biologically active agents that modulate capsaicin receptor function, the method comprising:

combining a candidate agent with any one of:

- 25 (a) a mammalian capsaicin receptor polypeptide;  
(b) a mammalian capsaicin receptor-related polypeptide;  
(c) a cell containing a nucleic acid encoding a mammalian capsaicin receptor polypeptide;  
(d) a cell containing a nucleic acid encoding a mammalian capsaicin receptor-related polypeptide;

or

- 30 (e) a non-human transgenic animal model for function of a capsaicin receptor gene comprising one of: (i) a knockout of a capsaicin receptor gene; (ii) an exogenous and stably transmitted mammalian capsaicin receptor gene sequence; or (iii) a capsaicin receptor promoter sequence operably linked to a reporter gene; and

determining the effect of said agent on capsaicin receptor function.

MEQASLDSESESPQENSCLDPPDRDPNCKPPPVKPHIFTTRSR 46  
 TRLFKGDSSEASPLDCPYEEGGLASCPIITVSSVLTIQRPDGPAS 93  
 VRPSSQDSVSAGEKPPRLYDRRSIFDAVAQSNQCQELLESLLPFLQRSK 140  
 KRLTDSSEFKDPE<sup>o</sup>TGKTCCLKAMNLHNHGQNDTIALLLDVARKTDSLK 187  
 QFVNASYTDSYY[KGQTALHIAIERRM<sup>o</sup>TLVTLLEN<sup>o</sup>CA<sup>o</sup>DVQAAN<sup>o</sup>GD 234  
 FFKKTKGRPGFY[FGELPLSLAACTNQLAIVKFL<sup>o</sup>LQNSWQPADISAR<sup>o</sup>D 281  
 SVGNTVLHALVEVADNTVDNTKFVTSMYNEILILGAKLHPTLKLEEIT 329  
 NRKGLTPLALAASSGKIGVLAYILQREIHEPECRH<sup>o</sup>LSRKFT<sup>o</sup>EWAYGP 376  
 VHSSLYDLSCIDTCEKNSVLEVIAYSSSETPNRHDMLLVEPLNRL<sup>o</sup>LQ 423  
 DKWDRFVKR[IFYFNFFVYCLYMIIFTAAAYY]RPVEGLPPYK<sup>o</sup>LKNTVG 470  
 DYFRVTGEI[LSVSGGVYFFFRGIGY]FLQRRPSLKSLFVD[SYSEILFFV] 518  
 QSLFMLVSVVLYF[SQRKEYVASMV]FSLAMGW<sup>o</sup>TNMLYYTRCGFQQMGI 564  
 YAVMI[EKMILRD]LCRFMFVYLVFLFCFSTAVV<sup>o</sup>TLIEDGKNNSLPMEST 612  
 PHKCRGSACKP[GNSYNSLYSTCLEL<sup>o</sup>FKFTIGMG]DLEFTENYD[FKAVF] 659  
 IILLAYVILTYILLNMLIALMG<sup>o</sup>ETVNKIAQESKNIWKLQRAITILDTE 709  
 KSFLKCMRKAFRSGKLLQVGFTPDGKDDYRWCFRVDEVNWT<sup>o</sup>TWNT 754  
 NVGIINEDPGNCEGVKRTL<sup>o</sup>SFSLRSGRVSGRNWKNFALVPLLRDAST 801  
 RDRHATQQEEVQLKHYTGSLKPEDAEVFKDSMVPCEK SEQ ID NO: 2 838

FIG. 1A




FIG. 1B

FIG. 1C

636	ELFKFTIGMGDL...EFTENYDFKAVFILLLLAYVILLT	670
ELFKFTIGMGEL...AFQEQLHFRGMVLLLLAYVLLT		
SLFWASFGGLVDSLVSFDLAGIKSFFTRMFWALLMFGCSYSVIN		
SLFWASFGGLVDSLVSFDLAGIKSFFTRMFWALLMFGCSYSVIN		
SLFWASIFGLINLLYVTNVKAKQHEFFTEFVCGATMFGCTYNVIS		
RTFIMTIGFESVLYREMSACDNFWMMKWIGKLIFFVIFETFV		
671	YILLNMLIALMGETVKNKIAQESKNIWKLRRAITIL	706
YILLNMLIALMSETVNSVATID		
IIVLLNMLIAMMSNSYQIISERADVEWKFAFSQLWM		
IIVLLNMLIAMMSNSYQIISERADTEWKFAFSQLWM		
LIVLLNMLIAMMNNNSYQLIADHADIETWKFAFSQLWM		
SILLQFNILLIAMMTRTYETIFFL		

rat VR1

human T12251

Caliphora z80230

Drosophila TRP

Bovine x99792

C. Elegans z72508

rat VR1

human T12251

Caliphora z80230

Drosophila TRP

Bovine x99792

C. Elegans z72508

(SEQ ID NO: 14)

(SEQ ID NO: 15)



(SEQ ID NO: 16)

(SEQ ID NO: 17)



(SEQ ID NO: 18)

(SEQ ID NO: 19)

3/12

capsaicin 1  $\mu\text{M}$ **FIG. 2****FIG. 3**

4/12

**FIG. 4****FIG. 5**

5/12

capsaicin 1  $\mu$ M




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D





FIG. 6E



FIG. 6F



Ca(-)

Ca(+)

6/12




FIG. 8






FIG. 7

7/12


**FIG. 9A**




**FIG. 10A**



**FIG. 9B**




**FIG. 10B**






8/12

FIG. 11



9/12

**FIG. 12**

10/12

FIG. 14





FIG. 13



11/12

FIG. 15A




FIG. 15B




FIG. 16






FIG. 17



12/12

FIG. 18



-1-

## SEQUENCE LISTING

<110> Julius, David J.  
Caterina, Michael J.  
Brake, Anthony J.

<120> NUCLEIC ACID SEQUENCES ENCODING  
CAPSAICIN RECEPTOR AND CAPSAICIN RECEPTOR-RELATED  
POLYPEPTIDES AND USES THEREOF

<130> 9076/084PCT

<140> Unassigned

<141> 1999-01-22

<150> 60/072,151

<151> 1998-01-22

<160> 48

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 2880

<212> DNA

<213> R. rattus

<220>

<221> CDS

<222> (81)...(2594)

<223> VR1 capsaicin receptor

<400> 1

```

cagctccaag gcacttgctc catttgggggt gtgcctgcac ctagctgggtt gcaaattggg      60
ccacagagga tctggaaagg atg gaa caa cgg gct agc tta gac tca gag gag      113
               Met Glu Gln Arg Ala Ser Leu Asp Ser Glu Glu
                   1               5               10

tct gag tcc cca ccc caa gag aac tcc tgc ctg gac cct cca gac aga      161
Ser Glu Ser Pro Pro Gln Glu Asn Ser Cys Leu Asp Pro Pro Asp Arg
                   15               20               25

gac cct aac tgc aag cca cct cca gtc aag ccc cac atc ttc act acc      209
Asp Pro Asn Cys Lys Pro Pro Pro Val Lys Pro His Ile Phe Thr Thr
                   30               35               40

agg agt cgt acc cgg ctt ttt ggg aag ggt gac tcg gag gag gcc tct      257
Arg Ser Arg Thr Arg Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Ser
                   45               50               55

ccc ctg gac tgc cct tat gag gaa ggc ggg ctg gct tcc tgc cct atc      305
Pro Leu Asp Cys Pro Tyr Glu Glu Gly Gly Leu Ala Ser Cys Pro Ile
                   60               65               70               75

```

-2-

atc act gtc agc tct gtt cta act atc cag agg cct ggg gat gga cct Ile Thr Val Ser Ser Val Leu Thr Ile Gln Arg Pro Gly Asp Gly Pro 80 85 90	353
gcc agt gtc agg ccg tca tcc cag gac tcc gtc tcc gct ggt gag aag Ala Ser Val Arg Pro Ser Ser Gln Asp Ser Val Ser Ala Gly Glu Lys 95 100 105	401
ccc ccg agg ctc tat gat cgc agg agc atc ttc gat gct gtg gct cag Pro Pro Arg Leu Tyr Asp Arg Arg Ser Ile Phe Asp Ala Val Ala Gln 110 115 120	449
agt aac tgc cag gag ctg gag agc ctg ctg ccc ttc ctg cag agg agc Ser Asn Cys Gln Glu Leu Glu Ser Leu Leu Pro Phe Leu Gln Arg Ser 125 130 135	497
aag aag cgc ctg act gac agc gag ttc aaa gac cca gag aca gga aag Lys Lys Arg Leu Thr Asp Ser Glu Phe Lys Asp Pro Glu Thr Gly Lys 140 145 150 155	545
acc tgt ctg cta aaa gcc atg ctc aat ctg cac aat ggg cag aat gac Thr Cys Leu Leu Lys Ala Met Leu Asn Leu His Asn Gly Gln Asn Asp 160 165 170	593
acc atc gct ctg ctc ctg gac gtt gcc cgg aag aca gac agc ctg aag Thr Ile Ala Leu Leu Leu Asp Val Ala Arg Lys Thr Asp Ser Leu Lys 175 180 185	641
cag ttt gtc aat gcc agc tac aca gac agc tac tac aag ggc cag aca Gln Phe Val Asn Ala Ser Tyr Thr Asp Ser Tyr Tyr Lys Gly Gln Thr 190 195 200	689
gca ctg cac att gcc att gaa cgg cgg aac atg acg ctg gtg acc ctc Ala Leu His Ile Ala Ile Glu Arg Arg Asn Met Thr Leu Val Thr Leu 205 210 215	737
ttg gtg gag aat gga gca gat gtc cag gct gcg gct aac ggg gac ttc Leu Val Glu Asn Gly Ala Asp Val Gln Ala Ala Ala Asn Gly Asp Phe 220 225 230 235	785
ttc aag aaa acc aaa ggg agg cct ggc ttc tac ttt ggt gag ctg ccc Phe Lys Lys Thr Lys Gly Arg Pro Gly Phe Tyr Phe Gly Glu Leu Pro 240 245 250	833
ctg tcc ctg gct gcg tgc acc aac cag ctg gcc att gtg aag ttc ctg Leu Ser Leu Ala Ala Cys Thr Asn Gln Leu Ala Ile Val Lys Phe Leu 255 260 265	881
ctg cag aac tcc tgg cag cct gca gac atc agc gcc cgg gac tca gtg Leu Gln Asn Ser Trp Gln Pro Ala Asp Ile Ser Ala Arg Asp Ser Val 270 275 280	929

-3-

ggc aac acg gtg ctt cat gcc ctg gtg gag gtg gca gat aac aca gtt	977
Gly Asn Thr Val Leu His Ala Leu Val Glu Val Ala Asp Asn Thr Val	
285 290 295	
gac aac acc aag ttc gtg aca agc atg tac aac gag atc ttg atc ctg	1025
Asp Asn Thr Lys Phe Val Thr Ser Met Tyr Asn Glu Ile Leu Ile Leu	
300 305 310 315	
ggg gcc aaa ctc cac ccc acg ctg aag ctg gaa gag atc acc aac agg	1073
Gly Ala Lys Leu His Pro Thr Leu Lys Leu Glu Glu Ile Thr Asn Arg	
320 325 330	
aag ggg ctc acg cca ctg gct ctg gct gct agc agt ggg aag atc ggg	1121
Lys Gly Leu Thr Pro Leu Ala Leu Ala Ala Ser Ser Gly Lys Ile Gly	
335 340 345	
gtc ttg gcc tac att ctc cag agg gag atc cat gaa ccc gag tgc cga	1169
Val Leu Ala Tyr Ile Leu Gln Arg Glu Ile His Glu Pro Glu Cys Arg	
350 355 360	
cac cta tcc agg aag ttc acc gaa tgg gcc tat ggg cca gtg cac tcc	1217
His Leu Ser Arg Lys Phe Thr Glu Trp Ala Tyr Gly Pro Val His Ser	
365 370 375	
tcc ctt tat gac ctg tcc tgc att gac acc tgt gaa aag aac tcg gtt	1265
Ser Leu Tyr Asp Leu Ser Cys Ile Asp Thr Cys Glu Lys Asn Ser Val	
380 385 390 395	
ctg gag gtg atc gct tac agc agc agt gag acc cct aac cgt cat gac	1313
Leu Glu Val Ile Ala Tyr Ser Ser Ser Glu Thr Pro Asn Arg His Asp	
400 405 410	
atg ctt ctc gtg gaa ccc ttg aac cga ctc cta cag gac aag tgg gac	1361
Met Leu Leu Val Glu Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp Asp	
415 420 425	
aga ttt gtc aag cgc atc ttc tac ttc aac ttc ttc gtc tac tgc ttg	1409
Arg Phe Val Lys Arg Ile Phe Tyr Phe Asn Phe Phe Val Tyr Cys Leu	
430 435 440	
tat atg atc atc ttc acc gcg gct gcc tac tat cgg cct gtg gaa ggc	1457
Tyr Met Ile Ile Phe Thr Ala Ala Ala Tyr Tyr Arg Pro Val Glu Gly	
445 450 455	
ttg ccc ccc tat aag ctg aaa aac acc gtt ggg gac tat ttc cga gtc	1505
Leu Pro Pro Tyr Lys Leu Lys Asn Thr Val Gly Asp Tyr Phe Arg Val	
460 465 470 475	
acc gga gag atc ttg tct gtg tca gga gga gtc tac ttc ttc ttc cga	1553
Thr Gly Glu Ile Leu Ser Val Ser Gly Gly Val Tyr Phe Phe Phe Arg	
480 485 490	
ggg att caa tat ttc ctg cag agg cga cca tcc ctc aag agt ttg ttt	1601
Gly Ile Gln Tyr Phe Leu Gln Arg Arg Pro Ser Leu Lys Ser Leu Phe	
495 500 505	



- 4 -

gtg gac agc tac agt gag ata ctt ttc ttt gta cag tcg ctg ttc atg Val Asp Ser Tyr Ser Glu Ile Leu Phe Phe Val Gln Ser Leu Phe Met 510 515 520	1649
ctg gtg tct gtg gta ctg tac ttc agc caa cgc aag gag tat gtg gct Leu Val Ser Val Val Leu Tyr Phe Ser Gln Arg Lys Glu Tyr Val Ala 525 530 535	1697
tcc atg gtg ttc tcc ctg gcc atg ggc tgg acc aac atg ctc tac tat Ser Met Val Phe Ser Leu Ala Met Gly Trp Thr Asn Met Leu Tyr Tyr 540 545 550 555	1745
acc cga gga ttc cag cag atg ggc atc tat gct gtc atg att gag aag Thr Arg Gly Phe Gln Gln Met Gly Ile Tyr Ala Val Met Ile Glu Lys 560 565 570	1793
atg atc ctc aga gac ctg tgc cgg ttt atg ttc gtc tac ctc gtg ttc Met Ile Leu Arg Asp Leu Cys Arg Phe Met Phe Val Tyr Leu Val Phe 575 580 585	1841
ttg ttt gga ttt tcc aca gct gtg gtg aca ctg att gag gat ggg aag Leu Phe Gly Phe Ser Thr Ala Val Val Thr Leu Ile Glu Asp Gly Lys 590 595 600	1889
aat aac tct ctg cct atg gag tcc aca cca cac aag tgc cgg ggg tct Asn Asn Ser Leu Pro Met Glu Ser Thr Pro His Lys Cys Arg Gly Ser 605 610 615	1937
gcc tgc aag cca ggt aac tct tac aac agc ctg tat tcc aca tgt ctg Ala Cys Lys Pro Gly Asn Ser Tyr Asn Ser Leu Tyr Ser Thr Cys Leu 620 625 630 635	1985
gag ctg ttc aag ttc acc atc ggc atg ggc gac ctg gag ttc act gag Glu Leu Phe Lys Phe Thr Ile Gly Met Gly Asp Leu Glu Phe Thr Glu 640 645 650	2033
aac tac gac ttc aag gct gtc ttc atc atc ctg tta ctg gcc tat gtg Asn Tyr Asp Phe Lys Ala Val Phe Ile Ile Leu Leu Leu Ala Tyr Val 655 660 665	2081
att ctc acc tac atc ctt ctg ctc aac atg ctc att gct ctc atg ggt Ile Leu Thr Tyr Ile Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly 670 675 680	2129
gag acc gtc aac aag att gca caa gag agc aag aac atc tgg aag ctg Glu Thr Val Asn Lys Ile Ala Gln Glu Ser Lys Asn Ile Trp Lys Leu 685 690 695	2177
cag aga gcc atc acc atc ctg gat aca gag aag agc ttc ctg aag tgc Gln Arg Ala Ile Thr Ile Leu Asp Thr Glu Lys Ser Phe Leu Lys Cys 700 705 710 715	2225

- 5 -

atg agg aag gcc ttc cgc tct ggc aag ctg ctg cag gtg ggg ttc act 2273  
 Met Arg Lys Ala Phe Arg Ser Gly Lys Leu Leu Gln Val Gly Phe Thr  
 720 725 730

cct gac ggc aag gat gac tac cgg tgg tgt ttc agg gtg gac gag gta 2321  
 Pro Asp Gly Lys Asp Asp Tyr Arg Trp Cys Phe Arg Val Asp Glu Val  
 735 740 745

aac tgg act acc tgg aac acc aat gtg ggt atc atc aac gag gac cca 2369  
 Asn Trp Thr Thr Trp Asn Thr Asn Val Gly Ile Ile Asn Glu Asp Pro  
 750 755 760

ggc aac tgt gag ggc gtc aag cgc acc ctg agc ttc tcc ctg agg tca 2417  
 Gly Asn Cys Glu Gly Val Lys Arg Thr Leu Ser Phe Ser Leu Arg Ser  
 765 770 775

ggc cga gtt tca ggg aga aac tgg aag aac ttt gcc ctg gtt ccc ctt 2465  
 Gly Arg Val Ser Gly Arg Asn Trp Lys Asn Phe Ala Leu Val Pro Leu  
 780 785 790 795

ctg agg gat gca agc act cga gat aga cat gcc acc cag cag gaa gaa 2513  
 Leu Arg Asp Ala Ser Thr Arg Asp Arg His Ala Thr Gln Gln Glu Glu  
 800 805 810

gtt caa ctg aag cat tat acg gga tcc ctt aag cca gag gat gct gag 2561  
 Val Gln Leu Lys His Tyr Thr Gly Ser Leu Lys Pro Glu Asp Ala Glu  
 815 820 825

gtt ttc aag gat tcc atg gtc cca ggg gag aaa taatggacac tatgcaggga 2614  
 Val Phe Lys Asp Ser Met Val Pro Gly Glu Lys  
 830 835

tcaatgcggg gtctttgggt ggtctgctta gggaaccagc aggggttgacg ttatctgggt 2674  
 ccactctgtg cctgcctagg cacattccta ggacttcggc gggcctgctg tgggaactgg 2734  
 gaggtgtgtg ggaattgaga tgtgtatcca accatgatct ccaaacattt ggctttcaac 2794  
 tctttatgga ctttattaaa cagagtgaat ggcaaatctc tacttggaca cataaaaaaa 2854  
 aaaaaaaaaa aaaaaaaaaa aaaaaa 2880

&lt;210&gt; 2

&lt;211&gt; 838

&lt;212&gt; PRT

&lt;213&gt; R. rattus

&lt;400&gt; 2

Met Glu Gln Arg Ala Ser Leu Asp Ser Glu Glu Ser Glu Ser Pro Pro  
 1 5 10 15  
 Gln Glu Asn Ser Cys Leu Asp Pro Pro Asp Arg Asp Pro Asn Cys Lys  
 20 25 30  
 Pro Pro Pro Val Lys Pro His Ile Phe Thr Thr Arg Ser Arg Thr Arg  
 35 40 45  
 Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Ser Pro Leu Asp Cys Pro  
 50 55 60  
 Tyr Glu Glu Gly Gly Leu Ala Ser Cys Pro Ile Ile Thr Val Ser Ser  
 65 70 75 80

-6-

Val Leu Thr Ile Gln Arg Pro Gly Asp Gly Pro Ala Ser Val Arg Pro  
 85 90 95  
 Ser Ser Gln Asp Ser Val Ser Ala Gly Glu Lys Pro Pro Arg Leu Tyr  
 100 105 110  
 Asp Arg Arg Ser Ile Phe Asp Ala Val Ala Gln Ser Asn Cys Gln Glu  
 115 120 125  
 Leu Glu Ser Leu Leu Pro Phe Leu Gln Arg Ser Lys Lys Arg Leu Thr  
 130 135 140  
 Asp Ser Glu Phe Lys Asp Pro Glu Thr Gly Lys Thr Cys Leu Leu Lys  
 145 150 155 160  
 Ala Met Leu Asn Leu His Asn Gly Gln Asn Asp Thr Ile Ala Leu Leu  
 165 170 175  
 Leu Asp Val Ala Arg Lys Thr Asp Ser Leu Lys Gln Phe Val Asn Ala  
 180 185 190  
 Ser Tyr Thr Asp Ser Tyr Tyr Lys Gly Gln Thr Ala Leu His Ile Ala  
 195 200 205  
 Ile Glu Arg Arg Asn Met Thr Leu Val Thr Leu Leu Val Glu Asn Gly  
 210 215 220  
 Ala Asp Val Gln Ala Ala Asn Gly Asp Phe Phe Lys Lys Thr Lys  
 225 230 235 240  
 Gly Arg Pro Gly Phe Tyr Phe Gly Glu Leu Pro Leu Ser Leu Ala Ala  
 245 250 255  
 Cys Thr Asn Gln Leu Ala Ile Val Lys Phe Leu Leu Gln Asn Ser Trp  
 260 265 270  
 Gln Pro Ala Asp Ile Ser Ala Arg Asp Ser Val Gly Asn Thr Val Leu  
 275 280 285  
 His Ala Leu Val Glu Val Ala Asp Asn Thr Val Asp Asn Thr Lys Phe  
 290 295 300  
 Val Thr Ser Met Tyr Asn Glu Ile Leu Ile Leu Gly Ala Lys Leu His  
 305 310 315 320  
 Pro Thr Leu Lys Leu Glu Glu Ile Thr Asn Arg Lys Gly Leu Thr Pro  
 325 330 335  
 Leu Ala Leu Ala Ala Ser Ser Gly Lys Ile Gly Val Leu Ala Tyr Ile  
 340 345 350  
 Leu Gln Arg Glu Ile His Glu Pro Glu Cys Arg His Leu Ser Arg Lys  
 355 360 365  
 Phe Thr Glu Trp Ala Tyr Gly Pro Val His Ser Ser Leu Tyr Asp Leu  
 370 375 380  
 Ser Cys Ile Asp Thr Cys Glu Lys Asn Ser Val Leu Glu Val Ile Ala  
 385 390 395 400  
 Tyr Ser Ser Ser Glu Thr Pro Asn Arg His Asp Met Leu Leu Val Glu  
 405 410 415  
 Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp Asp Arg Phe Val Lys Arg  
 420 425 430  
 Ile Phe Tyr Phe Asn Phe Phe Val Tyr Cys Leu Tyr Met Ile Ile Phe  
 435 440 445  
 Thr Ala Ala Ala Tyr Tyr Arg Pro Val Glu Gly Leu Pro Pro Tyr Lys  
 450 455 460  
 Leu Lys Asn Thr Val Gly Asp Tyr Phe Arg Val Thr Gly Glu Ile Leu  
 465 470 475 480  
 Ser Val Ser Gly Gly Val Tyr Phe Phe Phe Arg Gly Ile Gln Tyr Phe  
 485 490 495  
 Leu Gln Arg Arg Pro Ser Leu Lys Ser Leu Phe Val Asp Ser Tyr Ser  
 500 505 510

- 7 -

Glu Ile Leu Phe Phe Val Gln Ser Leu Phe Met Leu Val Ser Val Val  
 515 520 525  
 Leu Tyr Phe Ser Gln Arg Lys Glu Tyr Val Ala Ser Met Val Phe Ser  
 530 535 540  
 Leu Ala Met Gly Trp Thr Asn Met Leu Tyr Tyr Thr Arg Gly Phe Gln  
 545 550 555 560  
 Gln Met Gly Ile Tyr Ala Val Met Ile Glu Lys Met Ile Leu Arg Asp  
 565 570 575  
 Leu Cys Arg Phe Met Phe Val Tyr Leu Val Phe Leu Phe Gly Phe Ser  
 580 585 590  
 Thr Ala Val Val Thr Leu Ile Glu Asp Gly Lys Asn Asn Ser Leu Pro  
 595 600 605  
 Met Glu Ser Thr Pro His Lys Cys Arg Gly Ser Ala Cys Lys Pro Gly  
 610 615 620  
 Asn Ser Tyr Asn Ser Leu Tyr Ser Thr Cys Leu Glu Leu Phe Lys Phe  
 625 630 635 640  
 Thr Ile Gly Met Gly Asp Leu Glu Phe Thr Glu Asn Tyr Asp Phe Lys  
 645 650 655  
 Ala Val Phe Ile Ile Leu Leu Leu Ala Tyr Val Ile Leu Thr Tyr Ile  
 660 665 670  
 Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly Glu Thr Val Asn Lys  
 675 680 685  
 Ile Ala Gln Glu Ser Lys Asn Ile Trp Lys Leu Gln Arg Ala Ile Thr  
 690 695 700  
 Ile Leu Asp Thr Glu Lys Ser Phe Leu Lys Cys Met Arg Lys Ala Phe  
 705 710 715 720  
 Arg Ser Gly Lys Leu Leu Gln Val Gly Phe Thr Pro Asp Gly Lys Asp  
 725 730 735  
 Asp Tyr Arg Trp Cys Phe Arg Val Asp Glu Val Asn Trp Thr Thr Trp  
 740 745 750  
 Asn Thr Asn Val Gly Ile Ile Asn Glu Asp Pro Gly Asn Cys Glu Gly  
 755 760 765  
 Val Lys Arg Thr Leu Ser Phe Ser Leu Arg Ser Gly Arg Val Ser Gly  
 770 775 780  
 Arg Asn Trp Lys Asn Phe Ala Leu Val Pro Leu Leu Arg Asp Ala Ser  
 785 790 795 800  
 Thr Arg Asp Arg His Ala Thr Gln Gln Glu Val Gln Leu Lys His  
 805 810 815  
 Tyr Thr Gly Ser Leu Lys Pro Glu Asp Ala Glu Val Phe Lys Asp Ser  
 820 825 830  
 Met Val Pro Gly Glu Lys  
 835

&lt;210&gt; 3

&lt;211&gt; 2736

&lt;212&gt; DNA

&lt;213&gt; R. rattus

&lt;220&gt;

&lt;400&gt; 3

ggcggttaa	ac	ctgctctgtc	cactgtgtga	gacgaacagg	tggaggggtgg	acgacgcaga	60
gaaagctcgg	agcggggccgc	ggaggttccc	acagcccat	tactgtcagc	gttgagccgc		120
acccctccgg	gccgcacttc	ctctctcagt	ccccgtgcc	ggagagcccc	gctaggctcg		180
gtgatacctag	cctgcagttt	gccgccgcta	cacctgggt	tcagcctgcg	gggtcccagc		240

- 8 -

```

caggcctgcc cctgcggtat gagagaggaa ccttaacatc tccatctcta cagagggtttc 300
agctgtaagg agcatcctcc tctctcagga tgacttcagc ctccagcccc ccagctttca 360
ggctggagac ttccgatgga gatgaagagg gcaatgctga ggtgaacaag gggaagcagg 420
aaccgcccc catggagtca ccattccaga gggaggaccg gaattcctcc cctcagatca 480
aagtgaacct caacttcata aagagacctc ctaaaaacac ttctgctccc agccagcagg 540
agccagatcg gtttgaccgt gaccgactct tcagtgtggt ctcccggggt gtccccgagg 600
aactgactgg actgctagaa tacctgcgct ggaacagcaa gtacctcact gactctgcat 660
acacagaagg ctccactgga aagacgtgcc tgatgaaggc tgtgctgaac cttcaggatg 720
gggtcaatgc ctgcatcatg ccgctgctgc agattgacaa ggattccggc aatcccaagc 780
cctcgtcaa tgcccagtgc accgatgagt tctaccaagg ccacagtgcg ctgcacatcg 840
ccatagagaa gaggagcctg cagtgcgtga agctgctggt agagaatgga gcggatgttc 900
acctccgagc ctgtggccgc ttcttccaaa agcaccaagg aacttgtttc tattttggag 960
agctacctct ttctctggct gcgtgcacca agcagtggga tgtggtgacc tacctcctgg 1020
agaaccacaca ccagccggcc agcctggagg ccaccgactc cctgggcaac acagtcctgc 1080
atgctctggt aatgattgca gataactcgc ctgagaacag tgccctgggtg atccacatgt 1140
acgacgggct tctacaaatg ggggcgcgcc tctgccccac tgtgcagctt gaggaaatct 1200
ccaaccacca aggcctcaca cccctgaaac tagccgcaa ggaaggcaaa atcgagattt 1260
tcaggcacat tctgcagcgg gaattctcag gaccgtacca gcccctttcc cgaaagttaa 1320
ctgagtgggtg ttacggctct gtgcgggtat cgctgtacga cctgtcctct gtggacagct 1380
gggaaaagaa ctcggtgctg gagatcatcg cttttcattg caagagcccg aaccggcacc 1440
gcatggtggt tttagaacca ctgaacaagc ttctgcagga gaaatgggat cggctcgtct 1500
caagattctt cttcaacttc gcctgctact tggctctacat gttcatcttc accgtcgttg 1560
cctaccacca gccttccctg gatcagccag ccattccctc atcaaaaagcg acttttgagg 1620
aatccatgct gctgctgggc cacattctga tctgcttggt gggatattac ctcttactgg 1680
gccagctgtg gtacttttgg cggcggcgcc tgtttatctg gatctcattc atggacagct 1740
actttgaaat cctctttctc cttcaggctc tgctcacagt gctgtcccag gtgctgcgct 1800
tcatggagac tgaatggtac ctacccctgc tagtgttatc cctagtgtctg ggctggctga 1860
acctgcttta ctacacacgg ggctttcagc acacaggcat ctacagtgtc atgatccaga 1920
aggtcactct tcgagacctg ctccgtttcc tgctggtcta cctggtcttc cttttcggct 1980
ttgctgtagc cctagtaagc ttgagcagag agggccgaag tcccaaagcc cctgaagata 2040
acaactccac agtgacggaa cagcccacgg tgggcccagga ggaggagcca gctccatctc 2100
ggagcattct ggatgcctcc cttagagctgt tcaagtccac cattgggtatg ggggagctgg 2160
ctttccagga acagctgcgt tttcgtgggg tggctcctgct gttgctgttg gcctacgtcc 2220
ttctcaccta cgtcctgctg ctcaacatgc tcattgctct catgagcgaa actgtcaacc 2280
acgttgctga caacagctgg agcatctgga agttgcagaa agccatctct gtcttgagga 2340
tggaagaatg ttactggtgg tgccggagga agaaacatcg tgaaggagg ctgctgaaag 2400
tcggcaccag gggggatggt acccctgatg agcgtggtg cttcagggtg gaggaagtaa 2460
attgggctgc ttgggagaag actcttccca ccttatctga ggatccatca gggccaggca 2520
tactggtaa taaaaagaac ccaacctcta aaccggggaa gaacagtgcc tcagaggaag 2580
accatctgcc ctttcaggct ctccagtcct cctgatggcc cagatgcagc agcaggctgg 2640
caggatggag tagggaatct tcccagccac accagaggct actgaatttt ggtggaaata 2700
taaataatttt ttttcataa aaaaaaaaaa aaaaaa 2736

```

&lt;210&gt; 4

&lt;211&gt; 761

&lt;212&gt; PRT

&lt;213&gt; R. rattus

&lt;400&gt; 4

```

Met Thr Ser Ala Ser Ser Pro Pro Ala Phe Arg Leu Glu Thr Ser Asp
 1           5           10          15
Gly Asp Glu Glu Gly Asn Ala Glu Val Asn Lys Gly Lys Gln Glu Pro
          20          25          30
Pro Pro Met Glu Ser Pro Phe Gln Arg Glu Asp Arg Asn Ser Ser Pro
        35          40          45

```

-9-

Gln Ile Lys Val Asn Leu Asn Phe Ile Lys Arg Pro Pro Lys Asn Thr  
 50 55 60  
 Ser Ala Pro Ser Gln Gln Glu Pro Asp Arg Phe Asp Arg Asp Arg Leu  
 65 70 75 80  
 Phe Ser Val Val Ser Arg Gly Val Pro Glu Glu Leu Thr Gly Leu Leu  
 85 90 95  
 Glu Tyr Leu Arg Trp Asn Ser Lys Tyr Leu Thr Asp Ser Ala Tyr Thr  
 100 105 110  
 Glu Gly Ser Thr Gly Lys Thr Cys Leu Met Lys Ala Val Leu Asn Leu  
 115 120 125  
 Gln Asp Gly Val Asn Ala Cys Ile Met Pro Leu Leu Gln Ile Asp Lys  
 130 135 140  
 Asp Ser Gly Asn Pro Lys Pro Leu Val Asn Ala Gln Cys Thr Asp Glu  
 145 150 155 160  
 Phe Tyr Gln Gly His Ser Ala Leu His Ile Ala Ile Glu Lys Arg Ser  
 165 170 175  
 Leu Gln Cys Val Lys Leu Leu Val Glu Asn Gly Ala Asp Val His Leu  
 180 185 190  
 Arg Ala Cys Gly Arg Phe Phe Gln Lys His Gln Gly Thr Cys Phe Tyr  
 195 200 205  
 Phe Gly Glu Leu Pro Leu Ser Leu Ala Ala Cys Thr Lys Gln Trp Asp  
 210 215 220  
 Val Val Thr Tyr Leu Leu Glu Asn Pro His Gln Pro Ala Ser Leu Glu  
 225 230 235 240  
 Ala Thr Asp Ser Leu Gly Asn Thr Val Leu His Ala Leu Val Met Ile  
 245 250 255  
 Ala Asp Asn Ser Pro Glu Asn Ser Ala Leu Val Ile His Met Tyr Asp  
 260 265 270  
 Gly Leu Leu Gln Met Gly Ala Arg Leu Cys Pro Thr Val Gln Leu Glu  
 275 280 285  
 Glu Ile Ser Asn His Gln Gly Leu Thr Pro Leu Lys Leu Ala Ala Lys  
 290 295 300  
 Glu Gly Lys Ile Glu Ile Phe Arg His Ile Leu Gln Arg Glu Phe Ser  
 305 310 315 320  
 Gly Pro Tyr Gln Pro Leu Ser Arg Lys Phe Thr Glu Trp Cys Tyr Gly  
 325 330 335  
 Pro Val Arg Val Ser Leu Tyr Asp Leu Ser Ser Val Asp Ser Trp Glu  
 340 345 350  
 Lys Asn Ser Val Leu Glu Ile Ile Ala Phe His Cys Lys Ser Pro Asn  
 355 360 365  
 Arg His Arg Met Val Val Leu Glu Pro Leu Asn Lys Leu Leu Gln Glu  
 370 375 380  
 Lys Trp Asp Arg Leu Val Ser Arg Phe Phe Phe Asn Phe Ala Cys Tyr  
 385 390 395 400  
 Leu Val Tyr Met Phe Ile Phe Thr Val Val Ala Tyr His Gln Pro Ser  
 405 410 415  
 Leu Asp Gln Pro Ala Ile Pro Ser Ser Lys Ala Thr Phe Gly Glu Ser  
 420 425 430  
 Met Leu Leu Leu Gly His Ile Leu Ile Leu Leu Gly Gly Ile Tyr Leu  
 435 440 445  
 Leu Leu Gly Gln Leu Trp Tyr Phe Trp Arg Arg Arg Leu Phe Ile Trp  
 450 455 460  
 Ile Ser Phe Met Asp Ser Tyr Phe Glu Ile Leu Phe Leu Leu Gln Ala  
 465 470 475 480

-10-

Leu Leu Thr Val Leu Ser Gln Val Leu Arg Phe Met Glu Thr Glu Trp  
                     485                    490                    495  
 Tyr Leu Pro Leu Leu Val Leu Ser Leu Val Leu Gly Trp Leu Asn Leu  
                     500                    505                    510  
 Leu Tyr Tyr Thr Arg Gly Phe Gln His Thr Gly Ile Tyr Ser Val Met  
                     515                    520                    525  
 Ile Gln Lys Val Ile Leu Arg Asp Leu Leu Arg Phe Leu Leu Val Tyr  
                     530                    535                    540  
 Leu Val Phe Leu Phe Gly Phe Ala Val Ala Leu Val Ser Leu Ser Arg  
 545                    550                    555                    560  
 Glu Ala Arg Ser Pro Lys Ala Pro Glu Asp Asn Asn Ser Thr Val Thr  
                     565                    570                    575  
 Glu Gln Pro Thr Val Gly Gln Glu Glu Glu Pro Ala Pro Tyr Arg Ser  
                     580                    585                    590  
 Ile Leu Asp Ala Ser Leu Glu Leu Phe Lys Phe Thr Ile Gly Met Gly  
                     595                    600                    605  
 Glu Leu Ala Phe Gln Glu Gln Leu Arg Phe Arg Gly Val Val Leu Leu  
                     610                    615                    620  
 Leu Leu Leu Ala Tyr Val Leu Leu Thr Tyr Val Leu Leu Leu Asn Met  
 625                    630                    635                    640  
 Leu Ile Ala Leu Met Ser Glu Thr Val Asn His Val Ala Asp Asn Ser  
                     645                    650                    655  
 Trp Ser Ile Trp Lys Leu Gln Lys Ala Ile Ser Val Leu Glu Met Glu  
                     660                    665                    670  
 Asn Gly Tyr Trp Trp Cys Arg Arg Lys Lys His Arg Glu Gly Arg Leu  
                     675                    680                    685  
 Leu Lys Val Gly Thr Arg Gly Asp Gly Thr Pro Asp Glu Arg Trp Cys  
                     690                    695                    700  
 Phe Arg Val Glu Glu Val Asn Trp Ala Ala Trp Glu Lys Thr Leu Pro  
 705                    710                    715                    720  
 Thr Leu Ser Glu Asp Pro Ser Gly Pro Gly Ile Thr Gly Asn Lys Lys  
                     725                    730                    735  
 Asn Pro Thr Ser Lys Pro Gly Lys Asn Ser Ala Ser Glu Glu Asp His  
                     740                    745                    750  
 Leu Pro Leu Gln Val Leu Gln Ser Pro  
                     755                    760

&lt;210&gt; 5

&lt;211&gt; 273

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;400&gt; 5

tgcggtctcc	cnggggtgtc	cccgaggatc	tggctggact	tccnagagta	cctgagcaag	60
accagcaagt	acctcaccga	ctcggaatac	acagagggct	ncnacaggta	agacgtgcct	120
gatgaaggct	gtgctgaacc	ttaaggacgg	ggtcaatgcc	tgcattctgc	cactgctgca	180
gacgachgg	gactctggca	atcctcagcc	cctggtaaata	gcccagtgca	cagatgacta	240
ttaccgaggc	cacagcnctc	tgacatcgc	cat			273

&lt;210&gt; 6

&lt;211&gt; 768

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;400&gt; 6

-11-

tgggtgagct	acccctctct	ttggccgctt	gcaccaagca	gtgggatgtg	gtaagctacc	60
tcctggagaa	cccacaccag	cccgccagcc	tgcagnncac	tgactcccag	ggcaacacag	120
tcctgcatgc	cctagtgatg	atctcggaca	actcagctga	gaacattgca	ctggtgacca	180
gcatgtatga	tgggtctctc	caagctgggg	cccgncctct	gccctaccgt	gcnaagcttga	240
ggacatccgc	aacctgcagg	atctcacgcc	tctgnaannt	ggccgccaag	gagggcaaga	300
tcgrrwttty	maggcacatc	ctnnsmagcg	ggrrktttca	ggactgnagc	caccttttnc	360
ccgaaagttc	accgagtggg	ngctannkcg	gcctgtccgg	gntgtcgctg	tnatgacctg	420
gnnyttctnt	ggacagctgt	naggagaact	cagtgtctga	gatcattgcc	tttcattngc	480
aaragcccg	accgacaccg	aatggtcgtt	ttggagcccc	tgaacaaact	gctgcaggcn	540
gaaatgggat	ctgtctcatc	ccaagttctt	cttaaaacttc	ctgtgtaatc	tgatntacat	600
gttcattctt	amckctgttg	cctaccatca	gcctacccng	aagaagcagg	ccgccccctca	660
cctgaaagcg	gaggttggaa	actccatgct	gctgacgggc	cacatectta	tcctgctagg	720
ggggatctac	ctcctcgtgg	ggcaaaagtg	gaaatttttg	gggggaat		768

&lt;210&gt; 7

&lt;211&gt; 650

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;400&gt; 7

tgtttctctg	ccatcgagtg	gtacctgcc	ctgcttgtgt	ctgcgctggg	gctgggctgg	60
ctgaacctgc	tttactatac	acgtggcttc	cagcacacag	gcatctacag	tgctatgac	120
cagaagccct	ggtgagcctg	agccaggatt	ggcgccccga	agtcctaca	ggccccaatg	180
ccacagagtc	agtgcagccc	atggaggggac	aggaggacga	gggcaacggg	ggccagtaca	240
ggggatctct	ggwagcctcc	ttggagctct	tcaaattcac	catcggcctg	ggcgagctgg	300
ccttccagg	gcagctgcac	ttccgcggca	tnggtgctgc	tgctgctnct	ggcctacgt	360
ctgctcacct	acatcctgct	gctcaacatg	ctcatcgccc	tcatngagcg	agaccgtcaa	420
cankktcgcc	actgacagct	ggagcatctg	gaagctgcag	aaagncatct	nntgtcctgg	480
agatggagaa	tggctattgg	tggtgcanga	agaagcagcg	ggcaggtgtg	atgctgancg	540
ttggcactaa	gccagatggc	agcccs gatg	agcgctgggt	cttcagggtg	gaggaggtga	600
actgggcttc	atngggagca	gacgctgcct	acgctgtgtg	aggaccctgc		650

&lt;210&gt; 8

&lt;211&gt; 317

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;400&gt; 8

gagcttctcc	ctgcggtcaa	gcagagtttc	aggcagacac	tggaagaact	ttgccctggg	60
ccccctttta	agagaggcaa	gtctcgaata	ggcagctctg	tcagcccag	gaagtttata	120
tgcgacagtt	ttcagggtct	ctaaagccag	aggacgtga	ggtcttcaag	agtcctgccg	180
cttccggggg	gaagtggagg	cgtcacgcag	acagcactgt	caacactggg	ccttaggaga	240
ccccgttgcc	acggggggct	ctgagggaac	acagtgcctt	ttcagcagcc	ttgctgggtc	300
ttgctgcccc	gcatggt					317

&lt;210&gt; 9

&lt;211&gt; 65

&lt;212&gt; PRT

&lt;213&gt; H. sapiens

&lt;400&gt; 9

Ser	Phe	Ser	Leu	Arg	Ser	Ser	Arg	Val	Ser	Gly	Arg	His	Trp	Lys	Asn
1				5				10						15	
Phe	Ala	Leu	Val	Pro	Leu	Leu	Arg	Glu	Ala	Ser	Xaa	Arg	Xaa	Arg	Gln
		20						25						30	



-12-

Ser Ala Gln Pro Glu Glu Val Tyr Leu Arg Gln Phe Ser Gly Ser Leu  
           35                          40                          45  
 Lys Pro Glu Asp Ala Glu Val Phe Lys Ser Pro Ala Ala Ser Gly Glu  
       50                          55                          60  
 Lys  
 65

<210> 10  
 <211> 471  
 <212> DNA  
 <213> M. musculus

<400> 10  
 cctgcagaag agcancangc gcctgannga cagcgagttc aaagacccag agacgggaaa 60  
 gacctgtctg ctcaaagcca tgctcaatct gcacaatggg cagaacgaca ccattgctct 120  
 gtccttggaac attgcccggg agacagatag cctgaagcag ttgtcaatg ccagctacac 180  
 agacagctac tacaagggcc agacagcatt acacattgcc attgaaaggc ggaacatggc 240  
 nctggtgacc ctcttggtgg agaatggagc agatgtccag gctgctgctg acgggggactt 300  
 cttcnananaa accaanggga ggcttggtt ctactttggt gagctgcccc tgtccctggc 360  
 tgcgtgcacc aaccagctgg ccattgtgaa attcctgctg cagaactcct gggcagcctg 420  
 cagacatcag tggcncggga ttcngtgggc aacacngtgc tgcacnccct t 471

<210> 11  
 <211> 450  
 <212> DNA  
 <213> M. musculus

<400> 11  
 caagtgtcgg ggatctgcct tgcagggcca agttaattct ttacaacagc ctgtattcca 60  
 catgtctgga gctgttcaag ttcaccatcg gcatgggtga cctggagttc accgagaact 120  
 atgacttcaa ggctgtcttt catcactctg ttactggcct atgtgattct cacctacatc 180  
 ctctgtctca acatgctcat tgctctcatg ggcgagactg tcaacaagat tgcacaagag 240  
 agcaagaaca tctggaagct gcagcgagcc atcaccatcc tggatacaga gaagagtttc 300  
 ctgaagtgca tgaggaaggc cttccgctcc ggcaagctgc tgcaggtggg gttcacgccg 360  
 gacggcaagg atgacttccg gtggtgcttc aggggtggatg aggtgaactg gactacctgg 420  
 aacaccaacg tgggcatcat caacgaggac 450

<210> 12  
 <211> 18  
 <212> DNA  
 <213> R. rattus

<400> 12  
 gaccagcaag tacctcac 18

<210> 13  
 <211> 49  
 <212> DNA  
 <213> R. rattus

<400> 13  
 ctcccatgca gccagttta cttcctccac cctgaagcac cagcgctca 49

<210> 14  
 <211> 71

-13-

&lt;212&gt; PRT

&lt;213&gt; R. rattus

&lt;400&gt; 14

Glu	Leu	Phe	Lys	Phe	Thr	Ile	Gly	Met	Gly	Asp	Leu	Glu	Phe	Thr	Glu
1				5					10					15	
Asn	Tyr	Asp	Phe	Lys	Ala	Val	Phe	Ile	Ile	Leu	Leu	Leu	Ala	Tyr	Val
			20					25					30		
Ile	Leu	Thr	Tyr	Ile	Leu	Leu	Leu	Asn	Met	Leu	Ile	Ala	Leu	Met	Gly
		35					40					45			
Glu	Thr	Val	Asn	Lys	Ile	Ala	Gln	Glu	Ser	Lys	Asn	Ile	Trp	Lys	Leu
	50					55					60				
Gln	Arg	Ala	Ile	Thr	Ile	Leu									
65						70									

&lt;210&gt; 15

&lt;211&gt; 57

&lt;212&gt; PRT

&lt;213&gt; H. sapiens

&lt;400&gt; 15

Glu	Leu	Phe	Lys	Phe	Thr	Ile	Gly	Met	Gly	Glu	Leu	Ala	Phe	Gln	Glu
1				5					10					15	
Gln	Leu	His	Phe	Arg	Gly	Met	Val	Leu	Leu	Leu	Leu	Ala	Tyr	Val	
			20					25				30			
Leu	Leu	Thr	Tyr	Ile	Leu	Leu	Leu	Asn	Met	Leu	Ile	Ala	Leu	Met	Ser
		35					40					45			
Glu	Thr	Val	Asn	Ser	Val	Ala	Thr	Asp							
	50					55									

&lt;210&gt; 16

&lt;211&gt; 75

&lt;212&gt; PRT

&lt;213&gt; Caliphora

&lt;400&gt; 16

Ser	Leu	Phe	Trp	Ala	Ser	Phe	Gly	Leu	Val	Asp	Leu	Val	Ser	Phe	Asp
1				5					10					15	
Leu	Ala	Gly	Ile	Lys	Ser	Phe	Thr	Arg	Phe	Trp	Ala	Leu	Leu	Met	Phe
			20					25				30			
Gly	Ser	Tyr	Ser	Val	Ile	Asn	Ile	Ile	Val	Leu	Leu	Asn	Met	Leu	Ile
		35					40					45			
Ala	Met	Met	Ser	Asn	Ser	Tyr	Gln	Ile	Ile	Ser	Glu	Arg	Ala	Asp	Val
	50					55					60				
Glu	Trp	Lys	Phe	Ala	Arg	Ser	Gln	Leu	Trp	Met					
65						70				75					

&lt;210&gt; 17

&lt;211&gt; 75

&lt;212&gt; PRT

&lt;213&gt; D. melanogaster

&lt;400&gt; 17

Ser	Leu	Phe	Trp	Ala	Ser	Phe	Gly	Leu	Val	Asp	Leu	Val	Ser	Phe	Asp
1				5					10					15	

Leu	Ala	Gly	Ile	Lys	Ser	Phe	Thr	Arg	Phe	Trp	Ala	Leu	Leu	Met	Phe
			20					25					30		
Gly	Ser	Tyr	Ser	Val	Ile	Asn	Ile	Ile	Val	Leu	Leu	Asn	Met	Leu	Ile
		35					40					45			
Ala	Met	Met	Ser	Asn	Ser	Tyr	Gln	Ile	Ile	Ser	Glu	Arg	Ala	Asp	Val
	50					55					60				
Glu	Trp	Lys	Phe	Ala	Arg	Ser	Gln	Leu	Trp	Met					
65					70					75					

```
<210> 18
<211> 75
<212> PRT
<213> B. taurus
```

<400> 18															
Ser	Leu	Phe	Trp	Ser	Ile	Phe	Gly	Leu	Ile	Asn	Leu	Tyr	Val	Thr	Asn
1				5					10					15	
Val	Lys	Ala	Gln	His	Glu	Phe	Thr	Glu	Phe	Val	Gly	Ala	Thr	Met	Phe
			20					25					30		
Gly	Thr	Tyr	Asn	Val	Ile	Ser	Leu	Val	Val	Leu	Leu	Asn	Met	Leu	Ile
		35					40					45			
Ala	Met	Met	Asn	Asn	Ser	Tyr	Gln	Leu	Ile	Ala	Asp	His	Ala	Asp	Ile
	50					55					60				
Glu	Trp	Lys	Phe	Ala	Arg	Thr	Lys	Leu	Trp	Met					
65					70					75					

```
<210> 19
<211> 61
<212> PRT
<213> C. elegans
```

<400> 19															
Arg	Thr	Phe	Ile	Met	Thr	Ile	Gly	Glu	Phe	Ser	Val	Leu	Tyr	Arg	Glu
1				5					10					15	
Met	Ser	Ala	Cys	Asp	Asn	Phe	Trp	Met	Lys	Trp	Ile	Gly	Lys	Leu	Ile
			20					25					30		
Phe	Val	Ile	Phe	Glu	Thr	Phe	Val	Ser	Ile	Leu	Gln	Phe	Asn	Leu	Leu
		35					40					45			
Ile	Ala	Met	Met	Thr	Arg	Thr	Tyr	Glu	Thr	Ile	Phe	Leu			
	50					55					60				

```
<210> 20
<211> 350
<212> DNA
<213> H. sapiens
```

```
<220>
<221> CDS
<222> (41)...(350)
<223> Human VR2 cDNA
```

<400> 20

gagaggtcct ggctggacnn ngcagcctcc tctcctagg atg acc tca ccc tcc  
Met Thr Ser Pro Ser  
1 5

-15-

```

agc tct cca gtt ttc agg ttg gag aca tta gat gga ggc caa gaa gat      103
Ser Ser Pro Val Phe Arg Leu Glu Thr Leu Asp Gly Gly Gln Glu Asp
          10                      15                      20

ggc tct gag gcg gac aga gga aag ctg gat ttt ggg agc ggg ctg cct      151
Gly Ser Glu Ala Asp Arg Gly Lys Leu Asp Phe Gly Ser Gly Leu Pro
          25                      30                      35

ccc atg gag tca cag ttc cag ggc gag gac cgg aaa ttc gcc cct cag      199
Pro Met Glu Ser Gln Phe Gln Gly Glu Asp Arg Lys Phe Ala Pro Gln
          40                      45                      50

ata aga gtc aac ctc aac tac cga aag gga aca ggt gcc agt cag cgg      247
Ile Arg Val Asn Leu Asn Tyr Arg Lys Gly Thr Gly Ala Ser Gln Pro
          55                      60                      65

gat cca aac cga ttt gac cga gat cgg ctc ttc aat gcg gtc tcc cgg      295
Asp Pro Asn Arg Phe Asp Arg Asp Arg Leu Phe Asn Ala Val Ser Arg
          70                      75                      80                      85

ggg gtc ccc gag gat ctg gct gga ctt cca gag tac ctg agc aag acc      343
Gly Val Pro Glu Asp Leu Ala Gly Leu Pro Glu Tyr Leu Ser Lys Thr
          90                      95                      100

agc aag t                                                                350
Ser Lys

```

<210> 21  
 <211> 764  
 <212> DNA  
 <213> H. sapiens

<220>  
 <221> CDS  
 <222> (3)...(764)

```

<400> 21
tc ggt gag cta ccc ctc tct ttg gcc gct tgc acc aag cag tgg gat      47
Gly Glu Leu Pro Leu Ser Leu Ala Ala Cys Thr Lys Gln Trp Asp
  1              5              10              15

gtg gta agc tac ctc ctg gag aac cca cac cag ccc gcc agc ctg cag      95
Val Val Ser Tyr Leu Leu Glu Asn Pro His Gln Pro Ala Ser Leu Gln
          20              25              30

gcc act gac tcc cag ggc aac aca gtc ctg cat gcc cta gtg atg atc      143
Ala Thr Asp Ser Gln Gly Asn Thr Val Leu His Ala Leu Val Met Ile
          35              40              45

tcg gac aac tca gct gag aac att gca ctg gtg acc agc atg tat gat      191
Ser Asp Asn Ser Ala Glu Asn Ile Ala Leu Val Thr Ser Met Tyr Asp
          50              55              60

```

-16-

ggg ctc ctc caa gct ggg gcc cgc ctc tgc cct acc gtg cag ctt gag	239
Gly Leu Leu Gln Ala Gly Ala Arg Leu Cys Pro Thr Val Gln Leu Glu	
65 70 75	
gac atc cgc aac ctg cag gat ctc acg cct ctg aag ctg gcc gcc aag	287
Asp Ile Arg Asn Leu Gln Asp Leu Thr Pro Leu Lys Leu Ala Ala Lys	
80 85 90 95	
gag ggc aag atc grr aty ttc aag gca cat cct tgc aag cgg gaa gtt	335
Glu Gly Lys Ile Xaa Xaa Phe Lys Ala His Pro Cys Lys Arg Glu Val	
100 105 110	
ttc agg act gaa gcc acc ttt tcc ccg aaa gtt cac cga gtg gtg gct	383
Phe Arg Thr Glu Ala Thr Phe Ser Pro Lys Val His Arg Val Val Ala	
115 120 125	
aat ggg gcc tgt ccg ggt tgt cgc tgt aat gac ctg ggc ttt ctg tgg	431
Asn Gly Ala Cys Pro Gly Cys Arg Cys Asn Asp Leu Gly Phe Leu Trp	
130 135 140	
aca gct gtg agg aga act cag tgc tgg rra tca ttg cct ttc att tgc	479
Thr Ala Val Arg Arg Thr Gln Cys Trp Xaa Ser Leu Pro Phe Ile Cys	
145 150 155	
aar agc ccg acc gac acc gaa tgg tgc ttt tgg agc ccc tga aca aac	527
Xaa Ser Pro Thr Asp Thr Glu Trp Ser Phe Trp Ser Pro * Thr Asn	
160 165 170	
tgc tgc agg cga aat ggg atc tgc tca tcc cca agt tct tct taa act	575
Cys Cys Arg Arg Asn Gly Ile Cys Ser Ser Pro Ser Ser Ser * Thr	
175 180 185	
tcc tgt gta atc tga tta cat gtt cat ctt cac cgc tgt tgc cta cca	623
Ser Cys Val Ile * Leu His Val His Leu His Arg Cys Cys Leu Pro	
190 195 200	
tca gcc tac cct gaa gaa gca ggc cgc ccc tca cct gaa agc gga ggt	671
Ser Ala Tyr Pro Glu Glu Ala Gly Arg Pro Ser Pro Glu Ser Gly Gly	
205 210 215 220	
tgg aaa ctc cat gct gct gac ggg cca cat cct tat cct gct agg ggg	719
Trp Lys Leu His Ala Ala Asp Gly Pro His Pro Tyr Pro Ala Arg Gly	
225 230 235	
gat cta cct cct cgt ggg gca aaa gtg gaa att ttg ggg ggg aat	764
Asp Leu Pro Pro Arg Gly Ala Lys Val Glu Ile Leu Gly Gly Asn	
240 245 250	

&lt;210&gt; 22

&lt;211&gt; 884

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

-17-

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (3)...(764)

&lt;223&gt; Human VR2 cDNA

&lt;400&gt; 22

tg ttt cct ggc cat cga gtg gta cct gcc cct gct tgt gtc tgc gct	47
Phe Pro Gly His Arg Val Val Pro Ala Pro Ala Cys Val Cys Ala	
1 5 10 15	
ggg gct ggg ctg gct gaa cct gct tta cta tac acg tgg ctt cca gca	95
Gly Ala Gly Leu Ala Glu Pro Ala Leu Leu Tyr Thr Trp Leu Pro Ala	
20 25 30	
cac agg cat cta cag tgt cat gat cca gaa gcc ctg gtg agc ctg agc	143
His Arg His Leu Gln Cys His Asp Pro Glu Ala Leu Val Ser Leu Ser	
35 40 45	
cag gat tgg cgc ccc gaa gct cct aca ggc ccc aat gcc aca gag tca	191
Gln Asp Trp Arg Pro Glu Ala Pro Thr Gly Pro Asn Ala Thr Glu Ser	
50 55 60	
gtg cag ccc atg gag gga cag gag gac gag ggc aac ggg gcc cag tac	239
Val Gln Pro Met Glu Gly Gln Glu Asp Glu Gly Asn Gly Ala Gln Tyr	
65 70 75	
agg ggt atc ctg gwa gcc tcc ttg gag ctc ttc aaa ttc acc atc ggc	287
Arg Gly Ile Leu Xaa Ala Ser Leu Glu Leu Phe Lys Phe Thr Ile Gly	
80 85 90 95	
atg ggc gag ctg gcc ttc cag gag cag ctg cac ttc cgc ggc atg gtg	335
Met Gly Glu Leu Ala Phe Gln Glu Gln Leu His Phe Arg Gly Met Val	
100 105 110	
ctg ctg ctg ctg ctg gcc tac gtg ctg ctc acc tac atc ctg ctg ctc	383
Leu Leu Leu Leu Leu Ala Tyr Val Leu Leu Thr Tyr Ile Leu Leu Leu	
115 120 125	
aac atg ctc atc gcc ctc wtg agc gag acc gtc aac agt gtc gcc act	431
Asn Met Leu Ile Ala Leu Xaa Ser Glu Thr Val Asn Ser Val Ala Thr	
130 135 140	
gac agc tgg agc atc tgg aag ctg cag aaa gcc atc tct gtc ctg gag	479
Asp Ser Trp Ser Ile Trp Lys Leu Gln Lys Ala Ile Ser Val Leu Glu	
145 150 155	
atg gag aat ggc tat tgg tgg tgc agg aag aag cag cgg gca ggt gtg	527
Met Glu Asn Gly Tyr Trp Trp Cys Arg Lys Lys Gln Arg Ala Gly Val	
160 165 170 175	
atg ctg acc gtt ggc act aag cca gat ggc agc ccs gat gag cgc tgg	575
Met Leu Thr Val Gly Thr Lys Pro Asp Gly Ser Xaa Asp Glu Arg Trp	
180 185 190	

-18-

tgc ttc agg gtg gag gag gtg aac tgg gct tca tgg gag cag acg ctg	623
Cys Phe Arg Val Glu Glu Val Asn Trp Ala Ser Trp Glu Gln Thr Leu	
195 200 205	
cct acg ctg tgt gag gac ccg tca ggg gca ggt gtc cct cga act ctc	671
Pro Thr Leu Cys Glu Asp Pro Ser Gly Ala Gly Val Pro Arg Thr Leu	
210 215 220	
gag aac cct gtc ctg gct tcc cct ccc aag gag gat gag gat ggt gcc	719
Glu Asn Pro Val Leu Ala Ser Pro Pro Lys Glu Asp Glu Asp Gly Ala	
225 230 235	
tct gag gaa aac tat gtg ccc gtc cag ctc ctc cag tcc aac tga	764
Ser Glu Glu Asn Tyr Val Pro Val Gln Leu Leu Gln Ser Asn *	
240 245 250	
tggcccagat gcagcaggag gccagaggac agagcagagg atctttccaa ccacatctgc	824
tggctctggg gtcccagtga attctggtgg caaatatata ttttactaa mmwmaaaac	884

<210> 23  
 <211> 727  
 <212> PRT  
 <213> H. sapiens

<400> 23

Met Thr Ser Pro Ser Ser Ser Pro Val Phe Arg Leu Glu Thr Leu Asp	
1 5 10 15	
Gly Gly Gln Glu Asp Gly Ser Glu Ala Asp Arg Gly Lys Leu Asp Phe	
20 25 30	
Gly Ser Gly Leu Pro Pro Met Glu Ser Gln Phe Gln Gly Glu Asp Arg	
35 40 45	
Lys Phe Ala Pro Gln Ile Arg Val Asn Leu Asn Tyr Arg Lys Gly Thr	
50 55 60	
Gly Ala Ser Gln Pro Asp Pro Asn Arg Phe Asp Arg Asp Arg Leu Phe	
65 70 75 80	
Asn Ala Val Ser Arg Gly Val Pro Glu Asp Leu Ala Gly Leu Pro Glu	
85 90 95	
Tyr Leu Ser Lys Thr Ser Lys Tyr Leu Thr Asp Ser Glu Tyr Thr Glu	
100 105 110	
Gly Ser Thr Gly Lys Thr Cys Leu Met Lys Ala Val Leu Asn Leu Lys	
115 120 125	
Asp Gly Val Asn Ala Cys Ile Leu Pro Leu Leu Gln Ile Asp Arg Asp	
130 135 140	
Ser Gly Asn Pro Gln Pro Leu Val Asn Ala Gln Cys Thr Asp Asp Tyr	
145 150 155 160	
Tyr Arg Gly His Ser Ala Leu His Ile Ala Ile Glu Lys Arg Ser Leu	
165 170 175	
Gln Cys Val Lys Leu Leu Val Glu Asn Gly Ala Asn Val His Ala Arg	
180 185 190	
Ala Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa	
195 200 205	
Gly Glu Leu Pro Leu Ser Leu Ala Ala Cys Thr Lys Gln Trp Asp Val	
210 215 220	
Val Ser Tyr Leu Leu Glu Asn Pro His Gln Pro Ala Ser Leu Gln Ala	
225 230 235 240	

-19-

Thr Asp Ser Gln Gly Asn Thr Val Leu His Ala Leu Val Met Ile Ser  
 245 250 255  
 Asp Asn Ser Ala Glu Asn Ile Ala Leu Val Thr Ser Met Tyr Asp Gly  
 260 265 270  
 Leu Leu Gln Ala Gly Ala Arg Leu Cys Pro Thr Val Gln Leu Glu Asp  
 275 280 285  
 Ile Arg Asn Leu Gln Asp Leu Thr Pro Leu Lys Leu Ala Ala Lys Glu  
 290 295 300  
 Gly Lys Ile Xaa Ile Phe Xaa Arg His Ile Leu Ala Ser Gly Lys Phe  
 305 310 315 320  
 Ser Gly Leu Lys Pro Pro Phe Pro Arg Lys Phe Thr Glu Trp Trp Leu  
 325 330 335  
 Met Gly Pro Val Arg Val Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
 340 345 350  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
 355 360 365  
 Pro Asp Arg His Arg Met Val Val Leu Glu Pro Leu Asn Lys Leu Leu  
 370 375 380  
 Gln Ala Lys Trp Asp Leu Leu Ile Pro Lys Phe Phe Leu Asn Phe Leu  
 385 390 395 400  
 Cys Asn Leu Xaa Tyr Met Phe Ile Phe Thr Ala Val Ala Tyr His Gln  
 405 410 415  
 Pro Thr Leu Lys Lys Gln Ala Ala Pro His Leu Lys Ala Glu Val Gly  
 420 425 430  
 Asn Ser Met Leu Leu Thr Gly His Ile Leu Ile Leu Leu Gly Gly Ile  
 435 440 445  
 Tyr Leu Leu Val Gly Gln Lys Trp Lys Phe Trp Xaa Xaa Xaa Xaa Xaa  
 450 455 460  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Phe Pro Gly His Arg Val  
 465 470 475 480  
 Val Pro Ala Pro Ala Cys Val Cys Ala Gly Ala Gly Leu Ala Glu Pro  
 485 490 495  
 Ala Leu Leu Tyr Thr Trp Leu Pro Ala His Arg His Leu Gln Cys His  
 500 505 510  
 Asp Pro Glu Ala Leu Val Ser Leu Ser Gln Asp Trp Arg Pro Glu Ala  
 515 520 525  
 Pro Thr Gly Pro Asn Ala Thr Glu Ser Val Gln Pro Met Glu Gly Gln  
 530 535 540  
 Glu Asp Glu Gly Asn Gly Ala Gln Tyr Arg Gly Ile Leu Xaa Ala Ser  
 545 550 555 560  
 Leu Glu Leu Phe Lys Phe Thr Ile Gly Met Gly Glu Leu Ala Phe Gln  
 565 570 575  
 Glu Gln Leu His Phe Arg Gly Met Val Leu Leu Leu Leu Ala Tyr  
 580 585 590  
 Val Leu Leu Thr Tyr Ile Leu Leu Leu Asn Met Leu Ile Ala Leu Xaa  
 595 600 605  
 Ser Glu Thr Val Asn Ser Val Ala Thr Asp Ser Trp Ser Ile Trp Lys  
 610 615 620  
 Leu Gln Lys Ala Ile Ser Val Leu Glu Met Glu Asn Gly Tyr Trp Trp  
 625 630 635 640  
 Cys Arg Lys Lys Gln Arg Ala Gly Val Met Leu Thr Val Gly Thr Lys  
 645 650 655  
 Pro Asp Gly Ser Pro Asp Glu Arg Trp Cys Phe Arg Val Glu Glu Val  
 660 665 670



-20-

Asn Trp Ala Ser Trp Glu Gln Thr Leu Pro Thr Leu Cys Glu Asp Pro  
 675 680 685  
 Ser Gly Ala Gly Val Pro Arg Thr Leu Glu Asn Pro Val Leu Ala Ser  
 690 695 700  
 Pro Pro Lys Glu Asp Glu Asp Gly Ala Ser Glu Glu Asn Tyr Val Pro  
 705 710 715 720  
 Val Gln Leu Leu Gln Ser Asn  
 725

<210> 24  
 <211> 2845  
 <212> DNA  
 <213> chicken

<220>  
 <221> CDS  
 <222> (121)...(2649)  
 <223> chicken VR1 cDNA

<400> 24  
 ttccctataga acagagggag tttgctgaga ctggactatt ctgcagaaga caggatattt 60  
 ttgcaatatt tggatgatcg acggagggac acagaaaact tgaaaggctg cttgttcac 120  
 atg tct tcc att ctt gag aag atg aag aaa ttt ggc agt tct gac ata 168  
 Met Ser Ser Ile Leu Glu Lys Met Lys Lys Phe Gly Ser Ser Asp Ile  
 1 5 10 15  
 gaa gaa tct gaa gtg aca gat gaa cac acg gat ggg gaa gac tca gca 216  
 Glu Glu Ser Glu Val Thr Asp Glu His Thr Asp Gly Glu Asp Ser Ala  
 20 25 30  
 ctg gaa aca gct gac aac ctc cag ggt aca ttc agc aac aag gtg cag 264  
 Leu Glu Thr Ala Asp Asn Leu Gln Gly Thr Phe Ser Asn Lys Val Gln  
 35 40 45  
 cca tcc aaa agc aac atc ttt gca aga cgt gga cgg ttt gtg atg ggg 312  
 Pro Ser Lys Ser Asn Ile Phe Ala Arg Arg Gly Arg Phe Val Met Gly  
 50 55 60  
 gat tgt gac aag gac atg gct cca atg gac tcc ttt tac cag atg gat 360  
 Asp Cys Asp Lys Asp Met Ala Pro Met Asp Ser Phe Tyr Gln Met Asp  
 65 70 75 80  
 cac ctg atg gca cct tct gtc atc aaa ttt cat gcc aat atg gag agg 408  
 His Leu Met Ala Pro Ser Val Ile Lys Phe His Ala Asn Met Glu Arg  
 85 90 95  
 ggg aaa ctt cac aag ctc ctg tca aca gac tcc atc aca ggc tgc tct 456  
 Gly Lys Leu His Lys Leu Leu Ser Thr Asp Ser Ile Thr Gly Cys Ser  
 100 105 110  
 gaa aaa gct ttc aaa ttt tat gac cgc aga agg atc ttt gat gct gta 504  
 Glu Lys Ala Phe Lys Phe Tyr Asp Arg Arg Arg Ile Phe Asp Ala Val  
 115 120 125

-21-

gcc cga ggc agc aca aag gac ctg gat gat ctg ctg ctc tat cta aat	552
Ala Arg Gly Ser Thr Lys Asp Leu Asp Asp Leu Leu Leu Tyr Leu Asn	
130 135 140	
agg acc ttg aag cat ctc aca gat gat gaa ttc aaa gaa cca gaa act	600
Arg Thr Leu Lys His Leu Thr Asp Asp Glu Phe Lys Glu Pro Glu Thr	
145 150 155 160	
ggg aaa acc tgc tta ctg aaa gcc atg ctg aat cta cat gat ggg aaa	648
Gly Lys Thr Cys Leu Leu Lys Ala Met Leu Asn Leu His Asp Gly Lys	
165 170 175	
aat gat acc att ccc ttg ctg ctg gat att gca aag aaa act gga act	696
Asn Asp Thr Ile Pro Leu Leu Leu Asp Ile Ala Lys Lys Thr Gly Thr	
180 185 190	
ctg aaa gag ttt gta aat gca gaa tat act gac aac tat tac aaa ggc	744
Leu Lys Glu Phe Val Asn Ala Glu Tyr Thr Asp Asn Tyr Tyr Lys Gly	
195 200 205	
cag act gca ctc cac att gcc att gag aga agg aac atg tac ctg gta	792
Gln Thr Ala Leu His Ile Ala Ile Glu Arg Arg Asn Met Tyr Leu Val	
210 215 220	
aaa ctc ttg gtc cag aat gga gca gat gtt cat gca aga gca tgt ggg	840
Lys Leu Leu Val Gln Asn Gly Ala Asp Val His Ala Arg Ala Cys Gly	
225 230 235 240	
gag ttc ttc agg aaa atc aaa ggg aaa cct ggt ttt tat ttt gga gag	888
Glu Phe Phe Arg Lys Ile Lys Gly Lys Pro Gly Phe Tyr Phe Gly Glu	
245 250 255	
ctg ccc ctg tcc ctg gct gcc tgc acc aat cag ctc tgc att gtg aaa	936
Leu Pro Leu Ser Leu Ala Ala Cys Thr Asn Gln Leu Cys Ile Val Lys	
260 265 270	
ttt ctc ctt gag aac ccc tac cag gct gct gac att gct gct gag gac	984
Phe Leu Leu Glu Asn Pro Tyr Gln Ala Ala Asp Ile Ala Ala Glu Asp	
275 280 285	
tcc atg ggc aat atg gtt ctg cat act ctg gtg gag att gca gat aat	1032
Ser Met Gly Asn Met Val Leu His Thr Leu Val Glu Ile Ala Asp Asn	
290 295 300	
act aag gat aat acc aag ttc gtt acg aag atg tac aat aac ata ttg	1080
Thr Lys Asp Asn Thr Lys Phe Val Thr Lys Met Tyr Asn Asn Ile Leu	
305 310 315 320	
atc ctt ggt gcc aaa ata aat cct atc ctg aag ttg gaa gaa ctc acc	1128
Ile Leu Gly Ala Lys Ile Asn Pro Ile Leu Lys Leu Glu Glu Leu Thr	
325 330 335	
aac aaa aaa ggg ctg act cca tta acg ttg gca gcc aaa aca ggg aag	1176
Asn Lys Lys Gly Leu Thr Pro Leu Thr Leu Ala Ala Lys Thr Gly Lys	
340 345 350	

-22-

ata ggg att ttc gct tac atc ctc aga cga gag atc aaa gat cct gaa Ile Gly Ile Phe Ala Tyr Ile Leu Arg Arg Glu Ile Lys Asp Pro Glu 355 360 365	1224
tgc aga cac ttg tct agg aag ttc act gaa tgg gct tat gga cct gtc Cys Arg His Leu Ser Arg Lys Phe Thr Glu Trp Ala Tyr Gly Pro Val 370 375 380	1272
cat tca tct ctt tat gac ctg tcc tgc ata gac aca tgt gag aaa aat His Ser Ser Leu Tyr Asp Leu Ser Cys Ile Asp Thr Cys Glu Lys Asn 385 390 395 400	1320
tca gtg ctt gaa att att gcc tac agt agt gaa aca cca aat cgt cat Ser Val Leu Glu Ile Ile Ala Tyr Ser Ser Glu Thr Pro Asn Arg His 405 410 415	1368
gag atg ctg ctg gta gag ccc ctt aac agg cta ctg caa gac aag tgg Glu Met Leu Leu Val Glu Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp 420 425 430	1416
gac cga ttt gtc aag cac tta ttt tac ttc aac ttc ttt gta tat gca Asp Arg Phe Val Lys His Leu Phe Tyr Phe Asn Phe Phe Val Tyr Ala 435 440 445	1464
att cat atc agc atc ctc acc aca gct gcc tac tac aga cct gtg cag Ile His Ile Ser Ile Leu Thr Thr Ala Ala Tyr Tyr Arg Pro Val Gln 450 455 460	1512
aag ggg gac aag cct ccc ttc gct ttt ggt cac agc act ggg gaa tat Lys Gly Asp Lys Pro Pro Phe Ala Phe Gly His Ser Thr Gly Glu Tyr 465 470 475 480	1560
ttt cga gtg act gga gag ata ctg agt gta ttg gga gga ctg tat ttt Phe Arg Val Thr Gly Glu Ile Leu Ser Val Leu Gly Gly Leu Tyr Phe 485 490 495	1608
ttt ttc aga ggg ata cag tat ttt gtg cag agg cgc cca tca ttg aag Phe Phe Arg Gly Ile Gln Tyr Phe Val Gln Arg Arg Pro Ser Leu Lys 500 505 510	1656
acg ctg ata gtt gac agt tac agt gaa gtt ctt ttc ttc gtt cac tct Thr Leu Ile Val Asp Ser Tyr Ser Glu Val Leu Phe Phe Val His Ser 515 520 525	1704
ttg ctc ctc ctg agc tct gtg gtg ctg tac ttc tgt ggc cag gaa ctg Leu Leu Leu Leu Ser Ser Val Val Leu Tyr Phe Cys Gly Gln Glu Leu 530 535 540	1752
tat gtg gct tcc atg gtc ttc tcc ttg gct ctg ggc tgg gct aac atg Tyr Val Ala Ser Met Val Phe Ser Leu Ala Leu Gly Trp Ala Asn Met 545 550 555 560	1800

-23-

cta tac tac acc cgt ggc ttc cag cag atg ggc att tac tct gtc atg	1848
Leu Tyr Tyr Thr Arg Gly Phe Gln Gln Met Gly Ile Tyr Ser Val Met	
565 570 575	
att gca aag atg atc cta aga gac tta tgt cgc ttc atg ttt gtc tat	1896
Ile Ala Lys Met Ile Leu Arg Asp Leu Cys Arg Phe Met Phe Val Tyr	
580 585 590	
cta gta ttc ctc ttg gga ttt tcc aca gct gtg gtg act tta att gaa	1944
Leu Val Phe Leu Leu Gly Phe Ser Thr Ala Val Val Thr Leu Ile Glu	
595 600 605	
gat gac aat gag ggg cag gac aca aat agc tct gaa tat gcc cga tgc	1992
Asp Asp Asn Glu Gly Gln Asp Thr Asn Ser Ser Glu Tyr Ala Arg Cys	
610 615 620	
agc cat acg aaa cga ggc cgc aca tcc tat aac agt ctg tat tat acc	2040
Ser His Thr Lys Arg Gly Arg Thr Ser Tyr Asn Ser Leu Tyr Tyr Thr	
625 630 635 640	
tgc ttg gaa ctt ttc aag ttc act att ggg atg gga gac ctg gag ttt	2088
Cys Leu Glu Leu Phe Lys Phe Thr Ile Gly Met Gly Asp Leu Glu Phe	
645 650 655	
aca gag aac tac agg ttc aag tct gtg ttt gtc atc ctt ttg gtt ctc	2136
Thr Glu Asn Tyr Arg Phe Lys Ser Val Phe Val Ile Leu Leu Val Leu	
660 665 670	
tat gtc atc ctt acg tac atc ctc ctg ctc aat atg ctt att gca ctg	2184
Tyr Val Ile Leu Thr Tyr Ile Leu Leu Leu Asn Met Leu Ile Ala Leu	
675 680 685	
atg gga gaa act gtg agc aaa att gca cag gag agc aag agc atc tgg	2232
Met Gly Glu Thr Val Ser Lys Ile Ala Gln Glu Ser Lys Ser Ile Trp	
690 695 700	
aaa ctc cag agg ccc atc acg atc ttg gat att gaa aac agc tac ttg	2280
Lys Leu Gln Arg Pro Ile Thr Ile Leu Asp Ile Glu Asn Ser Tyr Leu	
705 710 715 720	
aac tgt ttg agg cgc tca ttc cga tct gga aaa aga gtc ttg gtg gga	2328
Asn Cys Leu Arg Arg Ser Phe Arg Ser Gly Lys Arg Val Leu Val Gly	
725 730 735	
atc aca cct gat ggc caa gat gat tac aga tgg tgc ttt aga gtt gat	2376
Ile Thr Pro Asp Gly Gln Asp Asp Tyr Arg Trp Cys Phe Arg Val Asp	
740 745 750	
gaa gtg aac tgg tcc acg tgg aat aca aat ttg ggc ata atc aac gaa	2424
Glu Val Asn Trp Ser Thr Trp Asn Thr Asn Leu Gly Ile Ile Asn Glu	
755 760 765	
gat cct ggg tgc tct ggt gac ctc aaa cga aat ccc agt tac tgt att	2472
Asp Pro Gly Cys Ser Gly Asp Leu Lys Arg Asn Pro Ser Tyr Cys Ile	
770 775 780	

-24-

aag cct ggt aga gtt tca ggg aaa aat tgg aaa act ttg gtt cca ctt 2520  
 Lys Pro Gly Arg Val Ser Gly Lys Asn Trp Lys Thr Leu Val Pro Leu  
 785 790 795 800

tta aga gat gga agc agg aga gaa gaa aca cca aaa cta cca gaa gaa 2568  
 Leu Arg Asp Gly Ser Arg Arg Glu Glu Thr Pro Lys Leu Pro Glu Glu  
 805 810 815

atc aaa tta aaa ccc att ttg gaa cct tat tat gag cca gag gat tgt 2616  
 Ile Lys Leu Lys Pro Ile Leu Glu Pro Tyr Tyr Glu Pro Glu Asp Cys  
 820 825 830

gag aca ttg aag gaa tcg ctt cca aag tca gtc tgatcttttg tttttaagaa 2669  
 Glu Thr Leu Lys Glu Ser Leu Pro Lys Ser Val  
 835 840

ggttaattct agttgtttgt gttggttctt acaaggagga caattaaaac gcttccttca 2729  
 taagagcggg gatttatgga aaaaggccaa agaagctagg aaatgactgt gtgcaaggat 2789  
 tcattaagta tcttgaataa actacttgtt gtttaaaaaa aaaaaaaaaa aaaaaa 2845

&lt;210&gt; 25

&lt;211&gt; 843

&lt;212&gt; PRT

&lt;213&gt; chicken

&lt;400&gt; 25

Met Ser Ser Ile Leu Glu Lys Met Lys Lys Phe Gly Ser Ser Asp Ile  
 1 5 10 15  
 Glu Glu Ser Glu Val Thr Asp Glu His Thr Asp Gly Glu Asp Ser Ala  
 20 25 30  
 Leu Glu Thr Ala Asp Asn Leu Gln Gly Thr Phe Ser Asn Lys Val Gln  
 35 40 45  
 Pro Ser Lys Ser Asn Ile Phe Ala Arg Arg Gly Arg Phe Val Met Gly  
 50 55 60  
 Asp Cys Asp Lys Asp Met Ala Pro Met Asp Ser Phe Tyr Gln Met Asp  
 65 70 75 80  
 His Leu Met Ala Pro Ser Val Ile Lys Phe His Ala Asn Met Glu Arg  
 85 90 95  
 Gly Lys Leu His Lys Leu Leu Ser Thr Asp Ser Ile Thr Gly Cys Ser  
 100 105 110  
 Glu Lys Ala Phe Lys Phe Tyr Asp Arg Arg Arg Ile Phe Asp Ala Val  
 115 120 125  
 Ala Arg Gly Ser Thr Lys Asp Leu Asp Asp Leu Leu Leu Tyr Leu Asn  
 130 135 140  
 Arg Thr Leu Lys His Leu Thr Asp Asp Glu Phe Lys Glu Pro Glu Thr  
 145 150 155 160  
 Gly Lys Thr Cys Leu Leu Lys Ala Met Leu Asn Leu His Asp Gly Lys  
 165 170 175  
 Asn Asp Thr Ile Pro Leu Leu Leu Asp Ile Ala Lys Lys Thr Gly Thr  
 180 185 190  
 Leu Lys Glu Phe Val Asn Ala Glu Tyr Thr Asp Asn Tyr Tyr Lys Gly  
 195 200 205  
 Gln Thr Ala Leu His Ile Ala Ile Glu Arg Arg Asn Met Tyr Leu Val  
 210 215 220

-25-

Lys Leu Leu Val Gln Asn Gly Ala Asp Val His Ala Arg Ala Cys Gly  
 225 230 235 240  
 Glu Phe Phe Arg Lys Ile Lys Gly Lys Pro Gly Phe Tyr Phe Gly Glu  
 245 250 255  
 Leu Pro Leu Ser Leu Ala Ala Cys Thr Asn Gln Leu Cys Ile Val Lys  
 260 265 270  
 Phe Leu Leu Glu Asn Pro Tyr Gln Ala Ala Asp Ile Ala Ala Glu Asp  
 275 280 285  
 Ser Met Gly Asn Met Val Leu His Thr Leu Val Glu Ile Ala Asp Asn  
 290 295 300  
 Thr Lys Asp Asn Thr Lys Phe Val Thr Lys Met Tyr Asn Asn Ile Leu  
 305 310 315 320  
 Ile Leu Gly Ala Lys Ile Asn Pro Ile Leu Lys Leu Glu Glu Leu Thr  
 325 330 335  
 Asn Lys Lys Gly Leu Thr Pro Leu Thr Leu Ala Ala Lys Thr Gly Lys  
 340 345 350  
 Ile Gly Ile Phe Ala Tyr Ile Leu Arg Arg Glu Ile Lys Asp Pro Glu  
 355 360 365  
 Cys Arg His Leu Ser Arg Lys Phe Thr Glu Trp Ala Tyr Gly Pro Val  
 370 375 380  
 His Ser Ser Leu Tyr Asp Leu Ser Cys Ile Asp Thr Cys Glu Lys Asn  
 385 390 395 400  
 Ser Val Leu Glu Ile Ile Ala Tyr Ser Ser Glu Thr Pro Asn Arg His  
 405 410 415  
 Glu Met Leu Leu Val Glu Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp  
 420 425 430  
 Asp Arg Phe Val Lys His Leu Phe Tyr Phe Asn Phe Phe Val Tyr Ala  
 435 440 445  
 Ile His Ile Ser Ile Leu Thr Thr Ala Ala Tyr Tyr Arg Pro Val Gln  
 450 455 460  
 Lys Gly Asp Lys Pro Pro Phe Ala Phe Gly His Ser Thr Gly Glu Tyr  
 465 470 475 480  
 Phe Arg Val Thr Gly Glu Ile Leu Ser Val Leu Gly Gly Leu Tyr Phe  
 485 490 495  
 Phe Phe Arg Gly Ile Gln Tyr Phe Val Gln Arg Arg Pro Ser Leu Lys  
 500 505 510  
 Thr Leu Ile Val Asp Ser Tyr Ser Glu Val Leu Phe Phe Val His Ser  
 515 520 525  
 Leu Leu Leu Leu Ser Ser Val Val Leu Tyr Phe Cys Gly Gln Glu Leu  
 530 535 540  
 Tyr Val Ala Ser Met Val Phe Ser Leu Ala Leu Gly Trp Ala Asn Met  
 545 550 555 560  
 Leu Tyr Tyr Thr Arg Gly Phe Gln Gln Met Gly Ile Tyr Ser Val Met  
 565 570 575  
 Ile Ala Lys Met Ile Leu Arg Asp Leu Cys Arg Phe Met Phe Val Tyr  
 580 585 590  
 Leu Val Phe Leu Leu Gly Phe Ser Thr Ala Val Val Thr Leu Ile Glu  
 595 600 605  
 Asp Asp Asn Glu Gly Gln Asp Thr Asn Ser Ser Glu Tyr Ala Arg Cys  
 610 615 620  
 Ser His Thr Lys Arg Gly Arg Thr Ser Tyr Asn Ser Leu Tyr Tyr Thr  
 625 630 635 640  
 Cys Leu Glu Leu Phe Lys Phe Thr Ile Gly Met Gly Asp Leu Glu Phe  
 645 650 655

-26-

Thr Glu Asn Tyr Arg Phe Lys Ser Val Phe Val Ile Leu Leu Val Leu  
                   660                  665                  670  
 Tyr Val Ile Leu Thr Tyr Ile Leu Leu Leu Asn Met Leu Ile Ala Leu  
                   675                  680                  685  
 Met Gly Glu Thr Val Ser Lys Ile Ala Gln Glu Ser Lys Ser Ile Trp  
                   690                  695                  700  
 Lys Leu Gln Arg Pro Ile Thr Ile Leu Asp Ile Glu Asn Ser Tyr Leu  
 705                  710                  715                  720  
 Asn Cys Leu Arg Arg Ser Phe Arg Ser Gly Lys Arg Val Leu Val Gly  
                   725                  730                  735  
 Ile Thr Pro Asp Gly Gln Asp Asp Tyr Arg Trp Cys Phe Arg Val Asp  
                   740                  745                  750  
 Glu Val Asn Trp Ser Thr Trp Asn Thr Asn Leu Gly Ile Ile Asn Glu  
                   755                  760                  765  
 Asp Pro Gly Cys Ser Gly Asp Leu Lys Arg Asn Pro Ser Tyr Cys Ile  
                   770                  775                  780  
 Lys Pro Gly Arg Val Ser Gly Lys Asn Trp Lys Thr Leu Val Pro Leu  
 785                  790                  795                  800  
 Leu Arg Asp Gly Ser Arg Arg Glu Glu Thr Pro Lys Leu Pro Glu Glu  
                   805                  810                  815  
 Ile Lys Leu Lys Pro Ile Leu Glu Pro Tyr Tyr Glu Pro Glu Asp Cys  
                   820                  825                  830  
 Glu Thr Leu Lys Glu Ser Leu Pro Lys Ser Val  
                   835                  840

<210> 26  
 <211> 135  
 <212> DNA  
 <213> H. sapiens

<400> 26  
 ttcaagttta cgatcgggat gggcgacctg gagttcactg agaactatga cttcaaggct 60  
 gtcttcatca tctgtctgct ggcctatgta attctcacct acatcctcct gctcaacatg 120  
 tttatcgctc tcatg 135

<210> 27  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> consensus

<400> 27  
 tttcaaagtt tcacgatcat atcgggatca tg 32

<210> 28  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> consensus

<400> 28

-27-

catgatcaga gatcgcgata tgatcagaca tagtt

35

&lt;210&gt; 29

&lt;211&gt; 135

&lt;212&gt; DNA

&lt;213&gt; chicken

&lt;400&gt; 29

ttcaagttca cgattgggat gggtgacctg gattttcatg aacatgccag attcagatac	60
tttgtcatgc ttctgtctgt gctttttgtg atcctcacct acatcctttt gctcaacatg	120
cttatagccc ttata	135

&lt;210&gt; 30

&lt;211&gt; 135

&lt;212&gt; DNA

&lt;213&gt; chicken

&lt;400&gt; 30

ttcaagttca ctattgggat gggagacctg gagtttacag agaactacag gttcaagtct	60
gtgtttgtca tccttttggt tctctatgtc atccttacgt acatcctcct gctcaatatg	120
cttatagccc taatg	135

&lt;210&gt; 31

&lt;211&gt; 28

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;400&gt; 31

tatctttcaa tctttctttc gtgatcta	28
--------------------------------	----

&lt;210&gt; 32

&lt;211&gt; 17

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;400&gt; 32

aaaaggggga ccagggc	17
--------------------	----

&lt;210&gt; 33

&lt;211&gt; 2544

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (14) ... (2530)

&lt;223&gt; Human VR1

&lt;400&gt; 33

ggatccagca agg atg aag aaa tgg agc agc aca gac ttg ggg gca gct	49
Met Lys Lys Trp Ser Ser Thr Asp Leu Gly Ala Ala	
1 5 10	



- 28 -

gcg gac cca ctc caa aag gac acc tgc cca gac ccc ctg gat gga gac Ala Asp Pro Leu Gln Lys Asp Thr Cys Pro Asp Pro Leu Asp Gly Asp 15 20 25	97
cct aac tcc agg cca cct cca gcc aag ccc cag ctc tcc acg gcc aag Pro Asn Ser Arg Pro Pro Pro Ala Lys Pro Gln Leu Ser Thr Ala Lys 30 35 40	145
agc cgc acc cgg ctc ttt ggg aag ggt gac tgc gag gag gct ttc ccg Ser Arg Thr Arg Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Phe Pro 45 50 55 60	193
gtg gat tgc cct cac gag gaa ggt gag ctg gac tcc tgc ccg acc atc Val Asp Cys Pro His Glu Glu Gly Glu Leu Asp Ser Cys Pro Thr Ile 65 70 75	241
aca gtc agc cct gtt atc acc atc cag agg cca gga gac ggc ccc acc Thr Val Ser Pro Val Ile Thr Ile Gln Arg Pro Gly Asp Gly Pro Thr 80 85 90	289
ggt gcc agg ctg ctg tcc cag gac tct gtc gcc gcc agc acc gag aag Gly Ala Arg Leu Leu Ser Gln Asp Ser Val Ala Ala Ser Thr Glu Lys 95 100 105	337
acc ctc agg ctc tat gat cgc agg agt atc ttt gaa gcc gtt gct cag Thr Leu Arg Leu Tyr Asp Arg Arg Ser Ile Phe Glu Ala Val Ala Gln 110 115 120	385
aat aac tgc cag gat ctg gag agc ctg ctg ctc ttc ctg cag aag agc Asn Asn Cys Gln Asp Leu Glu Ser Leu Leu Leu Phe Leu Gln Lys Ser 125 130 135 140	433
aag aag cac ctc aca gac aac gag ttc aaa gac cct gag aca ggg aag Lys Lys His Leu Thr Asp Asn Glu Phe Lys Asp Pro Glu Thr Gly Lys 145 150 155	481
acc tgt ctg ctg aaa gcc atg ctc aac ctg cac gac gga cag aac acc Thr Cys Leu Leu Lys Ala Met Leu Leu His Asp Gly Gln Asn Thr 160 165 170	529
acc atc ccc ctg ctc ctg gag atc gcg cgg caa acg gac agc ctg aag Thr Ile Pro Leu Leu Leu Glu Ile Ala Arg Gln Thr Asp Ser Leu Lys 175 180 185	577
gag ctt gtc aac gcc agc tac acg gac agc tac tac aag ggc cag aca Glu Leu Val Asn Ala Ser Tyr Thr Asp Ser Tyr Tyr Lys Gly Gln Thr 190 195 200	625
gca ctg cac atc gcc atc gag aga cgc aac atg gcc ctg gtg acc ctc Ala Leu His Ile Ala Ile Glu Arg Arg Asn Met Ala Leu Val Thr Leu 205 210 215 220	673
ctg gtg gag aac gga gca gac gtc cag gct gcg gcc cat ggg gac ttc Leu Val Glu Asn Gly Ala Asp Val Gln Ala Ala Ala His Gly Asp Phe 225 230 235	721

-29-

ttt aag aaa acc aaa ggg cgg cct gga ttc tac ttc ggt gaa ctg ccc Phe Lys Lys Thr Lys Gly Arg Pro Gly Phe Tyr Phe Gly Glu Leu Pro 240 245 250	769
ctg tcc ctg gcc gcg tgc acc aac cag ctg ggc atc gtg aag ttc ctg Leu Ser Leu Ala Ala Cys Thr Asn Gln Leu Gly Ile Val Lys Phe Leu 255 260 265	817
ctg cag aac tcc tgg cag acg gcc gac atc agc gcc agg gac tcg gtg Leu Gln Asn Ser Trp Gln Thr Ala Asp Ile Ser Ala Arg Asp Ser Val 270 275 280	865
ggc aac acg gtg ctg cac gcc ctg gtg gag gtg gcc gac aac acg gcc Gly Asn Thr Val Leu His Ala Leu Val Glu Val Ala Asp Asn Thr Ala 285 290 295 300	913
gac aac acg aag ttt gtg acg agc atg tac aat gag att ctg atc ctg Asp Asn Thr Lys Phe Val Thr Ser Met Tyr Asn Glu Ile Leu Ile Leu 305 310 315	961
ggg gcc aaa ctg cac ccg acg ctg aag ctg gag gag ctc acc aac aag Gly Ala Lys Leu His Pro Thr Leu Lys Leu Glu Glu Leu Thr Asn Lys 320 325 330	1009
aag gga atg acg ccg ctg gct ctg gca gct ggg acc ggg aag atc ggg Lys Gly Met Thr Pro Leu Ala Leu Ala Ala Gly Thr Gly Lys Ile Gly 335 340 345	1057
gtc ttg gcc tat att ctc cag cgg gag atc cag gag ccc gag tgc agg Val Leu Ala Tyr Ile Leu Gln Arg Glu Ile Gln Glu Pro Glu Cys Arg 350 355 360	1105
cac ctg tcc agg aag ttc acc gag tgg gcc tac ggg ccc gtg cac tcc His Leu Ser Arg Lys Phe Thr Glu Trp Ala Tyr Gly Pro Val His Ser 365 370 375 380	1153
tcg ctg tac gac ctg tcc tgc atc gac acc tgc gag aag aac tcg gtg Ser Leu Tyr Asp Leu Ser Cys Ile Asp Thr Cys Glu Lys Asn Ser Val 385 390 395	1201
ctg gag gtg atc gcc tac agc agc agc gag acc cct aat cgc cac gac Leu Glu Val Ile Ala Tyr Ser Ser Ser Glu Thr Pro Asn Arg His Asp 400 405 410	1249
atg ctc ttg gtg gag ccg ctg aac cga ctc ctg cag gac aag tgg gac Met Leu Leu Val Glu Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp Asp 415 420 425	1297
aga ttc gtc aag cgc atc ttc tac ttc aac ttc ctg gtc tac tgc ctg Arg Phe Val Lys Arg Ile Phe Tyr Phe Asn Phe Leu Val Tyr Cys Leu 430 435 440	1345

-30-

tac atg atc atc ttc acc atg gct gcc tac tac agg ccc gtg gat ggc	1393
Tyr Met Ile Ile Phe Thr Met Ala Ala Tyr Tyr Arg Pro Val Asp Gly	
445 450 455 460	
ttg cct ccc ttt aag atg gaa aaa act gga gac tat ttc cga gtt act	1441
Leu Pro Pro Phe Lys Met Glu Lys Thr Gly Asp Tyr Phe Arg Val Thr	
465 470 475	
gga gag atc ctg tct gtg tta gga gga gtc tac ttc ttt ttc cga ggg	1489
Gly Glu Ile Leu Ser Val Leu Gly Gly Val Tyr Phe Phe Phe Arg Gly	
480 485 490	
att cag tat ttc ctg cag agg cgg ccg tcg atg aag acc ctg ttt gtg	1537
Ile Gln Tyr Phe Leu Gln Arg Arg Pro Ser Met Lys Thr Leu Phe Val	
495 500 505	
gac agc tac agt gag atg ctt ttc ttt ctg cag tca ctg ttc atg ctg	1585
Asp Ser Tyr Ser Glu Met Leu Phe Phe Leu Gln Ser Leu Phe Met Leu	
510 515 520	
gcc acc gtg gtg ctg tac ttc agc cac ctc aag gag tat gtg gct tcc	1633
Ala Thr Val Val Leu Tyr Phe Ser His Leu Lys Glu Tyr Val Ala Ser	
525 530 535 540	
atg gta ttc tcc ctg gcc ttg ggc tgg acc aac atg ctc tac tac acc	1681
Met Val Phe Ser Leu Ala Leu Gly Trp Thr Asn Met Leu Tyr Tyr Thr	
545 550 555	
cgc ggt ttc cag cag atg ggc atc tat gcc gtc atg ata gag aag atg	1729
Arg Gly Phe Gln Gln Met Gly Ile Tyr Ala Val Met Ile Glu Lys Met	
560 565 570	
atc ctg aga gac ctg tgc cgt ttc atg ttt gtc tac gtc gtc ttc ttg	1777
Ile Leu Arg Asp Leu Cys Arg Phe Met Phe Val Tyr Val Val Phe Leu	
575 580 585	
ttc ggg ttt tcc aca gcg gtg gtg acg ctg att gaa gac ggg aag aat	1825
Phe Gly Phe Ser Thr Ala Val Val Thr Leu Ile Glu Asp Gly Lys Asn	
590 595 600	
gac tcc ctg ccg tct gag tcc acg tcg cac agg tgg cgg ggg cct gcc	1873
Asp Ser Leu Pro Ser Glu Ser Thr Ser His Arg Trp Arg Gly Pro Ala	
605 610 615 620	
tgc agg ccc ccc gat agc tcc tac aac agc ctg tac tcc acc tgc ctg	1921
Cys Arg Pro Pro Asp Ser Ser Tyr Asn Ser Leu Tyr Ser Thr Cys Leu	
625 630 635	
gag ctg ttc aag ttc acc atc ggc atg ggc gac ctg gag ttc act gag	1969
Glu Leu Phe Lys Phe Thr Ile Gly Met Gly Asp Leu Glu Phe Thr Glu	
640 645 650	
aac tat gac ttc aag gct gtc ttc atc atc ctg ctg ctg gcc tat gta	2017
Asn Tyr Asp Phe Lys Ala Val Phe Ile Ile Leu Leu Leu Ala Tyr Val	
655 660 665	

- 31 -

att ctc acc tac atc ctc ctg ctc aac atg ctc atc gcc ctc atg ggt 2065  
 Ile Leu Thr Tyr Ile Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly  
 670 675 680

gag act gtc aac aag atc gca cag gag agc aag aac atc tgg aag ctg 2113  
 Glu Thr Val Asn Lys Ile Ala Gln Glu Ser Lys Asn Ile Trp Lys Leu  
 685 690 695 700

cag aga gcc atc acc atc ctg gac acg gag aag agc ttc ctt aag tgc 2161  
 Gln Arg Ala Ile Thr Ile Leu Asp Thr Glu Lys Ser Phe Leu Lys Cys  
 705 710 715

atg agg aag gcc ttc cgc tca ggc aag ctg ctg cag gtg ggg tac aca 2209  
 Met Arg Lys Ala Phe Arg Ser Gly Lys Leu Leu Gln Val Gly Tyr Thr  
 720 725 730

cct gat ggc aag gac gac tac cgg tgg tgc ttc agg gtg gac gag gtg 2257  
 Pro Asp Gly Lys Asp Asp Tyr Arg Trp Cys Phe Arg Val Asp Glu Val  
 735 740 745

aac tgg acc acc tgg aac acc aac gtg ggc atc atc aac gaa gac ccg 2305  
 Asn Trp Thr Thr Trp Asn Thr Asn Val Gly Ile Ile Asn Glu Asp Pro  
 750 755 760

ggc aac tgt gag ggc gtc aag cgc acc ctg agc ttc tcc ctg cgg tca 2353  
 Gly Asn Cys Glu Gly Val Lys Arg Thr Leu Ser Phe Ser Leu Arg Ser  
 765 770 775 780

agc aga gtt tca ggc aga cac tgg aag aac ttt gcc ctg gtc ccc ctt 2401  
 Ser Arg Val Ser Gly Arg His Trp Lys Asn Phe Ala Leu Val Pro Leu  
 785 790 795

tta aga gag gca agt gct cga gat agg cag tct gct cag ccc gag gaa 2449  
 Leu Arg Glu Ala Ser Ala Arg Asp Arg Gln Ser Ala Gln Pro Glu Glu  
 800 805 810

gtt tat ctg cga cag ttt tca ggg tct ctg aag cca gag gac gct gag 2497  
 Val Tyr Leu Arg Gln Phe Ser Gly Ser Leu Lys Pro Glu Asp Ala Glu  
 815 820 825

gtc ttc aag agt cct gcc gct tcc ggg gag aag tgaaagccga attc 2544  
 Val Phe Lys Ser Pro Ala Ala Ser Gly Glu Lys  
 830 835

&lt;210&gt; 34

&lt;211&gt; 839

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

Met Lys Lys Trp Ser Ser Thr Asp Leu Gly Ala Ala Ala Asp Pro Leu  
 1 5 10 15

-32-

Gln Lys Asp Thr Cys Pro Asp Pro Leu Asp Gly Asp Pro Asn Ser Arg  
 20 25 30  
 Pro Pro Pro Ala Lys Pro Gln Leu Ser Thr Ala Lys Ser Arg Thr Arg  
 35 40 45  
 Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Phe Pro Val Asp Cys Pro  
 50 55 60  
 His Glu Glu Gly Glu Leu Asp Ser Cys Pro Thr Ile Thr Val Ser Pro  
 65 70 75 80  
 Val Ile Thr Ile Gln Arg Pro Gly Asp Gly Pro Thr Gly Ala Arg Leu  
 85 90 95  
 Leu Ser Gln Asp Ser Val Ala Ala Ser Thr Glu Lys Thr Leu Arg Leu  
 100 105 110  
 Tyr Asp Arg Arg Ser Ile Phe Glu Ala Val Ala Gln Asn Asn Cys Gln  
 115 120 125  
 Asp Leu Glu Ser Leu Leu Leu Phe Leu Gln Lys Ser Lys Lys His Leu  
 130 135 140  
 Thr Asp Asn Glu Phe Lys Asp Pro Glu Thr Gly Lys Thr Cys Leu Leu  
 145 150 155 160  
 Lys Ala Met Leu Asn Leu His Asp Gly Gln Asn Thr Thr Ile Pro Leu  
 165 170 175  
 Leu Leu Glu Ile Ala Arg Gln Thr Asp Ser Leu Lys Glu Leu Val Asn  
 180 185 190  
 Ala Ser Tyr Thr Asp Ser Tyr Tyr Lys Gly Gln Thr Ala Leu His Ile  
 195 200 205  
 Ala Ile Glu Arg Arg Asn Met Ala Leu Val Thr Leu Leu Val Glu Asn  
 210 215 220  
 Gly Ala Asp Val Gln Ala Ala Ala His Gly Asp Phe Phe Lys Lys Thr  
 225 230 235 240  
 Lys Gly Arg Pro Gly Phe Tyr Phe Gly Glu Leu Pro Leu Ser Leu Ala  
 245 250 255  
 Ala Cys Thr Asn Gln Leu Gly Ile Val Lys Phe Leu Leu Gln Asn Ser  
 260 265 270  
 Trp Gln Thr Ala Asp Ile Ser Ala Arg Asp Ser Val Gly Asn Thr Val  
 275 280 285  
 Leu His Ala Leu Val Glu Val Ala Asp Asn Thr Ala Asp Asn Thr Lys  
 290 295 300  
 Phe Val Thr Ser Met Tyr Asn Glu Ile Leu Ile Leu Gly Ala Lys Leu  
 305 310 315 320  
 His Pro Thr Leu Lys Leu Glu Glu Leu Thr Asn Lys Lys Gly Met Thr  
 325 330 335  
 Pro Leu Ala Leu Ala Ala Gly Thr Gly Lys Ile Gly Val Leu Ala Tyr  
 340 345 350  
 Ile Leu Gln Arg Glu Ile Gln Glu Pro Glu Cys Arg His Leu Ser Arg  
 355 360 365  
 Lys Phe Thr Glu Trp Ala Tyr Gly Pro Val His Ser Ser Leu Tyr Asp  
 370 375 380  
 Leu Ser Cys Ile Asp Thr Cys Glu Lys Asn Ser Val Leu Glu Val Ile  
 385 390 395 400  
 Ala Tyr Ser Ser Ser Glu Thr Pro Asn Arg His Asp Met Leu Leu Val  
 405 410 415  
 Glu Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp Asp Arg Phe Val Lys  
 420 425 430  
 Arg Ile Phe Tyr Phe Asn Phe Leu Val Tyr Cys Leu Tyr Met Ile Ile  
 435 440 445

-33-

Phe Thr Met Ala Ala Tyr Tyr Arg Pro Val Asp Gly Leu Pro Pro Phe  
 450 455 460  
 Lys Met Glu Lys Thr Gly Asp Tyr Phe Arg Val Thr Gly Glu Ile Leu  
 465 470 475 480  
 Ser Val Leu Gly Gly Val Tyr Phe Phe Phe Arg Gly Ile Gln Tyr Phe  
 485 490 495  
 Leu Gln Arg Arg Pro Ser Met Lys Thr Leu Phe Val Asp Ser Tyr Ser  
 500 505 510  
 Glu Met Leu Phe Phe Leu Gln Ser Leu Phe Met Leu Ala Thr Val Val  
 515 520 525  
 Leu Tyr Phe Ser His Leu Lys Glu Tyr Val Ala Ser Met Val Phe Ser  
 530 535 540  
 Leu Ala Leu Gly Trp Thr Asn Met Leu Tyr Tyr Thr Arg Gly Phe Gln  
 545 550 555 560  
 Gln Met Gly Ile Tyr Ala Val Met Ile Glu Lys Met Ile Leu Arg Asp  
 565 570 575  
 Leu Cys Arg Phe Met Phe Val Tyr Val Val Phe Leu Phe Gly Phe Ser  
 580 585 590  
 Thr Ala Val Val Thr Leu Ile Glu Asp Gly Lys Asn Asp Ser Leu Pro  
 595 600 605  
 Ser Glu Ser Thr Ser His Arg Trp Arg Gly Pro Ala Cys Arg Pro Pro  
 610 615 620  
 Asp Ser Ser Tyr Asn Ser Leu Tyr Ser Thr Cys Leu Glu Leu Phe Lys  
 625 630 635 640  
 Phe Thr Ile Gly Met Gly Asp Leu Glu Phe Thr Glu Asn Tyr Asp Phe  
 645 650 655  
 Lys Ala Val Phe Ile Ile Leu Leu Leu Ala Tyr Val Ile Leu Thr Tyr  
 660 665 670  
 Ile Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly Glu Thr Val Asn  
 675 680 685  
 Lys Ile Ala Gln Glu Ser Lys Asn Ile Trp Lys Leu Gln Arg Ala Ile  
 690 695 700  
 Thr Ile Leu Asp Thr Glu Lys Ser Phe Leu Lys Cys Met Arg Lys Ala  
 705 710 715 720  
 Phe Arg Ser Gly Lys Leu Leu Gln Val Gly Tyr Thr Pro Asp Gly Lys  
 725 730 735  
 Asp Asp Tyr Arg Trp Cys Phe Arg Val Asp Glu Val Asn Trp Thr Thr  
 740 745 750  
 Trp Asn Thr Asn Val Gly Ile Ile Asn Glu Asp Pro Gly Asn Cys Glu  
 755 760 765  
 Gly Val Lys Arg Thr Leu Ser Phe Ser Leu Arg Ser Ser Arg Val Ser  
 770 775 780  
 Gly Arg His Trp Lys Asn Phe Ala Leu Val Pro Leu Leu Arg Glu Ala  
 785 790 795 800  
 Ser Ala Arg Asp Arg Gln Ser Ala Gln Pro Glu Glu Val Tyr Leu Arg  
 805 810 815  
 Gln Phe Ser Gly Ser Leu Lys Pro Glu Asp Ala Glu Val Phe Lys Ser  
 820 825 830  
 Pro Ala Ala Ser Gly Glu Lys  
 835

&lt;210&gt; 35

&lt;211&gt; 2380

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

-34-

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (19)...(2313)

&lt;223&gt; Human VR2

&lt;400&gt; 35

cagcctcctc ctcctagg atg acc tca ccc tcc agc tct cca gtt ttc agg	51
Met Thr Ser Pro Ser Ser Ser Pro Val Phe Arg	
1 5 10	
ttg gag aca tta gat gga ggc caa gaa gat ggc tct gag gcg gac aga	99
Leu Glu Thr Leu Asp Gly Gly Gln Glu Asp Gly Ser Glu Ala Asp Arg	
15 20 25	
gga aag ctg gat ttt ggg agc ggg ctg cct ccc atg gag tca cag ttc	147
Gly Lys Leu Asp Phe Gly Ser Gly Leu Pro Pro Met Glu Ser Gln Phe	
30 35 40	
cag ggc gag gac cgg aaa ttc gcc cct cag ata aga gtc aac ctc aac	195
Gln Gly Glu Asp Arg Lys Phe Ala Pro Gln Ile Arg Val Asn Leu Asn	
45 50 55	
tac cga aag gga aca ggt gcc agt cag ccg gat cca aac cga ttt gac	243
Tyr Arg Lys Gly Thr Gly Ala Ser Gln Pro Asp Pro Asn Arg Phe Asp	
60 65 70 75	
cga gat cgg ctc ttc aat gcg gtc tcc cgg ggt gtc ccc gag gat ctg	291
Arg Asp Arg Leu Phe Asn Ala Val Ser Arg Gly Val Pro Glu Asp Leu	
80 85 90	
gct gga ctt cca gag tac ctg agc aag acc agc aag tac ctc acc gac	339
Ala Gly Leu Pro Glu Tyr Leu Ser Lys Thr Ser Lys Tyr Leu Thr Asp	
95 100 105	
tcg gaa tac aca gag ggc tcc aca ggt aag acg tgc ctg atg aag gct	387
Ser Glu Tyr Thr Glu Gly Ser Thr Gly Lys Thr Cys Leu Met Lys Ala	
110 115 120	
gtg ctg aac ctt aag gac gga gtc aat gcc tgc att ctg cca ctg ctg	435
Val Leu Asn Leu Lys Asp Gly Val Asn Ala Cys Ile Leu Pro Leu Leu	
125 130 135	
cag atc gac agg gac tct ggc aat cct cag ccc ctg gta aat gcc cag	483
Gln Ile Asp Arg Asp Ser Gly Asn Pro Gln Pro Leu Val Asn Ala Gln	
140 145 150 155	
tgc aca gat gac tat tac cga ggc cac agc gct ctg cac atc gcc att	531
Cys Thr Asp Asp Tyr Tyr Arg Gly His Ser Ala Leu His Ile Ala Ile	
160 165 170	
gag aag agg agt ctg cag tgt gtg aag ctc ctg gtg gag aat ggg gcc	579
Glu Lys Arg Ser Leu Gln Cys Val Lys Leu Leu Val Glu Asn Gly Ala	
175 180 185	

- 35 -

aat gtg cat gcc cgg gcc tgc ggc cgc ttc ttc cag aag ggc caa ggg Asn Val His Ala Arg Ala Cys Gly Arg Phe Phe Gln Lys Gly Gln Gly 190 195 200	627
act tgc ttt tat ttc ggt gag cta ccc ctc tct ttg gcc gct tgc acc Thr Cys Phe Tyr Phe Gly Glu Leu Pro Leu Ser Leu Ala Ala Cys Thr 205 210 215	675
aag cag tgg gat gtg gta agc tac ctc ctg gag aac cca cac cag ccc Lys Gln Trp Asp Val Val Ser Tyr Leu Leu Glu Asn Pro His Gln Pro 220 225 230 235	723
gcc agc ctg cag gcc act gac tcc cag ggc aac aca gtc ctg cat gcc Ala Ser Leu Gln Ala Thr Asp Ser Gln Gly Asn Thr Val Leu His Ala 240 245 250	771
cta gtg atg atc tcg gac aac tca gct gag aac att gca ctg gtg acc Leu Val Met Ile Ser Asp Asn Ser Ala Glu Asn Ile Ala Leu Val Thr 255 260 265	819
agc atg tat gat ggg ctc ctc caa gct ggg gcc cgc ctc tgc cct acc Ser Met Tyr Asp Gly Leu Leu Gln Ala Gly Ala Arg Leu Cys Pro Thr 270 275 280	867
gtg cag ctt gag gac atc cgc aac ctg cag gat ctc acg cct ctg aag Val Gln Leu Glu Asp Ile Arg Asn Leu Gln Asp Leu Thr Pro Leu Lys 285 290 295	915
ctg gcc gcc aag gag ggc aag atc gag att ttc agg cac atc ctg cag Leu Ala Ala Lys Glu Gly Lys Ile Glu Ile Phe Arg His Ile Leu Gln 300 305 310 315	963
cgg gag ttt tca gga ctg agc cac ctt tcc cga aag ttc acc gag tgg Arg Glu Phe Ser Gly Leu Ser His Leu Ser Arg Lys Phe Thr Glu Trp 320 325 330	1011
tgc tat ggg cct gtc cgg gtg tcg ctg tat gac ctg gct tct gtg gac Cys Tyr Gly Pro Val Arg Val Ser Leu Tyr Asp Leu Ala Ser Val Asp 335 340 345	1059
agc tgt gag gag aac tca gtg ctg gag atc att gcc ttt cat tgc aag Ser Cys Glu Glu Asn Ser Val Leu Glu Ile Ile Ala Phe His Cys Lys 350 355 360	1107
agc ccg cac cga cac cga atg gtc gtt ttg gag ccc ctg aac aaa ctg Ser Pro His Arg His Arg Met Val Val Leu Glu Pro Leu Asn Lys Leu 365 370 375	1155
ctg cag gcg aaa tgg gat ctg ctc atc ccc aag ttc ttc tta aac ttc Leu Gln Ala Lys Trp Asp Leu Leu Ile Pro Lys Phe Phe Leu Asn Phe 380 385 390 395	1203
ctg tgt aat ctg atc tac atg ttc atc ttc acc gct gtt gcc tac cat Leu Cys Asn Leu Ile Tyr Met Phe Ile Phe Thr Ala Val Ala Tyr His 400 405 410	1251



-36-

cag cct acc ctg aag aag cag gcc gcc cct cac ctg aaa gcg gag gtt Gln Pro Thr Leu Lys Lys Gln Ala Ala Pro His Leu Lys Ala Glu Val 415 420 425	1299
gga aac tcc atg ctg ctg acg ggc cac atc ctt atc ctg cta ggg ggg Gly Asn Ser Met Leu Leu Thr Gly His Ile Leu Ile Leu Leu Gly Gly 430 435 440	1347
atc tac ctc ctc gtg ggc cag ctg tgg tac ttc tgg cgg cgc cac gtg Ile Tyr Leu Leu Val Gly Gln Leu Trp Tyr Phe Trp Arg Arg His Val 445 450 455	1395
ttc atc tgg atc tcg ttc ata gac agc tac ttt gaa atc ctc ttc ctg Phe Ile Trp Ile Ser Phe Ile Asp Ser Tyr Phe Glu Ile Leu Phe Leu 460 465 470 475	1443
ttc cag gcc ctg ctc aca gtg gtg tcc cag gtg ctg tgt ttc ctg gcc Phe Gln Ala Leu Thr Val Val Ser Gln Val Leu Cys Phe Leu Ala 480 485 490	1491
atc gag tgg tac ctg ccc ctg ctt gtg tct gcg ctg gtg ctg ggc tgg Ile Glu Trp Tyr Leu Pro Leu Leu Val Ser Ala Leu Val Leu Gly Trp 495 500 505	1539
ctg aac ctg ctt tac tat aca cgt ggc ttc cag cac aca ggc atc tac Leu Asn Leu Leu Tyr Tyr Thr Arg Gly Phe Gln His Thr Gly Ile Tyr 510 515 520	1587
agt gtc atg atc cag aag gtc atc ctg cgg gac ctg ctg cgc ttc ctt Ser Val Met Ile Gln Lys Val Ile Leu Arg Asp Leu Leu Arg Phe Leu 525 530 535	1635
ctg atc tac tta gtc ttc ctt ttc ggc ttc gct gta gcc ctg gtg agc Leu Ile Tyr Leu Val Phe Leu Phe Gly Phe Ala Val Ala Leu Val Ser 540 545 550 555	1683
ctg agc cag gag gct tgg cgc ccc gaa gct cct aca ggc ccc aat gcc Leu Ser Gln Glu Ala Trp Arg Pro Glu Ala Pro Thr Gly Pro Asn Ala 560 565 570	1731
aca gag tca gtg cag ccc atg gag gga cag gag gac gag ggc aac ggg Thr Glu Ser Val Gln Pro Met Glu Gly Gln Glu Asp Glu Gly Asn Gly 575 580 585	1779
gcc cag tac agg ggt atc ctg gaa gcc tcc ttg gag ctc ttc aaa ttc Ala Gln Tyr Arg Gly Ile Leu Glu Ala Ser Leu Glu Leu Phe Lys Phe 590 595 600	1827
acc atc ggc atg ggc gag ctg gcc ttc cag gag cag ctg cac ttc cgc Thr Ile Gly Met Gly Glu Leu Ala Phe Gln Glu Gln Leu His Phe Arg 605 610 615	1875

-37-

ggc atg gtg ctg ctg ctg ctg ctg gcc tac gtg ctg ctc acc tac atc 1923  
 Gly Met Val Leu Leu Leu Leu Leu Ala Tyr Val Leu Leu Thr Tyr Ile  
 620 625 630 635

ctg ctg ctc aac atg ctc atc gcc ctc atg agc gag acc gtc aac agt 1971  
 Leu Leu Leu Asn Met Leu Ile Ala Leu Met Ser Glu Thr Val Asn Ser  
 640 645 650

gtc gcc act gac agc tgg agc atc tgg aag ctg cag aaa gcc atc tct 2019  
 Val Ala Thr Asp Ser Trp Ser Ile Trp Lys Leu Gln Lys Ala Ile Ser  
 655 660 665

gtc ctg gag atg gag aat ggc tat tgg tgg tgc agg aag aag cag cgg 2067  
 Val Leu Glu Met Glu Asn Gly Tyr Trp Trp Cys Arg Lys Lys Gln Arg  
 670 675 680

gca ggt gtg atg ctg acc gtt ggc act aag cca gat ggc agc ccc gat 2115  
 Ala Gly Val Met Leu Thr Val Gly Thr Lys Pro Asp Gly Ser Pro Asp  
 685 690 695

gag cgc tgg tgc ttc agg gtg gag gag gtg aac tgg gct tca tgg gag 2163  
 Glu Arg Trp Cys Phe Arg Val Glu Glu Val Asn Trp Ala Ser Trp Glu  
 700 705 710 715

cag acg ctg cct acg ctg tgt gag gac ccg tca ggg gca ggt gtc cct 2211  
 Gln Thr Leu Pro Thr Leu Cys Glu Asp Pro Ser Gly Ala Gly Val Pro  
 720 725 730

cga act ctc gag aac cct gtc ctg gct tcc cct ccc aag gag gat gag 2259  
 Arg Thr Leu Glu Asn Pro Val Leu Ala Ser Pro Pro Lys Glu Asp Glu  
 735 740 745

gat ggt gcc tct gag gaa aac tat gtg ccc gtc cag ctc ctc cag tcc 2307  
 Asp Gly Ala Ser Glu Glu Asn Tyr Val Pro Val Gln Leu Leu Gln Ser  
 750 755 760

aac tga tggcccagat gcagcaggag gccagaggac agagcagagg atctttccaa 2363  
 Asn \*

ccacatctgc tggctct 2380

<210> 36  
 <211> 764  
 <212> PRT  
 <213> Homo sapiens

<400> 36  
 Met Thr Ser Pro Ser Ser Ser Pro Val Phe Arg Leu Glu Thr Leu Asp  
 1 5 10 15  
 Gly Gly Gln Glu Asp Gly Ser Glu Ala Asp Arg Gly Lys Leu Asp Phe  
 20 25 30  
 Gly Ser Gly Leu Pro Pro Met Glu Ser Gln Phe Gln Gly Glu Asp Arg  
 35 40 45

-38-

Lys Phe Ala Pro Gln Ile Arg Val Asn Leu Asn Tyr Arg Lys Gly Thr  
 50 55 60  
 Gly Ala Ser Gln Pro Asp Pro Asn Arg Phe Asp Arg Asp Arg Leu Phe  
 65 70 75 80  
 Asn Ala Val Ser Arg Gly Val Pro Glu Asp Leu Ala Gly Leu Pro Glu  
 85 90 95  
 Tyr Leu Ser Lys Thr Ser Lys Tyr Leu Thr Asp Ser Glu Tyr Thr Glu  
 100 105 110  
 Gly Ser Thr Gly Lys Thr Cys Leu Met Lys Ala Val Leu Asn Leu Lys  
 115 120 125  
 Asp Gly Val Asn Ala Cys Ile Leu Pro Leu Leu Gln Ile Asp Arg Asp  
 130 135 140  
 Ser Gly Asn Pro Gln Pro Leu Val Asn Ala Gln Cys Thr Asp Asp Tyr  
 145 150 155 160  
 Tyr Arg Gly His Ser Ala Leu His Ile Ala Ile Glu Lys Arg Ser Leu  
 165 170 175  
 Gln Cys Val Lys Leu Leu Val Glu Asn Gly Ala Asn Val His Ala Arg  
 180 185 190  
 Ala Cys Gly Arg Phe Phe Gln Lys Gly Gln Gly Thr Cys Phe Tyr Phe  
 195 200 205  
 Gly Glu Leu Pro Leu Ser Leu Ala Ala Cys Thr Lys Gln Trp Asp Val  
 210 215 220  
 Val Ser Tyr Leu Leu Glu Asn Pro His Gln Pro Ala Ser Leu Gln Ala  
 225 230 235 240  
 Thr Asp Ser Gln Gly Asn Thr Val Leu His Ala Leu Val Met Ile Ser  
 245 250 255  
 Asp Asn Ser Ala Glu Asn Ile Ala Leu Val Thr Ser Met Tyr Asp Gly  
 260 265 270  
 Leu Leu Gln Ala Gly Ala Arg Leu Cys Pro Thr Val Gln Leu Glu Asp  
 275 280 285  
 Ile Arg Asn Leu Gln Asp Leu Thr Pro Leu Lys Leu Ala Ala Lys Glu  
 290 295 300  
 Gly Lys Ile Glu Ile Phe Arg His Ile Leu Gln Arg Glu Phe Ser Gly  
 305 310 315 320  
 Leu Ser His Leu Ser Arg Lys Phe Thr Glu Trp Cys Tyr Gly Pro Val  
 325 330 335  
 Arg Val Ser Leu Tyr Asp Leu Ala Ser Val Asp Ser Cys Glu Glu Asn  
 340 345 350  
 Ser Val Leu Glu Ile Ile Ala Phe His Cys Lys Ser Pro His Arg His  
 355 360 365  
 Arg Met Val Val Leu Glu Pro Leu Asn Lys Leu Leu Gln Ala Lys Trp  
 370 375 380  
 Asp Leu Leu Ile Pro Lys Phe Phe Leu Asn Phe Leu Cys Asn Leu Ile  
 385 390 395 400  
 Tyr Met Phe Ile Phe Thr Ala Val Ala Tyr His Gln Pro Thr Leu Lys  
 405 410 415  
 Lys Gln Ala Ala Pro His Leu Lys Ala Glu Val Gly Asn Ser Met Leu  
 420 425 430  
 Leu Thr Gly His Ile Leu Ile Leu Leu Gly Gly Ile Tyr Leu Leu Val  
 435 440 445  
 Gly Gln Leu Trp Tyr Phe Trp Arg Arg His Val Phe Ile Trp Ile Ser  
 450 455 460  
 Phe Ile Asp Ser Tyr Phe Glu Ile Leu Phe Leu Phe Gln Ala Leu Leu  
 465 470 475 480

-39-

Thr Val Val Ser Gln Val Leu Cys Phe Leu Ala Ile Glu Trp Tyr Leu  
 485 490 495  
 Pro Leu Leu Val Ser Ala Leu Val Leu Gly Trp Leu Asn Leu Leu Tyr  
 500 505 510  
 Tyr Thr Arg Gly Phe Gln His Thr Gly Ile Tyr Ser Val Met Ile Gln  
 515 520 525  
 Lys Val Ile Leu Arg Asp Leu Leu Arg Phe Leu Leu Ile Tyr Leu Val  
 530 535 540  
 Phe Leu Phe Gly Phe Ala Val Ala Leu Val Ser Leu Ser Gln Glu Ala  
 545 550 555 560  
 Trp Arg Pro Glu Ala Pro Thr Gly Pro Asn Ala Thr Glu Ser Val Gln  
 565 570 575  
 Pro Met Glu Gly Gln Glu Asp Glu Gly Asn Gly Ala Gln Tyr Arg Gly  
 580 585 590  
 Ile Leu Glu Ala Ser Leu Glu Leu Phe Lys Phe Thr Ile Gly Met Gly  
 595 600 605  
 Glu Leu Ala Phe Gln Glu Gln Leu His Phe Arg Gly Met Val Leu Leu  
 610 615 620  
 Leu Leu Leu Ala Tyr Val Leu Leu Thr Tyr Ile Leu Leu Leu Asn Met  
 625 630 635 640  
 Leu Ile Ala Leu Met Ser Glu Thr Val Asn Ser Val Ala Thr Asp Ser  
 645 650 655  
 Trp Ser Ile Trp Lys Leu Gln Lys Ala Ile Ser Val Leu Glu Met Glu  
 660 665 670  
 Asn Gly Tyr Trp Trp Cys Arg Lys Lys Gln Arg Ala Gly Val Met Leu  
 675 680 685  
 Thr Val Gly Thr Lys Pro Asp Gly Ser Pro Asp Glu Arg Trp Cys Phe  
 690 695 700  
 Arg Val Glu Glu Val Asn Trp Ala Ser Trp Glu Gln Thr Leu Pro Thr  
 705 710 715 720  
 Leu Cys Glu Asp Pro Ser Gly Ala Gly Val Pro Arg Thr Leu Glu Asn  
 725 730 735  
 Pro Val Leu Ala Ser Pro Pro Lys Glu Asp Glu Asp Gly Ala Ser Glu  
 740 745 750  
 Glu Asn Tyr Val Pro Val Gln Leu Leu Gln Ser Asn  
 755 760

<210> 37  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens

<400> 37  
 ggcgacctgg agttcactga g

21

<210> 38  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens

<400> 38  
 gagcaggagg atgttaggtga g

21

<210> 39  
 <211> 226

- 40 -

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 39

ctgcagcttc	cagatgttct	tgctctcctg	tgcgatcttg	ttgacagtct	cacccatgag	60
ggcgatgagc	atgttgagca	ggaggatgta	ggtgagaatt	acataggcca	gcagcaggat	120
gatgaagaca	gccttgaagt	catagtcttc	agtgaactcc	aggtcgcca	tgccgatggt	180
gaacttgaac	agctccaggc	agggtggagta	caggctgttg	taggag		226

&lt;210&gt; 40

&lt;211&gt; 226

&lt;212&gt; DNA

&lt;213&gt; R. rattus

&lt;400&gt; 40

gagaatgttg	tccgacataa	ggtgtacaga	cctcgacaag	ttcaagtggg	agccgtaccc	60
gctggacctc	aagtgactct	tgatgctgaa	gttccgacag	aagtagtagg	acaatgaccg	120
gatacactaa	gagtggatgt	aggaagacga	gttgtacgag	taacgagagt	acccactctg	180
gcagttgttc	taacgtgttc	tctcgttctt	gtagaccttc	gacgtc		226

&lt;210&gt; 41

&lt;211&gt; 145

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 41

aagacctcag	cgctcctctg	cttcagagac	cctgnaaaac	tgctgcagat	aaacttcttc	60
gggctgagca	nactgcctat	ctcgagcact	tgctctcttt	aaaaggggga	ccagggcaaa	120
gttcttccag	tgtctgcttg	aaact				145

&lt;210&gt; 42

&lt;211&gt; 144

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 42

tcaaagtcgg	tctgtgacct	tcttgaaacg	ggaccagggg	gaaaattctc	tccgttcang	60
agctntatcc	gtcagacgag	tccgggtcct	tcaaataagac	gctgtcaaaa	gtcccagaga	120
tttcgggtctc	ctgcgactcc	agaa				144

&lt;210&gt; 43

&lt;211&gt; 219

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 43

cccactccaa	aaggacacct	gcccagaccc	cctggatgga	gaccctaact	ccaggccacc	60
tccagccaag	ccccagctct	ccacggccaa	gagccgcacc	cggctctttg	ggaaggggtga	120
ctcggaggag	gctttcccg	tggattgccc	tcacgaggaa	ggtgagctgg	actcctgccc	180
gaccatcaca	gtcagccctg	ttatcaccat	ccagaggcc			219

&lt;210&gt; 44

&lt;211&gt; 219

&lt;212&gt; DNA

&lt;213&gt; R. rattus

-41-

<400> 44  
 cccaccccaa gagaactcct gcctggaccc tccagacaga gaccctaact gcaagccacc 60  
 tccagtcaag cccacatct tcaactaccag gagtcgtacc cggctttttg ggaaggggtga 120  
 ctcggaggag gcctctcccc tggactgccc ttatgaggaa ggcgggctgg cttctgccc 180  
 tatcatcact gtcagctctg ttctaactat ccagaggcc 219

<210> 45  
 <211> 91  
 <212> PRT  
 <213> Homo sapiens

<400> 45  
 Met Lys Lys Trp Ser Ser Thr Asp Leu Gly Ala Ala Ala Asp Pro Leu  
 1 5 10 15  
 Gln Lys Asp Thr Cys Pro Asp Pro Leu Asp Gly Asp Pro Asn Ser Arg  
 20 25 30  
 Pro Pro Pro Ala Lys Pro Gln Leu Ser Thr Ala Lys Ser Arg Thr Arg  
 35 40 45  
 Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Phe Pro Val Asp Cys Pro  
 50 55 60  
 His Glu Glu Gly Glu Leu Asp Ser Cys Pro Thr Ile Thr Val Ser Pro  
 65 70 75 80  
 Val Ile Thr Ile Gln Arg Pro Arg Xaa Arg Pro  
 85 90

<210> 46  
 <211> 91  
 <212> PRT  
 <213> R. rattus

<400> 46  
 Met Glu Gln Arg Ala Ser Leu Asp Ser Glu Glu Ser Glu Ser Pro Pro  
 1 5 10 15  
 Gln Glu Asn Ser Cys Leu Asp Pro Pro Asp Arg Asp Pro Asn Cys Lys  
 20 25 30  
 Pro Pro Pro Val Lys Pro His Ile Phe Thr Thr Arg Ser Arg Thr Arg  
 35 40 45  
 Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Ser Pro Leu Asp Cys Pro  
 50 55 60  
 Tyr Glu Glu Gly Gly Leu Ala Ser Cys Pro Ile Ile Thr Val Ser Ser  
 65 70 75 80  
 Val Leu Thr Ile Gln Arg Pro Gly Asp Gly Pro  
 85 90

<210> 47  
 <211> 24  
 <212> DNA  
 <213> Homo sapiens

<400> 47  
 agaaatggag cagcacagac ttgg 24

<210> 48  
 <211> 23  
 <212> DNA

-42-

<213> Homo sapiens

<400> 48

tcacttctcc ccggaagcgg cag

23

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/01418

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 530/350; 435/69.1, 252.3, 254.11, 320.1, 325; 514/2

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/350; 435/69.1, 252.3, 254.11, 320.1, 325; 514/2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
None

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, Dialog, Medline, WPI

search terms: Capsaicin Receptor, Vanilloid Receptor, cloning, DNA, isolation, purification

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	CATERINA et al. The Capsaicin Receptor: A Heat-Activated Ion Channel in the Pain Pathway. Nature. 23 October 1997, Vol. 389, pages 816-824, especially pages 817-819.	1, 2, 4(Species 1), 5, 6, 8(Species 1), 9-11, 17 ----- 4 (Species 14), 8 (Species 14)
X	SZALLASI et al. Characterization of a Peripheral Vanilloid (Capsaicin) Receptor in the Urinary Bladder of the Rat. Life Sciences. 1993, Vol. 52, No. 20, pages 221-226, especially pages 223, 224.	1, 2, 4(species 1), 17



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

08 APRIL 1999

Date of mailing of the international search report

12 MAY 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Sally P. Teng

Telephone No. (703) 308-0196



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/01418

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WARDLE et al. Pharmacological Characterization of the Vanilloid Receptor in the Rat Isolated Vas Deferens. J. Pharm. Pharmacol. 1996, Vol. 48, pages 285-291, especially pages 285-287.	1, 2, 4(species 1), 17
A	US 5,021,450 A (BLUMBERG) 04 June 1991, columns 7 and 9.	1-11, 17

Form PCT/ISA/210 (continuation of second sheet)(July 1992)★

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/01418

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:  
1-11, 17, species 1 and 14
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/01418

## A. CLASSIFICATION OF SUBJECT MATTER: IPC (6):

C07K 14/435, 14/705; C12N 15/12, 15/63, 15/70, 15/74, 15/79, 15/81; A61K 38/17

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-11 and 17, drawn to Capsaicin receptor and its encoding polynucleotide.

Group II, claim 12, drawn to antibody against the capsaicin receptor.

Group III, claims 13-16, drawn to method for identifying compounds that bind a capsaicin receptor.

Group IV, claim 18, drawn to a non-human transgenic animal.

Group V, claim 19, drawn to a method of screening for biologically active agents that modulate capsaicin receptor function.

The inventions listed as Groups I-V do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The special technical feature of Group I is the amino acid sequence and its encoding nucleic acid sequence. The special technical feature of Group II is the antibody that binds the receptor of Group I but does not share the same primary structure as the receptor of Group I. The special technical feature of Group III is a method of identifying a compound that bind the receptor of Group I. The special technical feature of Group IV is a transgenic animal having altered capsaicin receptor function. The special technical feature of Group V is a method of screening for biologically active agents that modulate capsaicin receptor function. The special technical feature of each Group is not the same or does not correspond to the special technical feature of any other group. The products of Groups I, II, and IV are structurally and functionally distinct, and the methods of Groups III and V require different method steps and reagents for achieving different goals. The Groups are not linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

- 1) SEQ ID NO: 1 and 2
- 2) SEQ ID NO: 3 and 4
- 3) SEQ ID NO: 5
- 4) SEQ ID NO: 6
- 5) SEQ ID NO: 7
- 6) SEQ ID NO: 8 and 9
- 7) SEQ ID NO: 10 and 11
- 8) SEQ ID NO: 20
- 9) SEQ ID NO: 21
- 10) SEQ ID NO: 22 and 23
- 11) SEQ ID NO: 24 and 25
- 12) SEQ ID NO: 26
- 13) SEQ ID NO: 27
- 14) SEQ ID NO: 33 and 34
- 15) SEQ ID NO: 35 and 36

The claims are deemed to correspond to the species listed above in the following manner:

claim 4: SEQ ID NO: 2, 4, 9, 11, 23, 25, 27, 34, and 36

claim 8: SEQ ID NO: 1, 3, 5, 6, 7, 8, 10, 20, 21, 22, 24, 26, 33, and 35

The following claims are generic: 1-19

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: The special technical

**INTERNATIONAL SEARCH REPORT**

International application No.  
PCT/US99/01418

feature of each species is its amino acid sequence and/or its encoding nucleic acid sequence. Each of the species is a capsaicin receptor comprising a unique amino acid sequence which is encoded by a unique nucleic acid sequence. The species are structurally and functionally distinct capsaicin receptors or they encode structurally and functionally distinct capsaicin receptors.